


# NATURAL PRODUCT COMMUNICATIONS

An International Journal for Communications and Reviews Covering all  
Aspects of Natural Products Research



Volume 10. Issue 10. Pages 1641-1796. 2015  
ISSN 1934-578X (printed); ISSN 1555-9475 (online)  
[www.naturalproduct.us](http://www.naturalproduct.us)

## EDITOR-IN-CHIEF

**DR. PAWAN K AGRAWAL**

Natural Product Inc.  
7963, Anderson Park Lane,  
Westerville, Ohio 43081, USA  
agrawal@naturalproduct.us

## EDITORS

**PROFESSOR ALEJANDRO F. BARRERO**

Department of Organic Chemistry,  
University of Granada,  
Campus de Fuente Nueva, s/n, 18071, Granada, Spain  
afbarre@ugr.es

**PROFESSOR ALESSANDRA BRACA**

Dipartimento di Chimica Bioorganica e Biofarmacia,  
Università di Pisa,  
via Bonanno 33, 56126 Pisa, Italy  
braca@farm.unipi.it

**PROFESSOR DE-AN GUO**

State Key Laboratory of Natural and Biomimetic Drugs,  
School of Pharmaceutical Sciences,  
Peking University,  
Beijing 100083, China  
gda5958@163.com

**PROFESSOR YOSHIHIRO MIMAKI**

School of Pharmacy,  
Tokyo University of Pharmacy and Life Sciences,  
Horinouchi 1432-1, Hachioji, Tokyo 192-0392, Japan  
mimaki@ps.toyaku.ac.jp

**PROFESSOR STEPHEN G. PYNE**

Department of Chemistry  
University of Wollongong  
Wollongong, New South Wales, 2522, Australia  
spyne@uow.edu.au

**PROFESSOR MANFRED G. REINECKE**

Department of Chemistry,  
Texas Christian University,  
Forts Worth, TX 76129, USA  
m.reinecke@tcu.edu

**PROFESSOR WILLIAM N. SETZER**

Department of Chemistry  
The University of Alabama in Huntsville  
Huntsville, AL 35809, USA  
wsetzer@chemistry.uah.edu

**PROFESSOR YASUHIRO TEZUKA**

Faculty of Pharmaceutical Sciences  
Hokuriku University  
Ho-3 Kanagawa-machi, Kanazawa 920-1181, Japan  
y-tezuka@hokuriku-u.ac.jp

**PROFESSOR DAVID E. THURSTON**

Institute of Pharmaceutical Science  
Faculty of Life Sciences & Medicine  
King's College London, Britannia House  
7 Trinity Street, London SE1 1DB, UK  
david.thurston@kcl.ac.uk

## HONORARY EDITOR

**PROFESSOR GERALD BLUNDEN**

The School of Pharmacy & Biomedical Sciences,  
University of Portsmouth,  
Portsmouth, PO1 2DT U.K.  
axuf64@dsl.pipex.com

## ADVISORY BOARD

Prof. Viqar Uddin Ahmad  
Karachi, Pakistan

Prof. Giovanni Appendino  
Novara, Italy

Prof. Yoshinori Asakawa  
Tokushima, Japan

Prof. Roberto G. S. Berlinck  
São Carlos, Brazil

Prof. Anna R. Bilia  
Florence, Italy

Prof. Maurizio Bruno  
Palermo, Italy

Prof. César A. N. Catalán  
Tucumán, Argentina

Prof. Josep Coll  
Barcelona, Spain

Prof. Geoffrey Cordell  
Chicago, IL, USA

Prof. Fatih Demirci  
Eskişehir, Turkey

Prof. Ana Cristina Figueiredo  
Lisbon, Portugal

Prof. Cristina Gracia-Viguera  
Murcia, Spain

Dr. Christopher Gray  
Saint John, NB, Canada

Prof. Dominique Guillaume  
Reims, France

Prof. Duvvuru Gunasekar  
Tirupati, India

Prof. Hisahiro Hagiwara  
Niigata, Japan

Prof. Tsukasa Iwashina  
Tsukuba, Japan

Prof. Leopold Jirovetz  
Vienna, Austria

Prof. Vladimir I Kalinin  
Vladivostok, Russia

Prof. Phan Van Kiem  
Hanoi, Vietnam

Prof. Niel A. Koorbanally  
Durban, South Africa

Prof. Chiaki Kuroda  
Tokyo, Japan

Prof. Hartmut Laatsch  
Göttingen, Germany

Prof. Marie Lacaille-Dubois  
Dijon, France

Prof. Shoen-Sheng Lee  
Taipei, Taiwan

Prof. Imre Mathe  
Szeged, Hungary

Prof. M. Soledade C. Pedras  
Saskatoon, Canada

Prof. Luc Pieters  
Antwerp, Belgium

Prof. Peter Proksch  
Düsseldorf, Germany

Prof. Phila Raharivelomanana  
Tahiti, French Polynesia

Prof. Luca Rastrelli  
Fisciano, Italy

Prof. Stefano Serra  
Milano, Italy

Prof. Monique Simmonds  
Richmond, UK

Dr. Bikram Singh  
Palampur, India

Prof. John L. Sorensen  
Manitoba, Canada

Prof. Johannes van Staden  
Scottsville, South Africa

Prof. Valentin Stonik  
Vladivostok, Russia

Prof. Winston F. Tinto  
Barbados, West Indies

Prof. Sylvia Urban  
Melbourne, Australia

Prof. Karen Valant-Vetschera  
Vienna, Austria

## INFORMATION FOR AUTHORS

Full details of how to submit a manuscript for publication in Natural Product Communications are given in Information for Authors on our Web site <http://www.naturalproduct.us>.

Authors may reproduce/republish portions of their published contribution without seeking permission from NPC, provided that any such republication is accompanied by an acknowledgment (original citation)-Reproduced by permission of Natural Product Communications. Any unauthorized reproduction, transmission or storage may result in either civil or criminal liability.

The publication of each of the articles contained herein is protected by copyright. Except as allowed under national "fair use" laws, copying is not permitted by any means or for any purpose, such as for distribution to any third party (whether by sale, loan, gift, or otherwise); as agent (express or implied) of any third party; for purposes of advertising or promotion; or to create collective or derivative works. Such permission requests, or other inquiries, should be addressed to the Natural Product Inc. (NPI). A photocopy license is available from the NPI for institutional subscribers that need to make multiple copies of single articles for internal study or research purposes.

**To Subscribe:** Natural Product Communications is a journal published monthly. 2015 subscription price: US\$2,595 (Print, ISSN# 1934-578X); US\$2,595 (Web edition, ISSN# 1555-9475); US\$2,995 (Print + single site online); US\$595 (Personal online). Orders should be addressed to Subscription Department, Natural Product Communications, Natural Product Inc., 7963 Anderson Park Lane, Westerville, Ohio 43081, USA. Subscriptions are renewed on an annual basis. Claims for nonreceipt of issues will be honored if made within three months of publication of the issue. All issues are dispatched by airmail throughout the world, excluding the USA and Canada.

## Effect of Supplementation with Wheat Bran Aqueous Extracts Obtained by Ultrasound-Assisted Technologies on the Sensory Properties and the Antioxidant Activity of Dry Pasta

Antonella Pasqualone<sup>a,\*</sup>, Laura Nunzia Delvecchio<sup>a</sup>, Giuseppe Gambacorta<sup>a</sup>, Barbara Laddomada<sup>b</sup>, Valeria Urso<sup>c</sup>, Agata Mazzaglia<sup>d</sup>, Paolo Ruisi<sup>c</sup> and Giuseppe Di Miceli<sup>c,e</sup>

<sup>a</sup>Dipartimento di Scienze del Suolo, della Pianta e degli Alimenti, Università degli Studi di Bari, via Amendola, 165/a, 70126 Bari, Italy

<sup>b</sup>Istituto di Scienze delle Produzioni Alimentari, C.N.R., via Monteroni, 73100 Lecce, Italy

<sup>c</sup>Fondazione Angelo e Salvatore Lima Mancuso, Piazza Marina, 61, 90100 Palermo, Italy

<sup>d</sup>Dipartimento di Agricoltura, Alimentazione e Ambiente, Università degli Studi di Catania, via S. Sofia, 98, 95123 Catania, Italy

<sup>e</sup>Dipartimento di Scienze Agrarie e Forestali, Università degli Studi di Palermo, viale delle Scienze, 90128 Palermo, Italy

antonella.pasqualone@uniba.it

Received: May 6<sup>th</sup>, 2015; Accepted: August 3<sup>rd</sup>, 2015

Phenolic compounds have antioxidant properties and activate endogenous detoxification defense systems able to scavenge the reactive species of oxygen. The external layers of wheat caryopsis, largely constituting by-products of the milling industry such as bran and various middlings, contain relevant amounts of phenolic compounds. The aim of the research has been to evaluate the effect of supplementation with wheat bran aqueous extracts, obtained by ultrasound-assisted technologies, on the sensory properties and antioxidant activity of dry pasta. The HPLC-DAD characterization of the extract evidenced the presence of ferulic and *p*-coumaric acids. The supplemented pasta showed significantly higher antioxidant activity and phenolic content than the control, coupled to good overall sensory judgment. In addition, two different pasta drying diagrams were adopted, and the comparison of the corresponding end-products allowed it to be pointed out that the processing technology has to be carefully set up to prevent possible detrimental effects on the antioxidant activity. The proposed utilization of bran might add value to a milling by-product that, otherwise, is mostly employed in animal feeding.

**Keywords:** Phenolic extracts, Ultrasound, Pasta, Bran, Functional foods.

In recent years the role played by reactive oxygen species in many diseases related to aging has been extensively reported [1]. Functional foods, providing health benefits beyond basic nutrition [2], are the object of increasing interest by both producers and health-conscious consumers. In particular, redox and antioxidant systems are among the most addressed targets in functional food science.

Phenolic compounds have been demonstrated to have clear antioxidant properties *in vitro* and *in vivo* [3]. The consumption of phenolics may hence contribute in preventing diseases related to aging [4,5]. The ascertained toxicity of synthetic phenolic antioxidants, such as butylated hydroxy anisole and butylated hydroxy toluene [6,7], focused attention towards the phenolic compounds commonly found in natural sources such as fruits, vegetables, and cereals [8]. In particular, the external layers of wheat caryopsis, largely constituting by-products of the milling industry such as bran and various middlings, contain relevant amounts of phenolic compounds [9-11]. These by-products are commonly destined for animal feeding. Alternatively, they are used in whole wheat bread- and pasta-making to an increasing, but still marginal, extent. However, the textural features of the end-product, such as pasta firmness and elasticity, are negatively affected by fiber [12,13], and an increase of cooking loss, swelling index, and water absorption has been reported [14].

The enrichment of fresh pasta with bran phenolic extracts, after preliminary KOH-induced hydrolysis, affects dough consistency

and pasta taste [15]. Ultrasound-assisted technology, an environmentally-friendly system for extracting bioactive compounds from natural sources [16], could be exploited to recover phenolic compounds from bran. The aim of this research has been to evaluate the effect of pasta supplementation with wheat bran aqueous extract, obtained by ultrasound-assisted technologies, on the sensory properties and antioxidant activity of the end-product.

The bran aqueous extract was submitted to quali-quantitative characterization by HPLC-DAD (Table 1). The most represented compound was ferulic acid, as already observed by other authors [17,18], followed by *p*-coumaric acid. None of the other phenolic acids usually found in wheat was detected. Higher levels of phenolics have been reported in the literature for extractions carried out by acid or basic hydrolysis of bran followed, after neutralization, by methanol or ethyl acetate extraction [17,19], but these methods are not suitable for food supplementation purposes.

**Table 1:** HPLC-DAD quali-quantitative characterization of aqueous extract obtained from wheat bran by means of ultrasound-assisted technologies.

Phenolic acid	Content(mg g <sup>-1</sup> d.m.)
<i>p</i> -Hydroxy benzoic acid	n.d.
Vanillic acid	n.d.
Syringic acid	n.d.
Sinapic acid	n.d.
Ferulic acid	0.31±0.01
<i>p</i> -Coumaric acid	0.25±0.01
<i>o</i> -Coumaric acid	n.d.

n.d. = not detected.

The semolina used in the trials showed good pasta-making features (Table 2). The gluten index was in the highest strength category ( $>80$ ) [20], ensuring good pasta consistency and resistance to overcooking [21]. The yellow index was above the ranges previously observed in quality surveys of Southern Italian semolina [22-24]. The bright yellow color, due to carotenoid pigments, is typical of semolina and is partly transferred to the end-product. The ash content fell within the limits of the current rules [25].

**Table 2:** Main quality characteristics of semolina used in the pasta-making trials.

Parameter	Value
Ash (% d.m.)	0.83±0.01
Dry gluten (% d.m.)	12.7±0.2
Gluten index	91±3
Yellow index	26.3±1.0

Semolina was processed with either aqueous bran extract (supplemented pasta) or water (control) into 'mezzi rigatoni'-shaped pasta, and dried according to two different drying diagrams (Fig. 1). The content of phenolic substances of the supplemented pasta was in the range reported for whole meal pasta [26] and was significantly higher in both the supplemented types than in the corresponding controls. The antioxidant activity of supplemented pasta was similar to the levels reported in soft wheat whole meal or in pigmented wheat [27,28], and was higher than the control only when the HT1 drying diagram was adopted. This result was probably due to the more severe thermal effect of the HT2 than HT1 process.

**Table 3:** Content of phenolic compounds and antioxidant activity of pasta supplemented with aqueous bran extract and of control pasta. Both pasta types were dried according to two different drying programs, HT1 and HT2, reported in Fig. 1.


Pasta type	Antioxidant activity		Total soluble phenolic compounds
	$\mu\text{mol Trolox g}^{-1}$ (d.m.)	DPPH SC%	mg FAE g <sup>-1</sup> (d.m.)
Control-HT1	0.42±0.003 <sup>b</sup>	10.62±0.42 <sup>b</sup>	1.03±0.01 <sup>b</sup>
Supplemented-HT1	0.49±0.002 <sup>a</sup>	12.77±0.13 <sup>a</sup>	1.16±0.01 <sup>a</sup>
Control-HT2	0.41±0.004 <sup>b</sup>	10.17±0.64 <sup>b</sup>	1.01±0.05 <sup>b</sup>
Supplemented-HT2	0.45±0.002 <sup>ab</sup>	11.56±0.23 <sup>ab</sup>	1.11±0.03 <sup>a</sup>

Different letters in the same column indicate significant differences at  $p < 0.05$ ; DPPH SC% = 2,2-diphenyl-1-picrylhydrazyl scavenging capacity percentage; Trolox = 6-hydroxy-2,5,7,8-tetramethylchroman-2-carboxylic acid.

Good cooking quality performances of pasta include low stickiness, and high firmness and elasticity. These features can be evaluated either by a texture analyzer or by sensory evaluation, the latter approach being more similar to the final consumer evaluation than the former. The sensory analysis of supplemented and control pasta was performed to evaluate 15 descriptors, defined in Table 4, so as to express an overall judgment. The addition of bran extract did not cause significant detrimental effects on textural characteristics, such as stickiness, elasticity, chewiness, and resistance to cutting (Table 5). Moreover, the supplemented pasta interestingly showed higher scores for the firmness sensory descriptor than control pasta. In addition, pasta color did not show significant differences between supplemented and control trials.

As far as taste was concerned, sweet and salty descriptors did not show differences by comparing supplemented pasta with the corresponding controls, whereas a slightly greater astringent note was perceived in pasta to which the bran extracts had been added, but without significant differences with the control. Regarding odor and flavor descriptors, no significant differences were observed. As a consequence, the overall evaluation was similar for supplemented pasta and control.

These findings demonstrated the effectiveness of the strategy of transferring phenolic compounds from bran to pasta by means of an aqueous extract, without encountering the typical drawbacks of



**Figure 1:** Variations of temperature and relative humidity in the drier during HT1 (top) and HT2 (bottom) pasta drying programs.

**Table 4:** Descriptors selected to describe the sensory properties of pasta.

Descriptor	Definition
Color	Intensity of amber-yellow color
Stickiness	Degree of adherence of pasta to the teeth (while chewing) and to the palate (by pressing a pasta piece onto palate, then removing it by the tongue)
Elasticity	Degree of recovery evaluated by stretching (spaghetti) or bending (other types) a pasta piece
Resistance to cutting	Resistance of pasta to being cut by a fork
Firmness	Degree of resistance while compressing a pasta piece by fingers
Odor of semolina	Aroma characteristic of durum wheat semolina perceived by orthonasal olfaction
Odor of cooked pasta	Aroma characteristic of cooked pasta perceived by orthonasal olfaction
Off-odor	Anomalous odor perceived by orthonasal olfaction
Flavor of semolina	Retronasal aroma characteristic of semolina perceived when pasta is in the mouth
Flavor of cooked pasta	Retronasal aroma characteristic of cooked pasta perceived when pasta is in the mouth
Off-flavor	Retronasal anomalous aroma perceived when pasta is in the mouth
Sweet taste	Taste of sucrose perceived in the mouth while chewing
Salty taste	Taste of sodium chloride perceived in the mouth while chewing
Astringency	Sensation of dryness and coarseness perceived in the mouth while chewing
Chewiness	Duration and number of chews needed before swallowing pasta
Overall judgment	Overall evaluation of pasta acceptability

whole meal pasta-making due to fiber interference with the gluten network [14]. The supplementation of pasta with aqueous phenolic extracts obtained from durum wheat bran by ultrasound-assisted technologies allowed us to obtain pasta with higher antioxidant activity than the control, coupled to good sensory properties. In addition, the proposed utilization of bran might valorize a milling by-product that otherwise is essentially employed in animal feeding.

It has to be remarked that the differences in the levels of antioxidant activity and phenolic compounds between supplemented pasta and control, although being statistically significant, were modest. However, these results can be considered satisfactory because the sensory properties of the end-product were not altered. Moreover, the typically high frequency of pasta consumption in the Italian diet could ensure beneficial health effects even for slight augmentations of pasta antioxidant activity. Finally, the comparison of two different drying diagrams allowed it to be pointed out that the processing technology has to be carefully set up to prevent possible detrimental effects on the antioxidant activity of pasta.

## Experimental

**Chemicals and reagents:** Authentic standards of phenolic acids (*p*-hydroxybenzoic acid, vanillic acid, syringic acid, *p*-coumaric acid,

*o*-coumaric acid, sinapic acid, and ferulic acid), 2,2-diphenyl-1-picrylhydrazyl (DPPH), and 6-hydroxy-2,5,7,8-tetramethylchroman-2-carboxylic acid (Trolox) were obtained from Sigma-Aldrich Chemical Co. (St. Louis, MO, USA), and Folin-Ciocalteu reagent, acetonitrile, and phosphoric acid from Carlo Erba Reagent (Milan, Italy).

**Ultrasound-assisted production of bran aqueous extract:** Durum wheat bran (3.5 kg) was mixed with tap water (35 L) and the suspension was submitted to ultrasound-assisted extraction at 20°C for 25 min in a pilot plant assembled by Weal (Milano, Italy) whose working principle and details are reported in Punzi *et al.* [29], then the suspension was filtered through a metal grid with 1 mm holes to recover the liquid phase. About 25 L of aqueous extract was obtained that was used immediately after its production.

**Table 5:** Sensory properties of pasta supplemented with aqueous bran extract and of control pasta. Both pasta types were dried according to two different drying programs, HT1 and HT2, as reported in Fig. 1.

Sensory descriptor	Pasta type			
	Control-HT1	Supplemented-HT1	Control-HT2	Supplemented-HT2
Color	4.75±1.00 <sup>a</sup>	4.66±1.29 <sup>a</sup>	4.50±1.27 <sup>a</sup>	3.83±1.92 <sup>a</sup>
Stickiness	5.00±2.33 <sup>a</sup>	5.33±1.21 <sup>a</sup>	4.83±1.60 <sup>a</sup>	4.75±2.23 <sup>a</sup>
Elasticity	6.25±1.73 <sup>a</sup>	6.83±1.48 <sup>a</sup>	5.91±1.50 <sup>a</sup>	6.33±1.03 <sup>a</sup>
Resistance to cutting	6.00±1.34 <sup>a</sup>	6.66±1.95 <sup>a</sup>	6.83±1.29 <sup>a</sup>	6.75±1.78 <sup>a</sup>
Firmness	4.50±1.75 <sup>a</sup>	6.08±1.78 <sup>b</sup>	3.83±1.95 <sup>a</sup>	6.00±2.11 <sup>b</sup>
Semolina odor	6.58±1.50 <sup>a</sup>	6.66±1.44 <sup>a</sup>	6.33±1.31 <sup>a</sup>	6.41±1.50 <sup>a</sup>
Cooked pasta odor	5.75±1.88 <sup>a</sup>	6.00±2.22 <sup>a</sup>	5.08±1.29 <sup>a</sup>	6.00±2.00 <sup>a</sup>
Off-odor	2.08±1.19 <sup>a</sup>	2.00±1.98 <sup>a</sup>	1.83±1.30 <sup>a</sup>	1.66±1.95 <sup>a</sup>
Sweet	4.50±1.24 <sup>a</sup>	4.33±2.28 <sup>a</sup>	4.91±1.56 <sup>a</sup>	4.91±1.97 <sup>a</sup>
Salty	2.08±1.40 <sup>a</sup>	2.75±1.73 <sup>a</sup>	2.16±1.22 <sup>a</sup>	2.25±2.30 <sup>a</sup>
Astringency	3.41±1.82 <sup>a</sup>	4.25±2.02 <sup>a</sup>	3.25±2.11 <sup>a</sup>	3.91±2.90 <sup>a</sup>
Chewiness	5.91±2.11 <sup>a</sup>	5.25±1.66 <sup>a</sup>	5.33±2.07 <sup>a</sup>	5.25±2.00 <sup>a</sup>
Semolina flavor	6.25±1.56 <sup>a</sup>	5.33±1.76 <sup>a</sup>	6.41±1.44 <sup>a</sup>	5.66±1.97 <sup>a</sup>
Cooked pasta flavor	4.50±1.71 <sup>a</sup>	4.33±1.23 <sup>a</sup>	5.00±1.68 <sup>a</sup>	5.08±1.50 <sup>a</sup>
Off-flavor	2.16±1.86 <sup>ab</sup>	3.41±1.53 <sup>b</sup>	1.75±2.21 <sup>a</sup>	2.16±2.39 <sup>ab</sup>
Overall judgment	5.50±1.09 <sup>a</sup>	4.58±1.17 <sup>a</sup>	5.50±1.51 <sup>a</sup>	5.41±1.44 <sup>a</sup>

Different letters in the same row indicate significant differences at  $p < 0.05$ .

**Pasta production:** Durum wheat (*Triticum durum* Desf.) cv ‘Vertola’ was used to perform the experiments. Grains were milled to semolina by means of a MLU 202 mill (Buhler, Uzwil, Switzerland), after conditioning at 17.5% moisture. Semolina was added to either the aqueous bran extract at 30% level (on semolina weight basis) to produce supplemented pasta, or 30% water, for control pasta. Specifically, about 5.5 kg semolina was kneaded with either water or aqueous bran extract for 15 min by using a MAC 60 VR 1-bar vacuum extruder (Italpast, Fidenza, Italy). The dough, extruded through a Teflon die, was shaped into ‘mezzi rigatoni’ having a length of 18.60±1.20 mm, diameter 6.66±0.16 mm, and thickness 1.00±0.08 mm. Then, each pasta type was divided into 2 batches and dried in a static dryer (LAB, Namad Impianti, Rome, Italy) according to 2 diverse high temperature (HT) drying programs (coded HT1 and HT2; reported in Fig. 1), both reaching the maximum temperature of 78°C but involving different variations of relative humidity (RH).

**HPLC quali-quantitative analysis of bran aqueous extract:** Quantitative measurements of individual phenolic acids were achieved using an Agilent 1100 (Santa Clara, CA, USA) liquid chromatograph equipped with a photodiode array detector (DAD). Wavelengths of 280, 295 and 320 nm were used for quantification of the phenolic acids. Phenolic acid separation was achieved by using a Phenomenex-luna (Macclesfield, UK) 5 µm C18 100 Å column (250 × 4.6 mm) and a column temperature of 30°C. The flow rate of the mobile phase was 1.0 mL/min, and the injection volume was 20 µL. A gradient-elution program was utilized with a mobile phase consisting of acetonitrile (solution A) and 1%, v/v, phosphoric acid in water (solution B) as follows: isocratic elution,

100% B, 0-30 min; linear gradient from 100% B to 85% B, 30-55 min; linear gradient from 85% B to 50% B, 55-80 min; linear gradient from 50% B to 30% B, 80-82 min; post time, 10 min before the next injection. The flow rate of the mobile phase was 1.0 mL/min, and the injection volume was 20 µL. Phenolic acids were quantified using calibration curves of phenolic acid standards (previously prepared as stock solutions at 2 mg mL<sup>-1</sup> in 80:20, v/v, ethanol/water and stored in the dark at -18°C, where it remained stable for over 3 months). The analyses were carried out in triplicate.

**Basic chemical and physical determinations of semolina and pasta:** Semolina was submitted to the determination of ash content according to the AACC method 08-12 [30]. Dry gluten and gluten index were determined as in the AACC method 38-12.02 [30]. Yellow index (corresponding to *b*\*) was determined by means of the reflectance colorimeter Chroma Meter CR-300 (Konica Minolta Sensing, Osaka, Japan).

**Quantification of total phenolic compounds and antioxidant activity of pasta:** The total phenolic compounds (specifically the fraction composed of soluble free phenolic acids and soluble phenolics bound to low molecular mass components) and the antioxidant activity, assessed by the DPPH radical scavenging capacity assay, were determined as in Pasqualone *et al.* [31]. All tests were carried out in triplicate.

**Sensory analysis:** Pasta was cooked in boiling distilled water at 1:10, w/v, pasta to water ratio, without the addition of salt. Optimum cooking time was determined according to the AACC Method 16-50 [30]. After cooking and draining, the samples were allowed to rest for 5 min. Subsequent analysis was carried out according to the UNI method 10957 [32], by a trained panel of 12 assessors, previously selected over a group of 16 people for their reliability, consistency, and discriminating ability. After a cycle of lectures on the basic notions of sensory analysis, the selection involved a series of tests to monitor the performance in recognizing the basic tastes, odors, flavors, and colors. Prior to actual sample evaluation, the selected panelists were trained via six pre-test sessions held to (i) familiarize them with the vocabulary regarding the descriptors of pasta, (ii) define the list of sensory terms to be evaluated in the samples studied, (iii) establish the order of their testing, (iv) define their intensity range, and (v) fix their scale anchors. Samples similar to those objects of the study were used in this phase. The list of sensory terms included descriptors of visual appearance (color), texture (stickiness, elasticity, resistance to cutting, firmness, chewiness), tactile in mouth (astringency), odor and flavor (semolina, cooked pasta, off-odor and off-flavor), taste (salty, sweet) and an overall judgment, defined in Table 4. They were rated on an anchored line scale that provided a 1-9 score range. The samples, presented in white dishes coded using a three-digit number, were distributed simultaneously and anonymously to all panel members. Evaluations were led in single boxes and two sensory testing sessions were carried out. The order of presentation was randomized among judges and sessions. All data were acquired by a direct computerized registration system (FIZZ Biosystemes, ver. 2.00 M, Couternon, France).

**Statistical analysis:** Data were submitted to one-way analysis of variance (ANOVA) by XLStat software (Addinsoft SARL, New York, NY, USA).

**Acknowledgments** - The authors acknowledge the financial support of MIUR in the framework of the Project PON/01\_01145/1-8 “ISCOCEM”.

## References

- [1] Lee J, Koo N, Min DB. (2004) Reactive oxygen species, aging, and antioxidative nutraceuticals. *Comprehensive Reviews in Food Science and Food Safety*, **3**, 21-33.
- [2] Roberfroid MB. (2000) Concepts and strategy of functional food science: the European perspective. *The American Journal of Clinical Nutrition*, **71**, 1660s-1664s.
- [3] Itagaki S, Kurokawa T, Nakata C, Saito Y, Oikawa S, Kobayashi M, Hirano T, Iseki K. (2009) *In vitro* and *in vivo* antioxidant properties of ferulic acid: A comparative study with other natural oxidation inhibitors. *Food Chemistry*, **114**, 466-471.
- [4] Bengmark S. (2006) Impact of nutrition on ageing and disease. *Current Opinions in Clinical Nutrition and Metabolic Care*, **9**, 2-7.
- [5] Manach C, Mazur A, Scalbert A. (2005) Polyphenols and prevention of cardiovascular diseases. *Current Opinion in Lipidology*, **16**, 77-84.
- [6] Branen AL. (1975) Toxicology and biochemistry of butylated hydroxy anisole and butylated hydroxy toluene. *Journal of the American Oil Chemical Society*, **52**, 59-63.
- [7] Ito N, Fukushima S, Hasegawa A, Shibata M, Ogiso T. (1983) Carcinogenicity of butylated hydroxy anisole in F344 rats. *Journal of the National Cancer Institute*, **70**, 343-352.
- [8] Naczki M, Shahidi F. (2006) Phenolics in cereals, fruits and vegetables: occurrence, extraction and analysis. *Journal of Pharmaceutical and Biomedical Analysis*, **41**, 1523-1542.
- [9] Onyeneho SN, Hettiarachchy NS. (1992) Antioxidant activity of durum wheat bran. *Journal of Agricultural and Food Chemistry*, **40**, 1496-1500.
- [10] Saadi A, Lempereur I, Sharonov S, Autran JC, Manfait M. (1998) Spatial distribution of phenolic materials in durum wheat grain as probed by confocal fluorescence spectral imaging. *Journal of Cereal Science*, **28**, 107-114.
- [11] Pasqualone A, Delvecchio LN, Mangini G, Blanco A. (2013) Effect of cultivar and roller milling process on the level of phenolic compounds in Italian durum wheat. In *Proceedings of 11th International Gluten Workshop. August 12-15, 2012, Beijing*. He Z, Wang D. (Eds). International Maize and Wheat Improvement Center (CIMMYT), Mexico, D.F., Mexico, 169-173.
- [12] Brennan CS, Tudorica CM. (2007) Fresh pasta quality as affected by enrichment of nonstarch polysaccharides. *Journal of Food Science*, **72**, S659-S665.
- [13] Edwards NM, Biliaderis CG, Dexter JE. (1995) Textural characteristics of wholewheat pasta and pasta containing non-starch polysaccharides. *Journal of Food Science*, **60**, 1321-1324.
- [14] Foschia M, Peressini D, Sensidoni A, Brennan MA, Brennan CS. (2015) How combinations of dietary fibres can affect physicochemical characteristics of pasta. *LWT-Food Science and Technology*, **61**, 41-46.
- [15] Delvecchio LN, Pasqualone A. (2013) Production trials of fresh pasta enriched with phenolic compounds extracted from wheat by KOH-induced hydrolysis. *Progress in Nutrition*, **14**, 247-251.
- [16] Wang L, Weller CL. (2006) Recent advances in extraction of nutraceuticals from plants. *Trends in Food Science and Technology*, **17**, 300-312.
- [17] Verma B, Hucl P, Chibbar RN. (2009) Phenolic acid composition and antioxidant capacity of acid and alkali hydrolysed wheat bran fractions. *Food Chemistry*, **116**, 947-954.
- [18] Laddomada B, Durante M, Minervini F, Garbetta A, Cardinali A, D'Antuono I, Caretto S, Blanco A, Mita G. (2015) Phytochemical characterization and anti-inflammatory activity of extracts from the whole-meal flour of Italian durum wheat cultivars. *International Journal of Molecular Sciences*, **16**, 3512-3527.
- [19] Lempereur I, Rouau X, Abecassis J. (1997) Genetic and agronomic variation in arabinoxylan and ferulic acid contents of durum wheat (*Triticum durum* L.) grain and its milling fractions. *Journal of Cereal Science*, **25**, 103-110.
- [20] Cubadda R, Carcea M, Pasqui LA. (1992) Suitability of the gluten index method for assessing gluten strength in durum wheat and semolina. *Cereal Foods World*, **37**, 866-869.
- [21] D'Egidio MG, Mariani BM, Nardi S, Novaro P, Cubadda R. (1990) Chemical and technological variables and their relationships: a predictive value equation for pasta cooking quality. *Cereal Chemistry*, **67**, 275-281.
- [22] Brescia MA, Sacco D, Sgarbetta A, Pasqualone A, Simeone R, Peri G, Sacco A. (2007) Characterisation of different typical Italian breads by means of traditional, spectroscopic and image analyses. *Food Chemistry*, **104**, 429-438.
- [23] Pasqualone A, Caponio F, Simeone R. (2004) Quality evaluation of re-milled durum wheat semolinas used for bread-making in Southern Italy. *European Food Research and Technology*, **219**, 630-634.
- [24] Raffo A, Pasqualone A, Sinesio F, Paoletti F, Quaglia G, Simeone R. (2003) Influence of durum wheat cultivar on the sensory profile and staling rate of Altamura bread. *European Food Research and Technology*, **218**, 49-55.
- [25] Decreto del Presidente della Repubblica 9 febbraio 2001, n. 187 Regolamento per la revisione della normativa sulla produzione e commercializzazione di sfarinati e paste alimentari, a norma dell'articolo 50 della legge 22 febbraio 1994, n. 146. *Official Journal of Italian Republic*, **117**, 6-12.
- [26] Hirawan R, Ser WY, Arntfield SD, Beta T. (2010) Antioxidant properties of commercial, regular-and whole-wheat spaghetti. *Food Chemistry*, **119**, 258-264.
- [27] Verma B, Hucl P, Chibbar RN. (2008) Phenolic content and antioxidant properties of bran in 51 wheat cultivars. *Cereal Chemistry*, **85**, 544-549.
- [28] Liu Q, Qiu Y, Beta T. (2010) Comparison of antioxidant activities of different colored wheat grains and analysis of phenolic compounds. *Journal of Agricultural and Food Chemistry*, **58**, 9235-9241.
- [29] Punzi R, Paradiso A, Fasciano C, Trani A, Faccia M, de Pinto MC, Gambacorta G. (2014) Phenols and antioxidant activity *in vitro* and *in vivo* of aqueous extracts obtained by ultrasound-assisted extraction from artichoke by-products. *Natural Product Communications*, **9**, 1315-1318.
- [30] American Association of Cereal Chemists. (2000) *Approved Methods of the American Association of Cereal Chemists*. AACC, St Paul, MN.
- [31] Pasqualone A, Delvecchio LN, Mangini G, Taranto F, Blanco A. (2014) Variability of total soluble phenolic compounds and antioxidant activity in a collection of tetraploid wheat. *Agricultural and Food Science*, **23**, 307-316.
- [32] UNI-Ente Italiano di Normazione. (2003) *Metodo 10957. Analisi sensoriale. Metodo per la definizione del profilo sensoriale degli alimenti e delle bevande*. UNI, Milan, Italy.

<b>A Novel Ellagic Acid Derivative from <i>Desbordesia glaucescens</i></b> Faustine L. DongmoMafodong, Apollinaire Tsopmo, Maurice D. Awouafack, Tchuenguem T. Roland, Jean P. Dzoyem and Pierre Tane	1709
<b>Proliferative Constituents from the Leaves of <i>Micromelum integerrimum</i></b> Yuan-bin Hu, Jun-ying Sun, Tang-yun Yu, Jing-cheng Wang, Jin-shan He, Yue-lai Zhou and Xian-wen Li	1711
<b>Xylactam B, A New Isobenzofuranone from an Endophytic <i>Xylaria</i> sp.</b> Nelum P. K. G. Piyasena, Anja Schöffler and Hartmut Laatsch	1715
<b>Separation of Aeruginosin-865 from Cultivated Soil Cyanobacterium (<i>Nostoc</i> sp.) by Centrifugal Partition Chromatography \combined with Gel Permeation Chromatography</b> José Cheel, Mirjana Minceva, Petra Urajová, Rabya Aslam, Pavel Hrouzek and Jiří Kopecký	1719
<b>Effect of Maturation Degree on Composition of Fatty Acids and Tocopherols of Fruit Oil from <i>Pistacia atlantica</i> Growing Wild in Algeria</b> Hamid Guenane, Isabelle Bombarda, Mohamed Didi OuldElhadj and Mohamed Yousfi	1723
<b>Chemoenzymatic Total Synthesis of (<i>R</i>)-4-Dodecanolide and (<i>R</i>)-4-Octanolide</b> Chittamuru Sreelakshmi, Adari Bhaskar Rao, Mangamoori Lakshmi Narasu, Anugu Srinivas Reddy, Palakondur Janardhan Reddy and Basi V. SubbaReddy	1729
<b>Inspired by Nature: The Use of Plant-derived Substrate/Enzyme Combinations to Generate Antimicrobial Activity <i>in situ</i></b> Ethiene Castellucci Estevam, Sharon Griffin, Muhammad Jawad Nasim, Dariusz Zieliński, Justyna Aszyk, Magdalena Osowicka, Natalia Dawidowska, Rinaldi Idroes, Agnieszka Bartoszek and Claus Jacob	1733
<b>Effect of Supplementation with Wheat Bran Aqueous Extracts Obtained by Ultrasound-Assisted Technologies on the Sensory Properties and the Antioxidant Activity of Dry Pasta</b> Antonella Pasqualone, Laura Nunzia Delvecchio, Giuseppe Gambacorta, Barbara Laddomada, Valeria Urso, Agata Mazzaglia, Paolo Ruisi and Giuseppe Di Miceli	1739
<b>Development of SCAR Markers Based on Improved RAPD Amplification Fragments and Molecular Cloning for Authentication of Herbal Medicines <i>Angelica sinensis</i>, <i>Angelica acutiloba</i> and <i>Levisticum officinale</i></b> Chun Zhang, Zhiqiang Mei, Jingliang Cheng, Yin He, Md. Asaduzzaman Khan, Peiyi Luo, Saber Imani and Junjiang Fu	1743
<b>Chemical Composition of <i>Blumea lacera</i> Essential Oil from Nepal. Biological Activities of the Essential Oil and (<i>Z</i>)-Lachnophyllum Ester</b> Prabodh Satyal, Bhuwan K. Chhetri, Noura S. Dosoky, Samon Shrestha, Ambika Poudel and William N. Setzer	1749
<b>Essential Oil Composition and Antibacterial Activity of <i>Hyptis colombiana</i> from the Venezuelan Andes</b> Mayalín Flores, Luis Rojas, Rosa Aparicio, María Eugenia Lucena and Alfredo Usubillaga	1751
<b>Repellent and Anti-quorum Sensing Activity of Six Aromatic Plants Occurring in Colombia</b> Leonor Cervantes-Ceballos, Karina Caballero-Gallardo and Jesus Olivero-Verbel	1753
<b>Chemical Composition and Larvicidal Activity of Greek Myrtle Essential Oils against <i>Culex pipiens</i> biotype <i>molestus</i></b> Aikaterini Koutsaviti, Irene Lignou, Ioannis Bazos, George Koliopoulos, Antonios Michaelakis, Athanassios Giatropoulos and Olga Tzakou	1759
<b>Chemical Composition Variability of the Herb Essential Oil in the Ontogenesis of <i>Artemisia campestris</i> subsp. <i>campestris</i></b> Anna Lis, Martyna Kowal and Joanna Kończak	1763

### Accounts/Reviews

<b>Sesquiterpenes from Essential Oils and Anti-Inflammatory Activity</b> Rita de Cássia da Silveira e Sá, Luciana Nalane Andrade and Damião Pergentino de Sousa	1767
<b>The Interaction of Alpha-synuclein with Membranes and its Implication in Parkinson's Disease: A Literature Review</b> Azucena Gonzalez-Horta	1775
<b>Herbal Supplements and Hepatotoxicity: A Short Review</b> Haszianaliza Haslan, Fariha Hajj Suhaimi and Srijit Das	1779
<b>Vibrational Circular Dichroism: Recent Advances for the Assignment of the Absolute Configuration of Natural Products</b> Eleuterio Burgueño-Tapia and Pedro Joseph-Nathan	1785

# Natural Product Communications

## 2015

Volume 10, Number 10

### Contents

#### Natural Products from Endophytes

(Guest Editors: John A. Johnson and Christopher A. Gray)

##### Original Paper

- Antimycobacterial Natural Products from Endophytes of the Medicinal Plant *Aralia nudicaulis***  
Haixin Li, Brandon Doucet, Andrew J. Flewelling, Stéphanie Jean, Duncan Webster, Gilles A. Robichaud, John A. Johnson and Christopher A. Gray 1641
- Microsporols A–C from the Plant Endophytic Fungus *Pestalotiopsis microspora***  
Xianfu Wu, Yadan Wang, Shuchun Liu, Xinzhong Liu and Liangdong Guo 1643
- Isolation of Phomopsolide A and 6(E)-Phomopsolide A as Antimycobacterial Natural Products from an Unidentified Endophyte of the Canadian Medicinal Plant *Heracleum maximum***  
Trevor N. Clark, Katelyn T. Ellsworth, Stéphanie Jean, Duncan Webster, Gilles A. Robichaud, John A. Johnson and Christopher A. Gray 1647
- Pyroloquin A, a 3-Decalinoyltetramic Acid with Selective Biological Activity, Isolated from Amazonian Cultures of the Novel Endophyte *Diaporthales* sp. E6927E**  
Eric V. Patridge, Alicia Darnell, Kaury Kucera, Gillian M. Phillips, Heidi R. Bokesch, Kirk R. Gustafson, Daniel J. Spakowicz, Linda Zhou, William M. Hungerford, Mark Plummer, Denton Hoyer, Alexandra Narváez-Trujillo, Andrew J. Phillips and Scott A. Strobel 1649
- Cytotoxic Cytochalasins and Other Metabolites from Xylariaceae sp. FL0390, a Fungal Endophyte of Spanish Moss**  
Ya-ming Xu, Bharat P. Bashyal, Mangping X. Liu, Patricia Espinosa-Artiles, Jana M. U'Ren, A. Elizabeth Arnold and A. A. Leslie Gunatilaka 1655
- Endophytic Fungus *Nigrospora oryzae* from a Medicinal plant *Coccinia grandis*, a High Yielding New Source of Phenazine-1-carboxamide**  
Dharushana Thanabalasingam, N. Savitri Kumar, Lalith Jayasinghe and Yoshinori Fujimoto 1659
- Polyketides from an Endophytic *Aspergillus fumigatus* Isolate Inhibit the Growth of *Mycobacterium tuberculosis* and MRSA**  
Andrew J. Flewelling, Amanda I. Bishop, John A. Johnson and Christopher A. Gray 1661
- A 3-Vinyl Cephem Derivative, a Useful Intermediate in the Synthesis of Cepham Antibiotics, from *Aspergillus awamori* Associated with Banana Fruit**  
H.M.S.K.H. Bandara, N. Savitri Kumar, Lalith Jayasinghe, Hironori Masubuti and Yoshinori Fujimoto 1663
- Two New Cyclic Depsipeptides from the Endophytic Fungus *Fusarium* sp.**  
Fang Lv, Georgios Daletos, Wenhan Lin and Peter Proksch 1667

##### Accounts/Reviews

- Bioactive Secondary Metabolites Produced by the Fungal Endophytes of Conifers**  
Andrea A. Stierle and Donald B. Stierle 1671

##### Original Paper

- Anti-diabetic Effect of Friedelan Triterpenoids in Streptozotocin Induced Diabetic Rat**  
Amitava Mandal, Vaskar Das, Pranab Ghosh and Shilpi Ghosh 1683
- Colochirosides B<sub>1</sub>, B<sub>2</sub>, B<sub>3</sub> and C, Novel Sulfated Triterpene Glycosides from the Sea Cucumber *Colochirus robustus* (Cucumariidae, Dendrochirotida)**  
Alexandra S. Silchenko, Anatoly I. Kalinovsky, Sergey A. Avilov, Pelageya V. Andryjaschenko, Pavel S. Dmitrenok, Vladimir I. Kalinin, Ekaterina A. Yurchenko and Igor Yu. Dolmatov 1687
- (+)-Chenabinol (Revised NMR Data) and Two New Alkaloids from *Berberis vulgaris* and their Biological Activity**  
Zdeněk Novák, Anna Hošťálková, Lubomír Opletal, Lucie Nováková, Martina Hrabínová, Jiří Kuneš and Lucie Čahlíková 1695
- The Influence of Cultivars and Phenological Phases on the Accumulation of Nevadensin and Salvigenin in Basil (*Ocimum basilicum*)**  
Botond Bernhardt, Jenő Bernáth, Attila Gere, Zoltán Kókai, Bonifác Komáromi, Szilvia Tavaszi-Sárosi, László Varga, László Sipos and Krisztina Szabó 1699
- Isoorientin, a Selective Inhibitor of Cyclooxygenase-2 (COX-2) from the Tubers of *Pueraria tuberosa***  
Manne Sumalatha, Rachakunta Munikishore, Aluru Rammohan, Duvvuru Gunasekar, Kotha Anil Kumar, Kakularam Kumar Reddy, Rajaram Azad, Pallu Reddanna and Bernard Bodo 1703
- Secondary Metabolites of *Alchemilla persica* Growing in Iran (East Azarbaijan)**  
Fariba Heshmati Afshar, Filippo Maggi, Sara Ferrari, Gregorio Peron and Stefano Dall'Acqua 1705

Continued inside backcover