



## Review article

# Association between autism spectrum disorder and diabetes: systematic review and meta-analysis

Samuele Cortese<sup>a,b,c,d,e</sup>, Alessandra Gabellone<sup>f</sup>, Lucia Marzulli<sup>f</sup>, Iciar Iturmendi-Sabater<sup>g,h,i</sup>, Diego de La Chica-Duarte<sup>g</sup>, Isabella M. Piqué<sup>g,j,k</sup>, Marco Solmi<sup>a,l,m,n</sup>, Jae Il Shin<sup>o</sup>, Lucia Margari<sup>f</sup>, Gonzalo Arrondo<sup>a,p,\*</sup>

<sup>a</sup> Centre for Innovation in Mental Health, School of Psychology, Faculty of Environmental and Life Sciences, University of Southampton, Southampton, UK

<sup>b</sup> Clinical and Experimental Sciences (CNS and Psychiatry), Faculty of Medicine, University of Southampton, Southampton, UK

<sup>c</sup> Solent NHS Trust, Southampton, UK

<sup>d</sup> Hassenfeld Children's Hospital at NYU Langone, New York University Child Study Center, New York City, New York, USA

<sup>e</sup> Division of Psychiatry and Applied Psychology, School of Medicine, University of Nottingham, Nottingham, UK

<sup>f</sup> Department of Biomedical Sciences and Human Oncology, University of Bari "Aldo Moro", Bari, Italy

<sup>g</sup> School of Education and Psychology, University of Navarra, Pamplona, Spain

<sup>h</sup> Division of Psychology and Language Sciences University College London, London, UK

<sup>i</sup> Institute of Medical Science, University of Toronto, Toronto, Canada

<sup>j</sup> Facultad de Ciencias Biomédicas, Universidad Austral, Pilar, Argentina

<sup>k</sup> Centro de Desarrollo y Fortalecimiento de la Niñez, Adolescencia y Familia Dr. Ramón Carrillo, San Isidro, Argentina

<sup>l</sup> Department of Psychiatry, University of Ottawa, Ottawa, ON, Canada

<sup>m</sup> Department of Mental Health, The Ottawa Hospital, Ottawa, ON, Canada

<sup>n</sup> Ottawa Hospital Research Institute (OHRI) Clinical Epidemiology Program University of Ottawa, Ottawa, ON, Canada

<sup>o</sup> Department of Pediatrics, Yonsei University College of Medicine, Seoul, South Korea

<sup>p</sup> Mind-Brain Group, Institute for Culture and Society, University of Navarra, Pamplona, Spain

## ARTICLE INFO

## Keywords:

Autism

Diabetes

Meta-analysis

## ABSTRACT

There is mixed evidence on the link between autism spectrum disorder (ASD) and diabetes. We conducted the first systematic review/meta-analysis on their association. Based on a pre-registered protocol (PROSPERO: CRD42021261114), we searched Pubmed, Ovid, and Web of Science databases up to 6 December 2021, with no language/type of document restrictions. We assessed study quality using the Newcastle-Ottawa Scale (NOS). We included 24 studies (total: 3427,773 individuals; 237,529 with ASD and 92,832 with diabetes) in the systematic review and 20 in the meta-analysis (mean stars number on the NOS: 5.89/10). There was a significant association, albeit characterized by significant heterogeneity, when pooling unadjusted OR (1.535, 95% CI = 1.109–2.126), which remained significant when restricting the analysis to children and type 2 diabetes, but became non-significant when considering adjusted ORs (OR: 1.528, 95% CI = 0.954–2.448). No significant prospective association was found ( $n = 2$ ) on diabetes predicting ASD (HR: 1.232, 0.826–11.837). Therefore, the association between ASD and diabetes is likely confounded by demographic and clinical factors that should be systematically investigated in future studies.

## 1. Introduction

There is increasing evidence that conditions classically thought as disorders of the central nervous system are actually characterized by abnormalities in other physiological systems (Qureshi and Mehler, 2013). This has prompted a growing line of research on the association between neuropsychiatric and physical conditions (Cortese et al., 2020).

Among other neuropsychiatric conditions, autism spectrum disorder (ASD) has been the focus of research in this area due to the increasing awareness of its impact on the life of affected individuals and their families. Indeed, beyond its defining core symptoms, i.e., qualitative differences in social interaction/communication and repetitive, restrictive, or unusual sensori-motor behaviors/interest (Lord et al., 2018), ASD is often characterized by a range of comorbidities. In addition to

\* Correspondence to: Mind-Brain Group, Institute for Culture and Society (ICS), University of Navarra, Pamplona, Navarra 31009, Spain.

E-mail address: [garrondo@unav.es](mailto:garrondo@unav.es) (G. Arrondo).

<https://doi.org/10.1016/j.neubiorev.2022.104592>

Received 11 January 2022; Received in revised form 16 February 2022; Accepted 20 February 2022

Available online 22 February 2022

0149-7634/© 2022 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

neuropsychiatric conditions such as anxiety, ADHD, and depressive disorders (Lai et al., 2019), meta-analytic evidence shows a significant association between autism and important physical conditions including obesity (Kahathuduwa et al., 2019), dermatitis (Tsai et al., 2020), and food allergy (Li et al., 2021). Some studies have explored a possible cross-sectional or prospective association with diabetes. These studies were prompted by evidence of a significant association between gestational diabetes and ASD (Rowland and Wilson, 2021), possibly accounted for by the impact of hyperglycemia on neurodevelopment via oxidative stress (Wells et al., 2009).

The association of ASD to diabetes might be plausible from a neurobiological standpoint. In relation to the association with diabetes type 1, shared inflammatory factors might be considered from a theoretical perspective at least. Indeed, it is well established that diabetes type 1 is related to the destruction of insulin-producing beta cells via autoimmune-mediated mechanisms (DiMeglio et al., 2018). Interestingly, some studies have reported evidence of immunological dysfunctions (neuroinflammation, increased autoantibodies, cytokines and immunoglobulins or abnormalities in immune cells,) in individuals with ASD (Bjørklund et al., 2016; Masi et al., 2017). Therefore, it has been proposed that individuals with ASD are at increased risk of disorders mediated by immunological dysfunctions such as diabetes type 1, asthma, inflammatory bowel disease and atopic dermatitis (Zerbo et al., 2015). As for the association with diabetes type 2, which is related to insulin resistance and relative insulin deficiency (Chatterjee et al., 2017), it has been postulated that the link may be mediated by the significant association of ASD to obesity and dyslipidemia, as well as by the metabolic side effects of antipsychotic medications that are quite commonly prescribed in individuals with ASD (Tromans et al., 2021).

However, whilst a significant association with diabetes (both type 1 (e.g., Freeman et al., 2005), and 2 (e.g., Chen et al., 2016)) has been found in some studies, other studies, e.g. (Bethin et al., 2019), have failed to replicate these positive findings. Given this mixed evidence, there is a need to quantitatively synthesize available evidence. Therefore, the aim of this study was to conduct the first systematic review with meta-analyses of studies (cross-sectional or prospective) providing data on the link between ASD and diabetes.

## 2. Methods

We followed the 2020 PRISMA guidelines (Page et al., 2021), and pre-registered the protocol in PROSPERO (CRD42021261114). As per the original, pre-registered protocol, we aimed to address two questions: 1) Is there a significant association between autism spectrum disorder (ASD) and diabetes in children, adolescents, or adults? 2) Is there a significant association between diabetes in mothers during pregnancy and ASD in their children? As we deemed that reporting the results related to both objectives would have been impractical in a single paper, the current paper is related to question #1. Of note, recent meta-analytic evidence has been published in relation to question #2 (Rowland and Wilson, 2021). We will update this evidence and report the results of question #2 in a separate, subsequent paper.

### 2.1. Search

We searched PubMed (MEDLINE), Ovid databases (PsycINFO, EMBASE + EMBASE Classic, Ovid MEDLINE), and Web of Knowledge [Web of Science (Science Citation Index Expanded), Biological Abstracts, BIOSIS, Food Science and Technology Abstracts] from inception to 6 December 2021 with no restrictions in terms of type of document or language. The syntax for PubMed, adapted for the other databases, was as follows: (autis\* [tiab] or asperger [tiab] or "pervasive developmental disorder" [tiab]) AND diabet\*. References of relevant systematic reviews were hand searched to retrieve any additional eligible study.

### 2.2. Inclusion criteria

We included observational (cross-sectional or prospective) studies providing data on the strength of the association between ASD and diabetes in children, adolescents or adults. Eligible definitions of ASD were as follows: 1) a categorical diagnosis of ASD according to the DSM III, DSM-III-R, DSM-IV, DSM-IV-TR, and DSM-5 or ICD 9 and ICD-10; 2) diagnosis based on the ADOS (Autism Diagnostic Observation Schedule) or similar validated tools; 3) a positive answer to the question: "Did your doctor ever tell you that you have ASD?" or similar; 4) a diagnosis of ASD recorded in medical files/registries. We planned a sensitivity analysis restricted to formal diagnosis of ASD, confirmed by (semi) structured interviews as per DSM/ICD criteria. Studies including individuals of any age and of both genders were eligible. Studies were included regardless of the past or current treatment of the participants with psychotropics for symptoms of ASD or related impairments (e.g., sleep or ADHD symptoms). However, we planned to assess the feasibility of conducting a sensitivity analysis including only studies with ASD medication naïve participants. Control groups recruited due to having other disorders potentially related to a differential risk of diabetes or ASD were not accepted (e.g., groups of individuals with another psychiatric disorder). Siblings of ASD participants were accepted as controls. Studies including participants recruited in clinical settings or in the general population were included.

### 2.3. Outcomes

The main outcome was the unadjusted ratio (odds ratio-OR-, risk ratio-RR-, hazard ratio -HR- or equivalent, prioritizing the OR) expressing the association between ASD and diabetes. Secondary outcome was the maximally adjusted ratio, when available. The choice of the primary and secondary outcome was made based on the estimated number of eligible studies, expected to be higher for studies reported unadjusted estimates, and, hence, more suitable for secondary/sensitivity analyses, but arguably the adjusted ORs are more informative, as potentially less prone to biased derived from the selection of the groups.

### 2.4. Screening and data extraction

In order to minimize the possibility of missing relevant studies, four researchers (Alessandra Gabellone; Lucia Marzulli; Iciar Iturmendi-Sabater; Diego de La Chica-Duarte) carried out the screening stage independently and any study considered of interest by any of them was moved to the full-text evaluation phase. Full texts from retrieved references were independently and blindly coded for eligibility by two study authors. Any disagreement was resolved by the first and last author. Two authors independently extracted data from the retained studies. Per protocol, we planned to extract the following information/data: 1) Publication detail: year and language of publication, country where the study was conducted; 2) Design: type of study (including cross-sectional, case-control, cohort.); 3) Study temporality (cross-sectional, prospective, retrospective); 4) Patient enrolment (consecutive, non-consecutive); 5) Setting (clinical vs. population-based study); 6) Study participants details: number, mean age (SD), gender distribution, SES and ethnicity of participants with and without ASD and diabetes; 6) Psychiatric comorbidities of individuals with and without ASD (type and prevalence); 7) Method to establish the diagnosis of ASD (self-reported diagnosis, diagnosis recorded in medical files/registry, structured or semi-structured interview); 8) Medication status of individuals with and without ASD (type of medication and percentage of treated participants, during and prior to the study); 9) Outcome (OR, RR, HR or equivalent)- we extracted both adjusted and unadjusted ratios if available and we used the most adjusted (= maximally adjusted) ratios in the analyses, 10) Number of outcomes within the exposed and unexposed participants- this was used to obtain the unadjusted odds ratio whenever it was not provided.




Fig. 1. PRISMA 2020 flow chart- selection of studies.

## 2.5. Risk of bias/quality assessment

Two authors independently performed the assessment of risk of bias in the included studies using the Newcastle–Ottawa Scale, NOS (Wells et al., 2000). Any discrepancies in the rating of study quality/risk of bias was resolved by consensus between the two authors. If this was not possible, the first and last authors made a judgement on rating and acted as arbitrators.

## 2.6. Data synthesis

Odds ratios expressing the association between ASD and diabetes were extracted when available or calculated from available data. We conducted two analyses: one focused on non-adjusted odds ratios, and another one on ratios adjusted (by study authors of individual studies) for possible confounders, that inevitably varied from study to study. Meta-analyses used random-effects models (with the DerSimonian and Laird procedure) because it allows the true population effect size to differ among studies. In the adjusted analysis we pooled HRs together to ORs as both diabetes and ASD occur unfrequently. Nevertheless, studies reporting HRs were later removed in a sensitivity analysis.

Per protocol, we planned to assess the feasibility of conducting the following subgroup meta-analyses: 1) analyzing children/adolescents and adults separately; 2) analyzing clinical and population-based studies separately; 3) including only studies with current (as opposed to lifetime) prevalence of ASD and diabetes; 4) including only studies with a diagnosis of ASD confirmed by structured/semi-structured interviews; 5) including only studies with ASD medications naive participants; 6) excluding studies that used siblings as controls; 7) including only longitudinal studies separately. We also planned to perform a meta-regression analysis including unadjusted ratios as outcome and year of study publication, gender, and the rating on the Newcastle-Ottawa Scale as regressors. The meta-analyses and meta-regressions were weighted by the reciprocal of the variance of the effect size, which gives greater weight to larger studies. We used the Q (indicating the presence, but not the degree, of heterogeneity) and the  $I^2$  index (percentage of variance

due to true heterogeneity) to estimated indices of heterogeneity of effect sizes. We used Egger's test to assess publication bias and Duval and Tweedie's trim and fill procedure to adjust for it. Analyses were performed using Comprehensive Meta-Analysis v3 (Biostat Inc, 2021).

## 3. Results

### 3.1. Overview of included studies

The search in the electronic databases yielded 2678 references. Fleiss' Kappa, a measure of the agreement between more than two raters where agreement due to chance is factored out, was 0.52, indicating a moderate agreement (Zaiontz, 2021a). Cohen's Kappa when judging inclusion versus exclusion in the full text screening phase was 0.60, which is also considered a moderate rate of agreement (Zaiontz, 2021b). After the screening process, 24 studies (reported in 27 publications) were included in the qualitative review (Alabaf et al., 2019; Barrett et al., 2021; Bilder et al., 2017; Butwicka et al., 2015; Chen et al., 2009, 2016, 2013; Croen et al., 2015; Dybdal et al., 2018; Flygare Wallén et al., 2018; Gurney et al., 2006; Kohane et al., 2012; Liu et al., 2021; McDermott et al., 2007; McManus et al., 2009; Schott et al., 2021; Shedlock et al., 2016; Spann et al., 2019; Supekar et al., 2017; Tyler et al., 2011; Vohra et al., 2017; Weir et al., 2021; Weiss et al., 2018; Zerbo et al., 2015). Fig. 1 shows the screening process. Table S1 reports the list of studies excluded after checking the full text, with reason for exclusion. Table S2 details the included studies reported in multiple records (papers).


The total sample size of the studies included in the systematic review was 3427,773 individuals, out of which 237,529 had ASD and 92,832 diabetes. The main study characteristics are reported in Table 1. Table S3 provides additional information on the cases and controls in each study.

Retained studies were all published in English, between 2006 and 2021, and were carried out in North America (n = 12), Northern Europe (Sweden n = 4, Denmark n = 1, Finland n = 1, United Kingdom n = 1) and Taiwan (n = 3), with an additional study being an internet-based transnational survey. Most studies were based on administrative

**Table 1**  
Characteristics of the included studies. NOS: Total number of stars in the Newcastle-Ottawa scale.

| Author year                       | Country        | Database                                                                                                       | Design          | Diagnosis of ASD                    | Diagnosis of diabetes               | Diabetes subtype | Age                                                        | Gender (% M)       | Total N | ASD N  | Diabetes N | NOS          |
|-----------------------------------|----------------|----------------------------------------------------------------------------------------------------------------|-----------------|-------------------------------------|-------------------------------------|------------------|------------------------------------------------------------|--------------------|---------|--------|------------|--------------|
| Alabaf et al., 2019               | Sweden         | Child and Adolescent Twin Study 1992–2006                                                                      | Cross-sectional | Clinical (A-TAC)                    | Self-report                         | Not specified    | 9–12 years                                                 | 50.92%             | 22329   | 301    | 93         | 7            |
| Barrett et al., 2021              | USA            | National Survey of Children's Health 2006–2019                                                                 | Cross-sectional | Self-report                         | Self-report                         | Not specified    | Children and adolescents 3–17 years                        | Not specified      | 131774  | 3601   | 593        | 4            |
| Bilder et al., 2017               | USA            | MarketScan® Research Databases                                                                                 | Cross-sectional | Registers                           | Registers                           | T1&T2            | Adults ≥ 18                                                | 39.32%             | 29786   | 37     | 7060       | 7            |
| Butwicka et al., 2015             | Sweden         | Multiple national registries                                                                                   | Longitudinal    | Registers                           | Registers                           | T1               | < 18 years                                                 | Not specified      | 1713733 | 7038   | 17122      | 9            |
| Chen et al., 2009                 | Taiwan         | National Health Insurance Research Databases, 1997–2004                                                        | Cross-sectional | Registers                           | Registers                           | T1&T2            | Children < 19 years                                        | Not specified      | 37831   | 3440   | 102        | 7            |
| Chen et al., 2013                 | Taiwan         | National Health Insurance Research Databases, 1996–2010                                                        | Cross-sectional | Registers                           | Registers                           | T1               | Any age                                                    | Not specified      | 7994    | 1598   | 8          | 7            |
| Chen et al., 2016                 | Taiwan         | National Health Insurance Research Databases, 2002–2009                                                        | Longitudinal    | Registers                           | Registers                           | T2               | Adolescents (10–17 years) and young adults (18–29 years)   | Not specified      | 30610   | 6122   | 187        | 9            |
| Croen et al., 2015                | USA            | Kaiser Permanente in Northern California, 2008–2012                                                            | Cross-sectional | Registers                           | Registers                           | Not specified    | Adults ≥ 18                                                | Not specified      | 16577   | 1507   | 767        | 7            |
| Dybdal et al., 2018               | Denmark        | Multiple national registries 1996–2013                                                                         | Longitudinal    | Registers                           | Registers                           | T1               | Children. Girls M = 9.6, SD = 4.4; Boys M = 10.2, SD = 4.6 | 52%                | 40672   | 520    | 5084       | 9            |
| Gurney et al., 2006               | USA            | National Survey of Children's Health 2003–2004                                                                 | Cross-sectional | Self-report                         | Self-report                         | Not specified    | Children and adolescents 3–17 years                        | Not specified      | 85272   | 483    | 256        | 4            |
| Liu et al., 2021                  | Sweden         | Multiple national registries 1973–2013                                                                         | Longitudinal    | Registers                           | Registers                           | T1               | Children and adolescents                                   | Not specified      | 92730   | 738    | 8430       | 9            |
| Schott et al., 2021               | USA            | Medicaid Analytic eXtract , 2008–2012                                                                          | Cross-sectional | Registers                           | Registers                           | Registers        | Not specified                                              | Adults 18–65 years | 74,1    | 667808 | 155617     | 57283        |
| Shedlock et al., 2016             | USA            | Military Health System 2000–2013                                                                               | Cross-sectional | Registers                           | Registers                           | T2               | Children. M = 8.83, SD = 3.44                              | 80%                | 292572  | 48762  | 1485       | 7            |
| Spann et al., 2019                | Finland        | Finnish Prenatal Study of Autism 1987–2007                                                                     | Cross-sectional | Registers                           | Registers                           | T1               | Any age                                                    | Not specified      | 22658   | 4600   | 180        | 7            |
| Tyler et al., 2011                | USA            | Individuals With Intellectual and Other Developmental Disabilities Electronic Health Record Analysis 2005–2008 | Cross-sectional | Registers                           | Registers                           | Not specified    | Adults ≥ 18                                                | 71.65%             | 314     | 108    | 23         | 7            |
| Vohra et al., 2017                | USA            | Medicaid Analytic eXtract 2000–2008                                                                            | Cross-sectional | Registers                           | Registers                           | Not specified    | Adults 22–64 years                                         | Not specified      | 7092    | 1772   | 313        | 7            |
| Flygare Wallén et al., 2018       | Sweden         | Central Administrative Database of Stockholm County, 1998–2015                                                 | Cross-sectional | Registers                           | Registers                           | Not specified    | Any age                                                    | Not specified      | 2010061 | 13921  | 117547     | 5            |
| Weir et al., 2021                 | Multiple       | Online physical health survey, 2018–2019                                                                       | Cross-sectional | Self-report                         | Self-report                         | T2               | Adults ≥ 18                                                | Not specified      | 2368    | 1156   | 172        | 4            |
| Weiss et al., 2018                | Canada         | Health Care Access Research and Developmental Disabilities                                                     | Cross-sectional | Registers                           | Registers                           | Not specified    | Adults 18–24 years                                         | Not specified      | 398358  | 5095   | 2641       | 6            |
| Zerbo et al., 2015                | USA            | Kaiser Permanente in Northern California, 1995–2006                                                            | Cross-sectional | Registers                           | Registers                           | T1               | M = 12.15, SD = 5.2                                        | 82%                | 33390   | 5565   | 64         | 7            |
| Kohane et al., 2012               | USA            | Shared Health Research Informatics Network                                                                     | Cross-sectional | Registers                           | Registers                           | T1               | 0–34 y                                                     | NR                 | 1246494 | 9105   | 4238       | 5            |
| Supekar et al., 2017              | USA            | Stanford Translational Research Integrated Database Environment                                                | Cross-sectional | Registers                           | Registers                           | T1               | Any                                                        | 44                 | 1847365 | 4790   | 10891      | 5            |
| McDermott et al., 2007            | USA            | Two local practices in Columbia                                                                                | Cross-sectional | Clinical notes and diagnostic codes | Clinical notes and diagnostic codes | Not specified    | Not reported                                               | Not reported       | 1873    | 45     | 272        | 4            |
| McManus et al., 2009 <sup>a</sup> | United Kingdom | Adult psychiatric morbidity in England Household Survey                                                        | Cross-sectional | Clinical (ADOS)                     | Not reported                        | Not specified    | Adults                                                     | Not reported       | 618     | 19     | 38         | Not reported |

<sup>a</sup> Data obtained from Tromans et al. (2021), which in turn obtained it directly from raw data of the survey.



**Fig. 2.** Forest plot. Meta-analysis of unadjusted ORs. Negative association means a decreased risk of ASD with diabetes while a positive association means an increased risk of ASD with diabetes.

registers, whereas four relied on self-reported surveys (Barrett et al., 2021; Gurney et al., 2006; McManus et al., 2009; Weir et al., 2021), and one on a twin database and clinical diagnoses (Alabaf et al., 2019). Four studies aimed to assess prospective associations (Butwicka et al., 2015; Chen et al., 2016; Dybdal et al., 2018; Liu et al., 2021) and were considered cohort studies for the purpose of risk of bias assessment, while the remaining studied cross-sectional associations and the case-control version of the NOS was used to evaluate their risk of bias. Regarding diabetes, eight studies dealt with the association between ASD and type 1 diabetes, three with type 2 diabetes, and the remaining with both types. Four studies included likely overlapping samples from Sweden (Alabaf et al., 2019; Butwicka et al., 2015; Flygare Wallén et al., 2018; Liu et al., 2021), and two from the USA (Schott et al., 2021; Vohra et al., 2017) so only the largest sample obtained at the country level was included in the meta-analysis (Butwicka et al., 2015; Schott et al., 2021). Similarly, there were three studies with likely overlapping samples from


Taiwan (Chen et al., 2009, 2016, 2013). They were not combined into the same analysis, but instead, contributed to specific sensitivity analyses. Hence, 20 studies were finally meta-analyzed.

Table S4 and S5 report the summary of the risk of bias/quality assessment for case-control and cohort study, respectively. The mean number of stars in the Newcastle-Ottawa scale for case-control studies was 5.89 (Standard deviation-SD = 1.52) and of 9 (SD = 0) for cohort studies.

Table S6 provides data on the outcome matrix, i.e., the number of individuals with and without ASD and with and without diabetes (2 × 2 matrix).

### 3.2. Results of the meta-analyses

When pooling unadjusted ORs ( $k = 18$ ), the summary effect indicated a significant association (OR: 1.58, 95% CI = 1.12–2.22), albeit



**Fig. 3.** Forest plot. Meta-analysis of maximally adjusted effect sizes. Negative association means a decreased risk of ASD with diabetes while a positive association means an increased risk of ASD with diabetes.



**Table 2**

Covariates adjusted for in relation to maximally adjusted outcomes retained in the analyses.

| Author_year               | Type | ES (95%CI)          | Covariates adjusted for statistically                                                                                                                                                                                                                                                                                                                                                              | Direction        |
|---------------------------|------|---------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------|
| Butwicka et al., 2015     | HR   | 1.70<br>(1.40,2.00) | Socioeconomic factors (maternal/paternal age at childbirth, maternal/paternal psychiatric history, maternal/paternal country of birth, level of education of higher educated parent), perinatal variables (gestational age, birth weight, being born small for gestational age, being born large for gestational age, Apgar score), and history of psychiatric disorders prior to the recruitment. | Diabetes > > ASD |
| Chen et al., 2016         | HR   | 3.25<br>(2.23,4.75) | Demographic data, use of atypical antipsychotics, and medical comorbidities                                                                                                                                                                                                                                                                                                                        | ASD > > Diabetes |
| Croen et al., 2015        | OR   | 2.18<br>(1.62,2.93) | Age, sex, and race/ethnicity                                                                                                                                                                                                                                                                                                                                                                       | NA               |
| Dybdal et al., 2018 (men) | HR   | 1.09<br>(0.82,1.46) | Age at inclusion                                                                                                                                                                                                                                                                                                                                                                                   | Diabetes > > ASD |
| Dybdal 2018 (women)       | HR   | 0.86<br>(0.49,1.50) | Age at inclusion                                                                                                                                                                                                                                                                                                                                                                                   | Diabetes > > ASD |
| Gurney et al., 2006       | OR   | 1.10<br>(0.40,3.10) | Age, sex, primary language, insurance, and household educational attainment                                                                                                                                                                                                                                                                                                                        | NA               |
| Schott et al., 2021       | OR   | 0.93<br>(0.91,0.96) | Age, race/ethnicity, months enrolled, Medicaid eligibility category, state and whether the individual had an Intellectual Disability diagnosis.                                                                                                                                                                                                                                                    | NA               |
| Spann et al., 2019        | OR   | 1.09<br>(0.77,1.56) | Maternal socioeconomic status, gestational age, and maternal and familial psychiatric diagnosis                                                                                                                                                                                                                                                                                                    | NA               |
| Vohra et al., 2017        | OR   | 0.74<br>(0.56,0.99) | Age, sex, and race/ethnicity                                                                                                                                                                                                                                                                                                                                                                       | NA               |
| Weir et al., 2021 (men)   | OR   | 1.51<br>(0.82,2.83) | Age, ethnicity, country of residence, education-level BMI, alcohol use, and smoking                                                                                                                                                                                                                                                                                                                | NA               |
| Weir et al., 2021 (women) | OR   | 1.69<br>(1.11,2.60) | Age, ethnicity, country of residence, education-level BMI, alcohol use, and smoking                                                                                                                                                                                                                                                                                                                | NA               |
| Weiss et al., 2018        | OR   | 2.78<br>(2.23,3.45) | Age, sex, rurality, and neighborhood income quintile                                                                                                                                                                                                                                                                                                                                               | NA               |

**Table 3**

Summary of results of sensitivity analyses. K: number of effect sizes. ES: pooled effect size. LBCI and UBCI: Lower and upper bound of the 95% confidence interval, pQ: p-value associated to the Q statistic of heterogeneity, PB: Evidence of publication bias (when  $p < 0.1$  for Egger test).

| Analysis                | K  | ES (95%CI)          | Q      | pQ     | I <sup>2</sup> | PB     |
|-------------------------|----|---------------------|--------|--------|----------------|--------|
| Children                | 7  | 1.49<br>(0.97,2.29) | 109.38 | 93.6%  | No             | 109.38 |
| Adults                  | 9  | 1.7<br>(1.13,2.57)  | 157.17 | 95.55% | Yes            | 157.17 |
| T1 diabetes             | 8  | 2.39<br>(2.21,2.58) | 47.54  | 87.38% | No             | 47.54  |
| T2 diabetes             | 8  | 1.44<br>(0.93,2.22) | 22.24  | 91.01% | No             | 22.24  |
| Registry diagnosis      | 7  | 1.35<br>(0.96,1.9)  | 718.84 | 98.19% | Yes            | 718.84 |
| Self-reported diagnosis | 3  | 2.57<br>(1.65,4.03) | 22.07  | 90.94% | No             | 22.07  |
| Epidemiological studies | 14 | 1.51<br>(1.05,2.19) | 828.69 | 98.19% | Yes            | 828.69 |
| Case-controlled studies | 3  | 2.2<br>(0.96,5.04)  | 799.87 | 98.25% | Yes            | 799.87 |

characterized by significant heterogeneity ( $Q = 829.29$ ,  $p < 0.05$ ,  $I^2$ : 97.95%) (Fig. 2) and evidence of publication bias (Fig. S1). When the trim and filled procedure was implemented, the association was no longer significant (OR: 0.99, 95% CI = 0.71–1.38). A very similar result was obtained when all adjusted effect sizes ( $k = 9$ ) were combined (OR: 1.532, 95% CI = 1.096–2.141,  $Q = 212.32$ ,  $I^2 = 95.29\%$ ), which also showed evidence of publication bias (Fig. 3 and S2, and Table 2 for a list of covariates adjusted for in each study). When aHRs were excluded, the summary effect size derived from pooling adjusted ORs ( $k = 7$ ) was lower and statistically not significant (OR: 1.528, 95% CI = 0.954–2.448), with significant heterogeneity ( $Q = 136.68$ ,  $p < 0.05$ ,  $I^2$ : 95.610) and publication bias (Fig. 4 and S3). Results were substantially replicated in the subgroup analysis focusing on children but not in adults, where the summary effect became not significant (Figs. S4 to S7).

Although the number of pooled studies was small, the association was significant for type 2 diabetes (OR: 2.57, 95% CI = 1.65–4.03) but not for type 1 (OR: 1.35, 95% CI = 0.96–1.90) as shown in Figs. S8 to S11. Since all studies involved diagnoses over the whole life or long time periods, we could not evaluate the effect of using only current diagnoses. Conversely, no study reported the number of medication-naïve individuals with ASD and the only study using siblings as controls was not included in the quantitative synthesis.


Furthermore, results held in the analyses focusing on population-based studies (only one study was in a clinical population), using diagnoses derived from administrative registries, or limiting the analysis to case-controlled studies (Figs. S12 to S19).

The meta-analysis of the studies reporting a prospective association (diabetes predicting ASD) showed a non-significant summary effect, with significant heterogeneity ( $Q = 9.617$ ,  $p < 0.008$ ,  $I^2 = 79.20$ ) (Fig. S20). There was only one study assessing the opposite temporal direction (ASD preceding diabetes) and it was significant (HR = 3.25, 95% CI = 2.23–4.75). Paucity of information/data prevented the conduct of the other planned analyses.

Based on the availability of data, we could only test, via meta-regression, the moderating effect of two variables: study quality rating, and year of publication, both of which were not significant ( $p > 0.05$ ).

#### 4. Discussion

To the best of our knowledge, this is the first meta-analysis of studies providing data on the association between ASD and diabetes. Whilst previous meta-analytic evidence established a significant association



**Fig. 4.** Forest plot. Meta-analysis of maximally adjusted effect sizes, excluding studies presenting hazard ratios. Negative association means a decreased risk of ASD with diabetes while a positive association means an increased risk of ASD with diabetes.

between gestational diabetes and ASD in offspring, studies assessing the association of ASD to diabetes type I or 2 in the same individual had not been meta-analyzed so far.

While our systematic review/meta-analysis points towards a statistically significant association between ASD and diabetes, robust evidence is currently lacking. Even though the primary analysis pooling all eligible studies showed a significant effect size, it was characterized by statistically significant heterogeneity, indicating the inappropriateness of considering the summary effect as representative of the true effect, and the need to gain insight into the sources of heterogeneity. While the analysis combining all adjusted effect sizes obtained similar results to our main analysis, the pooled effect size was reduced and was not statistically significant when only adjusted ORs (i.e., excluding aHRs) were combined. Of note, the covariates adjusted for inevitably varied from study to study. Furthermore, while the subgroup meta-analysis in children provided a significant effect this was not the case in the analysis in adults.

We also found no meta-analytic evidence that diabetes increases the risk of later ASD or viceversa, with just two studies showing no significant predictive role of diabetes on later ASD and a single study for the other direction.

High heterogeneity was a hallmark in all the analyses in our systematic review/meta-analysis, and we could not establish its sources in our subgroup or meta-regression analyses. Data also pointed towards publication bias in the main analysis, as well as many of the other analyses, which we tried to adjust for with the trim and fill procedure. However, it must be noted that procedures for evaluation and adjusting for such bias are less reliable with a reduced number of studies, especially when effect sizes are heterogeneous, as was our case. The risk of bias of included studies, as estimated by the Newcastle-Ottawa scale, was also variable, but consistently lower in cohort studies. However, risk of bias was not shown to be related to the direction of the effect sizes in the meta-regression analysis. Importantly, several studies failed to select an unbiased control group. Relatedly, the study by Schott et al. – the largest among the included ones – controlled for intellectual disability when selecting the control group for its ASD sample, as opposed to all other studies. Hence, these differences in design could be related to some of the discrepancies among effect sizes.

The results of the present systematic review/meta-analysis need to be considered in the light of a number of limitations. First, we overall included a relatively limited number of (heterogeneous) studies. Second, due to limitations in the reporting of individual studies we were only able to pool a very low number of studies in the analyses differentiating between type 1 and 2 diabetes, as we intended to do in a post hoc analysis. Third, due to lack of data, we could not assess a number of moderating factors (i.e., current- as opposed to lifetime- diagnosis of ASD and diabetes; effects of psychotropic medications) in the planned subgroup analyses as well as regressors in the planned meta-regression.

Fourth, whilst we could have categorized the confounding variables and using such categories in a meta-regression, given the inconsistency in the type and definition of confounding variables across studies (Table 2), we deemed that this would have been potentially misleading. Future studies should control for common and more consistently defined confounders to allow such meta-regression.

Therefore, despite representing the most comprehensive evidence synthesis to date, our systematic-review/meta-analysis points to the need of conducting large population-based studies controlling simultaneously, for a broad number of confounders, identified via directed acyclic graph, as done in previous work. For instance, following a systematic review and meta-analysis identifying confounding factors controlled in each study included in the systematic review on the association between asthma and ADHD, Cortese et al. performed a study in the Swedish registries controlling simultaneously for all the confounders identified via the systematic review (Cortese et al., 2018).

## 5. Conclusions

Despite its limitations, our work suggests that claims on a significant association between ASD and diabetes are currently not supported by robust evidence. Future research should focus on moderators that may explain significant associations in subgroups of individuals. Meanwhile, stakeholders and policy makers should be aware that any suggestion of systematically screening diabetes in individuals with ASD and vice versa is not grounded on solid evidence.

## Funding

This research received no external funding.

## CRedit authorship contribution statement

**Samuele Cortese:** Conceptualization, Methodology, Formal analysis, Investigation, Resources, Writing – original draft, Writing – review & editing, Supervision, Project administration, Funding acquisition. **Alessandra Gabellone:** Investigation, Writing – review and editing. **Lucia Marzulli:** Conceptualization, Methodology, Investigation, Writing – review & editing, Funding acquisition. **Iciar Iturmendi-Sabater:** Investigation, Writing – review & editing. **Diego de La Chica-Duarte:** Investigation, Writing – review & editing. **Isabella M. Piqué:** Investigation, Writing – review & editing. **Marco Solmi:** Conceptualization, Methodology, Writing – review & editing. **Jae Il Shin:** Conceptualization, Methodology, Writing – review & editing. **Lucia Margari:** Conceptualization, Methodology, Investigation, Writing – review & editing, Funding acquisition. **Gonzalo Arrondo:** Conceptualization, Methodology, Formal analysis, Investigation, Writing – original draft, Writing – review & editing, Supervision, Project administration.

## Institutional Review Board Statement

Not applicable.

## Informed Consent Statement

Not applicable.

## Conflicts of Interest

The authors declare no conflict of interest.

## Data Availability Statement

The data presented in this study are available in the [Supplementary material](#).

## Appendix A. Supporting information

Supplementary data associated with this article can be found in the online version at [doi:10.1016/j.neubiorev.2022.104592](https://doi.org/10.1016/j.neubiorev.2022.104592).

## References

- Alabaf, S., Gillberg, C., Lundström, S., Lichtenstein, P., Kerekes, N., Råstam, M., Anckarsäter, H., 2019. Physical health in children with neurodevelopmental disorders. *J. Autism Dev. Disord.* 49, 83–95. <https://doi.org/10.1007/s10803-018-3697-4>.
- Barrett, C.E., Bullard, K.M., Park, J., Koyama, A., Mercado, C., Claussen, A.H., Newsome, K., Pavkov, M.E., 2021. Prevalence of Mental, Behavioral, and Developmental Disorders among Youth with Diabetes, United States (2016–2019). *Diabetes* 70, 510–P. <https://doi.org/10.2337/db21-510-P>.
- Bethin, K.E., Kanapka, L.G., Laffel, L.M., Majidi, S., Chaytor, N.S., MacLeish, S., Adams, R., Foster, N.C., 2019. Autism spectrum disorder in children with Type 1 diabetes. *Diabet. Med.* 36, 1282–1286. <https://doi.org/10.1111/dme.14069>.
- Bilder, D.A., Kober, J.A., Cohen-Pfeiffer, J.L., Johnson, E.M., Jurecki, E.R., Grant, M.L., 2017. Neuropsychiatric comorbidities in adults with phenylketonuria: a retrospective cohort study. *Mol. Genet. Metab.* VO - 121 <https://doi.org/10.1016/j.ymgme.2017.03.002>.
- Biostat Inc., 2021. Comprehensive Meta-Analysis [WWW Document]. URL (<http://www.meta-analysis.com/index.php>) (accessed 12.12.21).
- Björklund, G., Saad, K., Chirumbolo, S., K, Kern, J., A, Geier, D., R, Geier, M., A, Urbina, M., 2016. Immune dysfunction and neuroinflammation in autism spectrum disorder. *Acta Neurobiol. Exp. (Wars)* 76, 257–268. <https://doi.org/10.21307/ane-2017-025>.
- Butwickia, A., Frisén, L., Almqvist, C., Zethelius, B., Lichtenstein, P., 2015. Risks of psychiatric disorders and suicide attempts in children and adolescents with type 1 diabetes: a population-based cohort study. *Diabetes Care* 38, 453–459. <https://doi.org/10.2337/dc14-0262>.
- Chatterjee, S., Khunti, K., Davies, M.J., 2017. Type 2 diabetes. *Lancet* 389, 2239–2251. [https://doi.org/10.1016/S0140-6736\(17\)30058-2](https://doi.org/10.1016/S0140-6736(17)30058-2).
- Chen, C.Y., Chen, K.H., Liu, C.Y., Huang, S.L., Lin, K.M., 2009. Increased risks of Congenital, Neurologic, and Endocrine disorders associated with Autism in preschool children: cognitive ability differences. *J. Pediatr.* 154. <https://doi.org/10.1016/j.jpeds.2008.09.043>.
- Chen, M.H., Su, T.P., Chen, Y.S., Hsu, J.W., Huang, K.L., Chang, W.H., Chen, T.J., Bai, Y. M., 2013. Comorbidity of allergic and autoimmune diseases in patients with autism spectrum disorder: a nationwide population-based study. *Res. Autism Spectr. Disord.* 7, 205–212. <https://doi.org/10.1016/j.rasd.2012.08.008>.
- Chen, M.-H., Lan, W.-H., Hsu, J.-W., Huang, K.-L., Su, T.-P., Li, C.-T., Lin, W.-C., Tsai, C.-F., Tsai, S.-J., Lee, Y.-C., Chen, Y.-S., Pan, T.-L., Chang, W.-H., Chen, T.-J., Bai, Y.-M., 2016. Risk of developing Type 2 diabetes in adolescents and young adults with Autism spectrum disorder: A nationwide longitudinal study. *Diabetes Care* 39, 788–793. <https://doi.org/10.2337/dc15-1807>.
- Cortese, S., Sun, S., Zhang, J., Sharma, E., Chang, Z., Kuja-Halkola, R., Almqvist, C., Larsson, H., Faraone, S.V., 2018. Association between attention deficit hyperactivity disorder and asthma: a systematic review and meta-analysis and a Swedish population-based study. *Lancet Psychiatry* 5, 717–726. [https://doi.org/10.1016/S2215-0366\(18\)30224-4](https://doi.org/10.1016/S2215-0366(18)30224-4).
- Cortese, S., Solmi, M., Arrondo, G., Cipriani, A., Fusar-Poli, P., Larsson, H., Correll, C., 2020. Association between mental disorders and somatic conditions: protocol for an umbrella review. *Evid. Based Ment. Heal.* 23, 135–139. <https://doi.org/10.1136/ebmental-2020-300158>.
- Croen, L.A., Zerbo, O., Qian, Y., Massolo, M.L., Rich, S., Sidney, S., Kripke, C., 2015. The health status of adults on the autism spectrum. *Autism* 19, 814–823. <https://doi.org/10.1177/1362361315577517>.
- DiMeglio, L.A., Evans-Molina, C., Oram, R.A., 2018. Type 1 diabetes. *Lancet* 391, 2449–2462. [https://doi.org/10.1016/S0140-6736\(18\)31320-5](https://doi.org/10.1016/S0140-6736(18)31320-5).
- Dybdal, D., Tolstrup, J.S., Sildorf, S.M., Boisen, K.A., Svensson, J., Skovgaard, A.M., Teilmann, G.K., 2018. Increasing risk of psychiatric morbidity after childhood onset type 1 diabetes: a population-based cohort study. *Diabetologia* 61, 831–838. <https://doi.org/10.1007/s00125-017-4517-7>.
- Flygare Wallén, E., Ljunggren, G., Carlsson, A.C., Pettersson, D., Wändell, P., 2018. High prevalence of diabetes mellitus, hypertension and obesity among persons with a recorded diagnosis of intellectual disability or autism spectrum disorder. *J. Intellect. Disabil. Res.* 62, 269–280. <https://doi.org/10.1111/jir.12462>.
- Freeman, S.J., Roberts, W., Daneman, D., 2005. Type 1 Diabetes and Autism: Is there a link? *Diabetes Care* 28, 925–926. <https://doi.org/10.2337/diacare.28.4.925>.
- Gurney, J.G., McPheeters, M.L., Davis, M.M., 2006. Parental report of health conditions and health care use among children with and without Autism. *Arch. Pediatr. Adolesc. Med.* 160, 825. <https://doi.org/10.1001/archpedi.160.8.825>.
- Kahathuduwa, C.N., West, B.D., Blume, J., Dharavath, N., Moustaid-Moussa, N., Mastergeorge, A., 2019. The risk of overweight and obesity in children with autism spectrum disorders: a systematic review and meta-analysis. *Obes. Rev.* 20, 1667–1679. <https://doi.org/10.1111/obr.12933>.
- Kohane, I.S., McMurry, A., Weber, G., MacFadden, D., Rappaport, L., Kunkel, L., Bickel, J., Wattanasin, N., Spence, S., Murphy, S., Churchill, S., 2012. The comorbidity burden of children and young adults with autism spectrum disorders. *PLoS One* 7. <https://doi.org/10.1371/journal.pone.0033224>.
- Lai, M.-C., Kasse, C., Besney, R., Bonato, S., Hull, L., Mandy, W., Szatmari, P., Ameis, S. H., 2019. Prevalence of co-occurring mental health diagnoses in the autism population: a systematic review and meta-analysis. *Lancet Psychiatry* 6, 819–829. [https://doi.org/10.1016/S2215-0366\(19\)30289-5](https://doi.org/10.1016/S2215-0366(19)30289-5).
- Li, H., Liu, H., Chen, X., Zhang, J., Tong, G., Sun, Y., 2021. Association of food hypersensitivity in children with the risk of autism spectrum disorder: a meta-analysis. *Eur. J. Pediatr.* 180, 999–1008. <https://doi.org/10.1007/s00431-020-03826-x>.
- Liu, S., Kuja-Halkola, R., Larsson, H., Lichtenstein, P., Ludvigsson, J.F., Svensson, A.M., Gudbjörnsdóttir, S., Tideman, M., Serlachius, E., Butwickia, A., 2021. Poor glycaemic control is associated with increased risk of neurodevelopmental disorders in childhood-onset type 1 diabetes: a population-based cohort study. *Diabetologia* 64, 767–777. <https://doi.org/10.1007/s00125-020-05372-5>.
- Lord, C., Elsabbagh, M., Baird, G., Veenstra-Vanderweele, J., 2018. Autism spectrum disorder. *Lancet* 392, 508–520. [https://doi.org/10.1016/S0140-6736\(18\)31129-2](https://doi.org/10.1016/S0140-6736(18)31129-2).
- Masi, A., Glozier, N., Dale, R., Guastella, A.J., 2017. The immune system, cytokines, and biomarkers in Autism spectrum disorder. *Neurosci. Bull.* 33, 194–204. <https://doi.org/10.1007/s12264-017-0103-8>.
- McDermott, S., Moran, R., Platt, T., Dasari, S., 2007. Prevalence of diabetes in persons with disabilities in primary care. *J. Dev. Phys. Disabil.* 19, 263–271. <https://doi.org/10.1007/s10882-007-9058-4>.
- McManus, S., Meltzer, H., Traolach, B., Bebbington, P., Jenkins, R., 2009. Adult Psychiatric Morbidity in England, 2007: results of a household survey. NHS Information Centre for health and social care.
- Page, M.J., McKenzie, J.E., Bossuyt, P.M., Boutron, I., Hoffmann, T.C., Mulrow, C.D., Shamseer, L., Tetzlaff, J.M., Akl, E.A., Brennan, S.E., Chou, R., Glanville, J., Grimshaw, J.M., Hróbjartsson, A., Lalu, M.M., Li, T., Loder, E.W., Mayo-Wilson, E., McDonald, S., McGuinness, L.A., Stewart, L.A., Thomas, J., Tricco, A.C., Welch, V.A., Whiting, P., Moher, D., 2021. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ*. <https://doi.org/10.1136/bmj.n71> n71.
- Qureshi, I.A., Mehler, M.F., 2013. Towards a 'systems'-level understanding of the nervous system and its disorders. *Trends Neurosci.* 36, 674–684. <https://doi.org/10.1016/j.tins.2013.07.003>.
- Rowland, J., Wilson, C.A., 2021. The association between gestational diabetes and ASD and ADHD: a systematic review and meta-analysis. *Sci. Rep.* 11, 5136. <https://doi.org/10.1038/s41598-021-84573-3>.
- Schott, W., Tao, S., Shea, L., 2021. Co-occurring conditions and racial-ethnic disparities: medicare enrolled adults on the autism spectrum. *Autism Res.* <https://doi.org/10.1002/aur.2644>.
- Shedlock, K., Susi, A., Gorman, G.H., Hisle-Gorman, E., Erdie-Lalena, C.R., Nylund, C.M., 2016. Autism spectrum disorders and metabolic complications of obesity. *J. Pediatr.* 178, 183–187. <https://doi.org/10.1016/j.jpeds.2016.07.055> e1.
- Spann, M.N., Timonen-Soivio, L., Suominen, A., Cheslack-Postava, K., McKeague, I.W., Sourander, A., Brown, A.S., 2019. Proband and familial Autoimmune diseases are associated with proband diagnosis of Autism spectrum disorders. *J. Am. Acad. Child Adolesc. Psychiatry* 58, 496–505. <https://doi.org/10.1016/j.jaac.2018.09.444>.
- Supekar, K., Iyer, T., Menon, V., 2017. The influence of sex and age on prevalence rates of comorbid conditions in autism. *Autism Res.* 10, 778–789. <https://doi.org/10.1002/aur.1741>.
- Tromans, S., Yao, G., Alexander, R., Mukaetova-Ladinska, E., Kiani, R., Al-Uzri, M., Chester, V., Carr, R., Morgan, Z., Vounzoulaki, E., Brugh, T., 2021. The prevalence of diabetes in autistic persons: a systematic review. *Clin. Pract. Epidemiol. Ment. Heal.* 16, 212–225. <https://doi.org/10.2174/1745017902016010212>.
- Tsai, T., Chao, Y., Hsieh, C., Huang, Y., 2020. Association between Atopic Dermatitis and Autism spectrum disorder: a systematic review and meta-analysis. *Acta Derm. Venereol.* 100 <https://doi.org/10.2340/00015555-3501> adv00146.
- Tyler, C.V., Schramm, S.C., Karafa, M., Tang, A.S., Jain, A.K., 2011. Chronic disease risks in young adults with autism spectrum disorder: Forewarned is forearmed. *Am. J. Intellect. Dev. Disabil.* 116, 371–380. <https://doi.org/10.1352/1944-7558-116.5.371>.
- Vohra, R., Madhavan, S., Sambamoorthi, U., 2017. Comorbidity prevalence, healthcare utilization, and expenditures of Medicaid enrolled adults with autism spectrum disorders. *Autism* 21, 995–1009. <https://doi.org/10.1177/1362361316665222>.



- Weir, E., Allison, C., Warrier, V., Baron-Cohen, S., 2021. Increased prevalence of non-communicable physical health conditions among autistic adults. *Autism* 25, 681–694. <https://doi.org/10.1177/1362361320953652>.
- Weiss, J.A., Isaacs, B., Diepstra, H., Wilton, A.S., Brown, H.K., McGarry, C., Lunskey, Y., 2018. Health concerns and health service utilization in a population cohort of young adults with Autism spectrum disorder. *J. Autism Dev. Disord.* 48, 36–44. <https://doi.org/10.1007/s10803-017-3292-0>.
- Wells, G., Shea, B., O'Connell, D., Peterson, J., Welch, V., Losos, M., Tugwell, P., 2000. The Newcastle-Ottawa Scale (NOS) for assessing the quality of non randomised studies in meta-analyses [WWW Document]. URL ([http://www.ohri.ca/program/clinical\\_epidemiology/oxford.asp](http://www.ohri.ca/program/clinical_epidemiology/oxford.asp)) (accessed 9.1.21).
- Wells, P.G., McCallum, G.P., Chen, C.S., Henderson, J.T., Lee, C.J.J., Perstin, J., Preston, T.J., Wiley, M.J., Wong, A.W., 2009. Oxidative stress in developmental origins of disease: teratogenesis, neurodevelopmental deficits, and cancer. *Toxicol. Sci.* 108, 4–18. <https://doi.org/10.1093/toxsci/kfn263>.
- Zaiontz, C., 2021a. Fleiss' Kappa [WWW Document]. Real Stat. Using Excel. URL (<https://www.real-statistics.com/reliability/interrater-reliability/fleiss-kappa/>) (accessed 12.12.21).
- Zaiontz, C., 2021b. Cohen's Kappa [WWW Document]. Real Stat. Using Excel. URL (<https://www.real-statistics.com/reliability/interrater-reliability/cohen-kappa/>) (accessed 12.12.21).
- Zerbo, O., Leong, A., Barcellos, L., Bernal, P., Fireman, B., Croen, L.A., 2015. Immune mediated conditions in autism spectrum disorders. *Brain Behav. Immun.* 46, 232–236. <https://doi.org/10.1016/j.bbi.2015.02.001>.