



Use of concomitant proton pump inhibitors, statins or metformin in patients treated with pembrolizumab for metastatic urothelial carcinoma: data from the ARON-2 retrospective study

Ondřej Fiala · Sebastiano Buti · Hideki Takeshita · Yohei Okada · Francesco Massari · Georgia Anguera Palacios, et al. [full author details at the end of the article]

Received: 4 May 2023 / Accepted: 3 August 2023 / Published online: 7 September 2023
© The Author(s), under exclusive licence to Springer-Verlag GmbH Germany, part of Springer Nature 2023

Abstract

Background Concomitant medications may potentially affect the outcome of cancer patients. In this sub-analysis of the ARON-2 real-world study (NCT05290038), we aimed to assess the impact of concomitant use of proton pump inhibitors (PPI), statins, or metformin on outcome of patients with metastatic urothelial cancer (mUC) receiving second-line pembrolizumab.

Methods We collected data from the hospital medical records of patients with mUC treated with pembrolizumab as second-line therapy at 87 institutions from 22 countries. Patients were assessed for overall survival (OS), progression-free survival (PFS), and overall response rate. We carried out a survival analysis by a Cox regression model.

Results A total of 802 patients were eligible for this retrospective study; the median follow-up time was 15.3 months. PPI users compared to non-users showed inferior PFS (4.5 vs. 7.2 months, $p=0.002$) and OS (8.7 vs. 14.1 months, $p<0.001$). Concomitant PPI use remained a significant predictor of PFS and OS after multivariate Cox analysis. The use of statins or metformin was not associated with response or survival.

Conclusions Our study results suggest a significant prognostic impact of concomitant PPI use in mUC patients receiving pembrolizumab in the real-world context. The mechanism of this interaction warrants further elucidation.

Keywords Urothelial cancer · Proton pump inhibitors · Statins · Metformin · ARON-2 study · Immunotherapy

Introduction

In recent years, the introduction of immune checkpoint inhibitors (ICIs) targeting the programmed cell death protein-1 (PD-1)/ligand-1 (PD-L1) for the treatment of advanced or metastatic urothelial carcinoma (mUC) has improved outcomes and quality of life of these patients [1, 2]. Currently, ICIs are approved in three indications for mUC: (1) first-line treatment in cisplatin-ineligible patients with high PD-L1-expressing tumors or any platinum-ineligible patients regardless of PD-L1 status (atezolizumab, pembrolizumab); (2) second-line treatment following platinum-based chemotherapy or progression within 12 months after neoadjuvant or adjuvant chemotherapy (pembrolizumab, nivolumab,

atezolizumab), and (3) switch maintenance therapy following platinum-based chemotherapy (avelumab) [2–5].

Patients with advanced solid cancers often have other relevant comorbidities, implying a significant number of concomitant medications. Drug–drug interactions as well as potential anticancer effects of commonly used drugs have become an area of increasing interest. Proton pump inhibitors (PPIs), statins, and metformin represent the most commonly prescribed drugs worldwide including cancer patients. Recent studies have reported possible detrimental effects of concomitant medications, including antibiotics, PPIs, and corticosteroids, on the efficacy of ICIs in patients with several malignancies [6–16], including mUC [13–17]. In particular, PPIs are among the most commonly prescribed drugs for patients with cancer to reduce gastrointestinal toxicity associated with certain anticancer drugs [18]. However, the potential risk of their direct and/or indirect interactions with anticancer agents is high in terms of bioavailability, pharmacokinetics, and immunological

Ondřej Fiala and Sebastiano Buti have contributed equally to this work.

Joaquim Bellmunt and Matteo Santoni are co-Senior Authors.

interference, the latter possibly mediated by microbiota [19, 20]. Consequently, the effectiveness of anticancer compounds may be altered by taking concomitant PPI alone or in combination with steroids and/or antibiotics [7, 16, 21]. The impact of PPI intake during ICI immunotherapy for mUC has been suggested very recently by three relatively small studies [13–17]. In contrast to PPIs, the concomitant administration of metformin or statins seems to be associated with favorable outcome in patients with advanced cancer treated with ICIs. Recent data suggest potential synergistic antitumor effects of the concomitant use of ICIs as monotherapy with the oral hypoglycemic agent metformin in patients with cancer and diabetes mellitus, possibly due to the elicitation of multiple cross-mechanisms between cancer metabolism and the host immune system in controlling cancer cell growth [22]. Nevertheless, when ICIs were used in the context of the first-line combination therapy in patients with advanced renal cell carcinoma, the role of concomitant metformin intake was not confirmed [23]. Additionally, concomitant statin exposure in cancer patients treated with ICIs may also lead to favorable outcomes. It has been demonstrated that statins exert a series of biological activities with high potential to enhance the effect of ICIs. Specifically, statins modulate the cancer cell metabolism and are also involved in multiple immune system functions including T-cell signaling, antigen presentation, immune cell migration and cytokine production [23–26]. To our knowledge, there are no data on a possible association between metformin or statin intake and outcome of mUC patients treated with ICIs.

The ARON project has been designed to create a global network to allow uro-oncologists to share and discuss their experiences on the use of immunotherapy and other emerging drugs in patients with genitourinary cancers. Specifically, the ARON-2 study (NCT05290038) was designed to globally collect real-world data on the use of pembrolizumab for mUC. The aim of the present analysis was to assess the association of concomitant use of PPIs, metformin, or statins with patient outcomes in the ARON-2 study population.

Patients and methods

Study design and patient population

We retrospectively analyzed data from patients with a cytologically and/or histologically confirmed mUC, treated with second-line pembrolizumab. The clinical data were collected between January 1, 2016, and October 1, 2022. We retrospectively reviewed anonymized data obtained from the hospital information systems sent by participating centers.

The study protocol "ARON-2" (No. 2022 39) was approved by the Ethical Committee of the Marche Region (Italy) on February 17, 2022, and complied with the International Ethical Guidelines for Biomedical Research, the Declaration of Helsinki, and local laws in each participating center. The informed consent with subsequent analysis of the follow-up data was obtained from all the participants.

Pembrolizumab was administered intravenously as a single agent at the standard approved schedule (200 mg every 3 weeks). The treatment was continued until disease progression, unacceptable toxicity, or patient refusal. None of the patients had received prior ICI therapy. Concomitant PPIs, statins, and metformin were administered orally at individualized doses under the supervision the patients' healthcare providers. Standard follow-up for patients receiving pembrolizumab generally consisted of periodic physical examinations, and laboratory analyses were usually carried out every 3–6 weeks. Computed tomography (CT) or magnetic resonance (MR) was performed at baseline and every 2–4 months thereafter, according to physicians' practice, or when disease progression was clinically suspected.

Study endpoints

The primary endpoint of the ARON-2 study was response, assessed by progression-free survival (PFS), overall survival (OS) and overall response rate (ORR). This sub-analysis evaluates the correlation between the concomitant use of PPIs, metformin or statins, and these outcomes of mUC patients treated with pembrolizumab. PFS was defined as the time from the start of pembrolizumab to progression or death from any cause, whichever occurred first. OS was defined as the time from the start of pembrolizumab until death or lost at follow-up. We considered as censored those patients without progression or death at the last follow-up. The objective response to pembrolizumab was assessed according to RECIST 1.1 principles, and data on complete (CR) or partial responses (PR), stable disease (SD) or progressive disease (PD) were collected and analyzed [27]. ORR was defined as the proportion of patients who achieved CR or PR per RECIST 1.1 criteria.

Statistical analysis

PFS and OS were estimated by using the Kaplan–Meier method with Rothman's 95% confidence intervals (CI) and compared using the log-rank test. The median follow-up was calculated with the Kaplan–Meier method. Univariate and multivariate analyses were performed by using Cox proportional hazards models. The Chi-square test was used to assess potential differences between PPIs, statin or metformin users versus non-users in terms of patient characteristics and ORR. Subsequently, survival analyses focusing

on the impact of PPI use in specific patient subgroups were performed. These subgroups were defined by: gender (males, females), age (< 65 years, ≥ 65 years), smoking (current and former smokers, non-smokers), histology (pure UC, other variants), ECOG-PS (0–1, ≥ 2), primary tumor site (lower urinary tract, upper urinary tract), type of metastatic spread (synchronous metastases = mUC initially diagnosed in stage IV, metachronous metastases = metastatic relapse of initially localized-stage UC), site of distant metastases (lymph node—non-regional, bone, lung, liver), pembrolizumab setting (Cohort A = mUC patients progressed after first-line chemotherapy, Cohort B = metastatic relapse within < 1 years from adjuvant/neoadjuvant therapy). Significance levels were set at a 0.05 value, and all *p* values were two-sided. The statistical analysis was performed by using MedCalc version 19.6.4 (MedCalc Software, Broekstraat 52, 9030 Mariakerke, Belgium).

Results

Baseline characteristics

In total, the study included 802 mUC patients. The baseline patient characteristics are summarized in Table 1. The median follow-up time was 15.3 months (95% CI 14.0–76.0). In 307 (38%) patients, pembrolizumab therapy was ongoing at the time of data cut-off; 430 (54%) patients had died at the time of data cut-off; 535 (67%) patients received pembrolizumab following progression during first-line platinum-based chemotherapy (**Cohort A**) and 267 (33%) after disease recurrence within < 12 months since the completion of adjuvant or neoadjuvant chemotherapy (**Cohort B**). One hundred and sixty-two (32.7%) of the 495 patients with progression on pembrolizumab received further therapies (103 in cohort A and 59 in cohort B).

Concomitant use of PPIs, statins or metformin was reported in 372 (46%), 185 (23%) and 98 (12%) patients, respectively. The use of metformin was more frequent in patients aged ≥ 65 years, while no differences were found in baseline patient characteristics according to the concomitant use of PPIs or statins (Table 1).

Survival analysis

Median PFS and OS for the whole cohort were 6.2 (95% CI 5.1–6.9) and 11.3 months (95% CI 9.5–13.0), respectively. Median PFS and OS for PPI users were 4.5 (95% CI 3.7–6.3) and 8.7 (95% CI 6.9–11.4) months versus 7.2 (95% CI 5.7–9.1) and 14.1 months (95% CI 10.4–16.2) for PPI non-users ($p=0.002$ and $p<0.001$, respectively) (Fig. 1). Median PFS and OS for statin users were 6.9 (95% CI 4.0–9.5) and 10.3 (95% CI 8.0–15.3) months versus 6.0 (95% CI 4.8–6.9)

and 11.3 months (95% CI 9.1–13.4) months for statin non-users ($p=0.715$ and $p=0.999$, respectively) (Fig. 1). Median PFS and OS for metformin users were 7.1 (95% CI 3.7–12.0) and 12.4 (95% CI 7.8–16.0) months versus 6.2 (95% CI 5.0–6.9) and 10.5 (95% CI 9.0–13.3) months for metformin non-users ($p=0.630$ and $p=0.896$, respectively) (Fig. 1). Forest plots summarizing HRs and 95% CIs for OS and PFS according to the assessed comedication are displayed in Fig. 2.

In the Cox multivariate analysis, the use of PPIs remains a significant and independent factor predicting both PFS (HR = 1.28, 95% CI 1.08–1.53, $p=0.006$) and OS (HR = 1.41, 95% CI 1.17–1.71, $p<0.001$). Other independent prognostic factors were synchronous metastatic disease, bone and liver metastases for PFS; and smoking status, synchronous metastatic disease, bone and liver metastases for OS (Table 2).

Subgroup survival analyses according to the concomitant use of PPIs

The association of concomitant use of PPIs with shorter OS was statistically significant in both males ($p=0.004$) and females ($p=0.019$) (Fig. 3); patients aged ≥ 65 years ($p=0.001$) (Fig. 3); current or former smokers ($p<0.001$) (Fig. 3); patients with ECOG-PS 0–1 ($p=0.004$); patients with pure UC histology ($p=0.002$) and those with other histological variants ($p=0.034$) (Fig. 4); patients with UC of lower urinary tract ($p=0.001$) (Fig. 4); patients with metachronous metastatic disease ($p=0.001$) (Fig. 4); patients with non-regional lymph node metastases ($p<0.001$) (Fig. 4); patients with bone metastases ($p=0.012$) (Fig. 4); and patients in cohort A ($p<0.001$) (Fig. 4). The difference in OS according to the use of PPIs was not statistically significant in patients aged < 65 years ($p=0.135$); non-smokers ($p=0.057$); patients with ECOG-PS ≥ 2 ($p=0.150$); patients with upper urinary tract tumors ($p=0.112$); patients with lung metastases ($p=0.179$); patients with liver metastases ($p=0.122$); patients with synchronous metastatic disease ($p=0.067$); and patients in cohort B ($p=0.262$). Similar to OS, the association of concomitant use of PPIs with shorter PFS was statistically significant in both males ($p=0.020$) and females ($p=0.031$) (Fig. 5); patients aged ≥ 65 years ($p=0.013$) and those aged < 65 years ($p=0.05$) (Fig. 5); current or former smokers ($p=0.006$) (Fig. 5); patients with ECOG-PS 0–1 ($p=0.017$); patients with pure UC histology ($p=0.006$) (Fig. 5); patients with UC of lower urinary tract ($p<0.001$) (Fig. 5); patients with metachronous ($p=0.025$) and those with synchronous metastatic disease ($p=0.016$); patients with non-regional lymph node metastases ($p<0.001$) (Fig. 6); and patients in cohort A ($p=0.008$) (Fig. 6). The difference in PFS according to the use of PPIs was not statistically significant in non-smokers ($p=0.12$);

Table 1 Patients' characteristics in the overall study population and stratified by concomitant medications

Patients	Overall 802 (%)	PPI users 372 (%)	PPI non- users 430 (%)	<i>p</i>	Statin users 185 (%)	Statin non- users 617 (%)	<i>P</i>	Metformin users 98 (%)	Metformin non-users 704 (%)	<i>P</i>
<i>Gender</i>										
Male	591 (74)	277 (74)	314 (73)	0.873	145 (78)	446 (72)	0.328	77 (79)	514 (73)	0.322
Female	211 (26)	95 (26)	116 (27)		40 (22)	171 (28)		21 (21)	190 (27)	
Age ≥ 65 years	595 (74)	285 (77)	310 (72)	0.418	151 (81)	444 (72)	0.134	83 (85)	512 (73)	0.038
ECOG-PS ≥ 2	138 (17)	74 (20)	64 (15)	0.353	40 (22)	98 (16)	0.281	13 (13)	125 (18)	0.330
Current or former smokers	521 (65)	253 (68)	268 (62)	0.375	130 (70)	391 (63)	0.296	63 (64)	458 (65)	0.883
<i>Primary tumor location</i>										
Upper urinary tract	214 (27)	108 (29)	106 (25)	0.525	44 (24)	170 (28)	0.520	23 (23)	191 (27)	0.515
Lower urinary tract	588 (73)	264 (71)	324 (75)		141 (76)	447 (72)		75 (77)	513 (73)	
<i>Tumor histology</i>										
Pure urothelial carcinoma	648 (81)	307 (83)	341 (79)	0.472	157 (85)	491 (80)	0.353	87 (89)	561 (80)	0.079
Minor or mixed variants	154 (19)	65 (17)	89 (21)		28 (15)	126 (20)		11 (11)	143 (20)	
Synchronous metastatic disease	246 (31)	113 (30)	133 (31)	0.878	50 (27)	196 (32)	0.439	29 (30)	217 (31)	0.878
<i>Common sites of metastasis</i>										
Lymph nodes (non-regional)	564 (70)	263 (71)	301 (70)	0.877	128 (69)	436 (71)	0.758	73 (74)	491 (70)	0.530
Lung	265 (33)	120 (32)	145 (34)	0.764	64 (35)	201 (33)	0.766	43 (44)	222 (32)	0.081
Bone	227 (28)	114 (31)	113 (26)	0.435	52 (28)	175 (28)	1.000	33 (34)	194 (28)	0.360
Liver	146 (18)	71 (19)	75 (17)	0.714	48 (26)	98 (16)	0.083	22 (22)	124 (18)	0.481
Brain	11 (1)	7 (2)	4 (1)	0.562	5 (3)	6 (1)	0.314	4 (4)	7 (1)	0.175
<i>Pembrolizumab setting</i>										
Cohort A (progressed after first-line chemotherapy)	535 (67)	265 (71)	270 (63)	0.230	115 (62)	420 (68)	0.375	68 (69)	467 (66)	0.651
Cohort B (recurred within < 1 year from adjuvant/ neoadjuvant therapy)	267 (33)	107 (29)	160 (37)		70 (38)	197 (32)		30 (31)	237 (34)	





Fig. 1 Overall survival and progression-free survival according to the use of concomitant proton pump inhibitors (PPIs), statins or metformin

Fig. 2 Forest plots showing hazard ratios (HR) and 95% confidence intervals (CI) for overall survival (OS) and progression-free survival (PFS) according to the use of concomitant proton pump inhibitors (PPIs), statins and metformin



patients with other histological UC variants ($p=0.120$); patients with ECOG-PS ≥ 2 ($p=0.179$); patients with upper urinary tract tumors ($p=0.595$); patients with bone metastases ($p=0.317$); lung metastases ($p=0.625$); patients with liver metastases ($p=0.550$) and those in cohort B ($p=0.150$). The survival data are summarized in detail in Table 3. Forest plots summarizing HRs and 95% CIs for OS and PFS according to the PPI use in all patient subgroups are displayed in Fig. 7.

Objective response

Eighty patients (10%) achieved CR, 168 (21%) PR, 197 (24%) SD and 357 (44%) PD, with an ORR of 31%. The OS was significantly different according to the type of response: NR (95% CI NR–NR), 34.4 months (95% CI 22.4–47.2), 15.6 months (95% CI 12.4–19.4) and 4.3 months (95% CI 3.8–30.4) in patients with CR, PR, SD and PD, respectively ($p<0.001$).

Stratified by concomitant medications, the ORR was 26% in PPI users and 36% in PPI non-users (CR = 8%, PR = 18%, SD = 24%, PD = 50% vs CR = 12%, PR = 24%, SD = 24%, PD = 40%; $p=0.127$). No difference was found in terms of ORR between statin users versus non-users (ORR = 35%, CR = 12%, PR = 23%, SD = 22%, PD = 43% versus ORR = 29%, CR = 9%, PR = 20%, SD = 26%, PD = 45%; $p=0.364$) or between metformin users versus non-users (ORR = 33%, CR = 11%, PR = 22%, SD = 24%, PD = 43%,

vs. ORR = 31%, CR = 10%, PR = 21%, SD = 25%, PD = 44%; $p=0.762$).

Discussion

In the last decade, immunotherapy has increasingly gained a key role in the treatment of several solid tumors and hematological malignancies. In addition, ICIs demonstrated a certain efficacy in various settings in solid tumors: neoadjuvant, adjuvant, first-line or successive lines of therapy. A perfect example of this expanding use of ICIs represents advanced UC. Since the initial second-line strategy approval for patients who progressed on a prior platinum-based chemotherapy, the use of ICIs in mUC has been quickly extended to the first-line setting in PD-L1-positive cisplatin-ineligible patients or any platinum-ineligible patients regardless of PD-L1 expression as well as to the switch maintenance therapy in platinum responders [28].

As we witness this indisputable therapeutic revolution, an emerging and important issue that remains to be solved is whether the concomitant administration of other drugs may impair or eventually enhance the efficacy of ICIs. This is a crucial question also considering that UC mostly affects elderly patients who may present other disorders and use several concomitant medications.

In our study, the concomitant use of PPIs was significantly associated with both shorter PFS and OS in mUC

Table 2 Univariate and multivariate Cox analyses for overall survival (OS) and progression-free survival (PFS)

Overall survival	Univariate cox regression		Multivariate cox regression	
	HR (95% CI)	<i>p</i> value	HR (95% CI)	<i>p</i> value
Gender (females vs. males)	1.22 (0.99–1.50)	0.062		
Age (≥ 65 years vs. < 65 years)	1.07 (0.86–1.33)	0.530		
Smoking (smokers vs. non-smokers)	0.81 (0.66–0.98)	0.030	0.78 (0.64–0.95)	0.012
Histology (mixed vs. pure UC)	1.04 (0.81–1.32)	0.775		
Upper vs. lower urinary tract	1.10 (0.90–1.36)	0.353		
Synchronous metastatic disease (yes vs. no)	1.32 (1.08–1.61)	0.007	1.33 (1.09–1.63)	0.005
Lymph node metastases (Y vs. N)	0.86 (0.70–1.06)	0.151		
Bone metastases (Y vs. N)	1.53 (1.25–1.88)	<0.001	1.51 (1.22–1.84)	<0.001
Liver metastases (Y vs. N)	1.46 (1.16–1.84)	0.001	1.39 (1.10–1.75)	0.006
Proton pump inhibitors (Y vs. N)	1.41 (1.17–1.71)	<0.001	1.41 (1.17–1.70)	<0.001
Statins (Y vs. N)	1.00 (0.80–1.25)	0.999		
Metformin (Y vs. N)	0.98 (0.73–1.31)	0.896		
Progression-free survival	Univariate cox regression		Multivariate cox regression	
	HR (95% CI)	<i>p</i> value	HR (95% CI)	
Gender (females vs. males)	0.99 (0.81–1.22)	0.957		
Age (≥ 65 years vs. < 65 years)	0.93 (0.76–1.14)	0.485		
Smoking (smokers vs. no-smokers)	0.93 (0.77–1.11)	0.409		
Histology (mixed vs. pure UC)	1.03 (0.83–1.29)	0.771		
Upper vs. lower urinary tract	1.04 (0.86–1.27)	0.681		
Synchronous metastatic disease (yes vs. no)	1.28 (1.06–1.55)	0.009	1.24 (1.02–1.49)	0.027
Lymph node metastases (Y vs. N)	0.86 (0.71–1.04)	0.130		
Bone metastases (Y vs. N)	1.48 (1.22–1.79)	<0.001	1.42 (1.17–1.71)	<0.001
Liver metastases (Y vs. N)	1.51 (1.22–1.86)	<0.001	1.44 (1.17–1.79)	<0.001
Proton pump inhibitors (Y vs. N)	1.32 (1.10–1.58)	0.002	1.28 (1.08–1.53)	0.006
Statins (Y vs. N)	0.96 (0.78–1.19)	0.716		
Metformin (Y vs. N)	0.94 (0.71–1.23)	0.935		

ECOG-PS = Eastern Cooperative Oncology Group-Performance Status; UC = Urothelial Carcinoma

patients receiving pembrolizumab therapy. Furthermore, its independent adverse prognostic role was confirmed in the multivariate analysis. The use of statins or metformin did not affect the response or survival outcomes of these patients.

PPIs, statins and metformin represent the most commonly prescribed medications in the global population; however, their potential to affect the efficacy of ICIs in cancer patients has not been fully elucidated.

The ability of PPIs to influence the efficacy of ICIs may be mainly related to the modifications induced by PPIs in gut microbiome, which plays a crucial role in patients treated with ICIs for solid tumors [29]. It has been shown that PPI use can induce marked alterations to the intestinal microbiome including changes in composition, reduction of the alpha diversity, small intestinal bacterial overgrowth and manifestation of oral bacteria in more distal parts of the intestine [30–32]. In this regard, Imhann et al. [33] observed that PPI use is associated with the presence of multiple oral bacteria over-represented in the fecal microbiome, with an

overall significant increase in bacteria, including *Enterococcus*, *Streptococcus*, *Staphylococcus* and *Escherichia coli*. There has been an accumulating body of evidence that the concomitant use of PPIs could have detrimental effect on ICI therapy in various cancer types [16–18]. In particular, this issue has been underexplored in the field of mUC. Several retrospective studies have been reported recently. However, their crucial limitation is based on a relatively small number of included patients, especially those of PPI users, potentially introducing a significant bias. Thus, there has been a lack for solid data from large studies. The association of shorter PFS and OS with concomitant use of PPIs (HR 1.70, 95% CI 1.23–2.35, $p=0.001$ and HR 2.02, 95% CI 1.28–3.18, $p=0.003$, respectively) in a cohort of 227 mUC patients treated with pembrolizumab has been found in a retrospective study conducted by Fukuokaya et al. [13]. Similar data were reported in two other studies conducted by Tomisaki et al. and Okuyama et al. including 40 and 155 patients, respectively. These results are in agreement




Fig. 3 Overall survival according to concomitant use of proton pump inhibitors (PPIs) stratified by sex, smoking status and age


with our large study including 802 patients (372 PPI users). Aside from mUC patients treated with pembrolizumab, similar findings were found in a post hoc analysis of IMvigor210 and IMvigor211 clinical trials conducted by Hopkins et al. [9]. They found that PPI use was associated with shorter PFS and OS (HR 1.38, 95% CI 1.18–1.62, $p < 0.001$ and HR 1.52, 95% CI 1.27–1.83, $p < 0.001$, respectively) in advanced-stage UC patients treated with atezolizumab, while there was no association in those treated with chemotherapy [10].

On the other hand, a study by Kunimitsu et al. including 79 patients failed to show such a prognostic role of the use of PPIs in advanced-stage UC when treated with pembrolizumab [15]. Nonetheless, it is another study fundamentally limited by a small cohort size.

Notably, our data show that the use of PPIs was significantly associated with poor outcome only in patients with primary tumors of the lower urinary tract but not in those

with primary tumors of the upper urinary tract. This may be explained by a relatively small number and the overall poor prognosis of mUC patients with tumors originating in the upper urinary tract and their generally reduced benefit from ICIs [34] and a distinct genomic background of this specific subgroup [35]. Similar results have been recently reported by Taguchi et al., who successfully validated the prognostic capability of the drug score counted from the baseline use of antibiotics, PPIs and corticosteroids in advanced-stage mUC patients treated with pembrolizumab. In agreement with our study, they found that drug score was significantly correlated with survival outcomes in patients with primary tumors of the lower urinary tract but not in those with primary tumors of the upper urinary tract [36]. However, further elucidation of this issue is needed. We found no significant correlation between the PPI use and ORR, indirectly indicating that PPIs mainly impact duration of response (DR), rather

Fig. 4 Overall survival according to concomitant use of proton pump inhibitors (PPIs) stratified by tumor histology, primary tumor site, metachronous disease, lymph node or bone metastases and pembrolizumab setting (second-line therapy after progression on first-line platinum-based chemotherapy, Cohort A)



than objective response to therapy. However, DR was not assessed specifically in the present study.

Anticancer properties of statins such as the inhibition of tumor cell growth, invasion and metastatic potential have

been suggested in various experimental studies [37–39]. Notwithstanding, clinical studies evaluating the impact of statin use in cancer patients including those with UC have been conducted with inconclusive results. Moreover, there




Fig. 5 Progression-free survival according to concomitant use of proton pump inhibitors (PPIs) stratified by sex, age, smoking status, tumor histology and primary tumor site




Fig. 6 Progression-free survival according to concomitant use of proton pump inhibitors (PPIs) stratified by synchronous metastatic disease, lymph node metastases and pembrolizumab setting (second-line therapy after progression on first-line platinum-based chemotherapy, Cohort A)

are no data on the role of the statin use in mUC patients treated with ICIs. Ferro et al. suggested that statins may have a beneficial effect on recurrence rates in patients with high-grade non-muscle invasive urothelial bladder cancer in a multicenter study including 1510 patients [40]. Oppositely, another recent retrospective study by Haimerl et al. found no impact of statin use on bladder cancer recurrence or survival in a cohort of 972 patients [41]. In our study, we found no impact of the concomitant statin use in mUC patients treated with pembrolizumab.

Metformin represents another commonly used drug with suggested anticancer activity, which is related to both direct effects on cancer cells based on inhibition of cancer-related signaling pathways and indirect effects on the host based on lowering blood glucose and insulin as well as anti-inflammatory effects [42, 43]. Although several retrospective studies show the association of metformin use with favorable

prognosis of patients with cancer, there are no data on its role in patients with mUC treated with ICIs. Our data show no impact of the concomitant metformin use in mUC patients treated with pembrolizumab.



Our study presents several limitations, mainly due to its retrospective nature. A centralized review of radiological imaging was not performed. The dosage and duration of the investigated comedication exposure could not be assessed from the available data sources. Thus, we were not able to investigate a difference between chronic and temporary use after pembrolizumab treatment initiation. Furthermore, we had no available data on other concomitant medications (i.e. steroids, antibiotics) or patients' comorbidities that could affect the efficacy of pembrolizumab. Consequently, our results should be interpreted with caution and are in need of a further prospective validation. On the other hand, the major strength of our study is based on a large patient

Table 3 Summary of patient survival data according to the specific subgroups

	Median OS (95% CI)		<i>p</i> value	Median PFS (95% CI)		<i>p</i> value
	PPI users	PPI non-users		PPI users	PPI non-users	
Whole cohort	8.7 months (6.9–11.4)	14.1 months (10.4–16.2)	<i>p</i> < 0.001	4.5 months (3.7–6.3)	7.2 months (5.7–9.1)	<i>p</i> = 0.002
<i>Gender</i>						
Males	10.0 months (7.2–13.0)	15.1 months (11.1–17.0)	<i>p</i> = 0.004	5.0 months (3.8–6.4)	6.9 months (5.6–9.1)	<i>p</i> = 0.020
Females	6.3 months (4.6–46.4)	11.3 months (7.5–18.6)	<i>p</i> = 0.019	4.0 months (3.2–44.0)	8.3 months (4.6–12.4)	<i>p</i> = 0.031
<i>Age</i>						
< 65 years	9.7 months (5.1–16.8)	12.6 months (9.2–22.4)	<i>p</i> = 0.135	4.2 months (3.4–44.0)	6.9 months (4.7–9.5)	<i>p</i> = 0.05
≥ 65 years	8.7 months (6.4–11.4)	14.1 months (10.0–17.0)	<i>p</i> = 0.001	4.6 months (3.6–6.4)	7.5 months (5.6–10.0)	<i>p</i> = 0.013
<i>Smoking</i>						
Current or former smokers	10.0 months (6.9–12.4)	16.0 months (12.4–22.4)	<i>p</i> < 0.001	4.5 months (3.7–6.9)	7.5 months (6.1–11.4)	<i>p</i> = 0.006
Non-smokers	7.4 months (4.6–12.2)	10.2 months (7.5–14.1)	<i>p</i> = 0.057	4.5 months (3.4–6.6)	6.0 months (4.2–9.5)	<i>p</i> = 0.12
<i>Histology</i>						
Pure UC	8.9 months (7.0–11.5)	14.3 months (10.4–16.5)	<i>p</i> = 0.002	5.3 months (3.7–6.6)	7.2 months (5.7–9.4)	<i>p</i> = 0.006
Other variants	6.4 months (3.6–46.4)	12.4 months (8.3–23.4)	<i>p</i> = 0.034	3.9 months (3.2–44.0)	7.0 months (4.7–20.9)	<i>p</i> = 0.120
<i>ECOG-PS</i>						
0–1	11.5 months (8.9–14.9)	16.2 months (13.3–22.2)	<i>p</i> = 0.004	6.4 months (4.5–7.3)	8.5 months (6.7–10.4)	<i>p</i> = 0.017
≥ 2	3.2 months (2.6–4.1)	5.1 months (4.1–6.6)	<i>p</i> = 0.150	2.6 months (1.9–3.3)	3.5 months (2.5–4.8)	<i>p</i> = 0.179
<i>Primary tumor site</i>						
Lower urinary tract	9.7 months (7–11.7)	14.6 months (11.1–17.5)	<i>p</i> = 0.001	4.5 months (3.7–6.3)	7.7 months (6.2–9.9)	<i>p</i> < 0.001
Upper urinary tract	11.8 months (8.3–16.2)	7.2 months (4.8–12.1)	<i>p</i> = 0.112	5.3 months (3.4–8.0)	5.4 months (4.1–9.5)	<i>p</i> = 0.595
<i>Type of metastatic spread</i>						
Synchronous metastases	6.2 months (4.0–9.7)	10.0 months (7.4–47.2)	<i>p</i> = 0.067	3.6 months (2.6–5.8)	6.6 months (3.9–9.5)	<i>p</i> = 0.016
Metachronous metastases	10.2 months (7.8–13.0)	15.8 months (12.5–22.2)	<i>p</i> = 0.001	6.1 months (4.0–7.2)	7.7 months (5.8–9.9)	<i>p</i> = 0.025
<i>Site of distant metastases</i>						
Lymph node (non-regional)	8.7 months (6.4–11.7)	15.4 months (12.5–19.0)	<i>p</i> < 0.001	4.5 months (3.5–6.3)	8.6 months (6.4–12.4)	<i>p</i> < 0.001
Bone	5.8 months (3.9–7.0)	6.8 months (5.5–16.2)	<i>p</i> = 0.012	3.3 months (2.9–4.6)	3.8 months (3.2–5.6)	<i>p</i> = 0.317
Lung	8.1 months (5.8–11.7)	11.3 months (7.4–15.5)	<i>p</i> = 0.179	6.2 months (3.5–7.2)	5.5 months (4.1–7.7)	<i>p</i> = 0.625
Liver	4.5 months (3.6–8.1)	10.0 months (6.7–16.0)	<i>p</i> = 0.122	3.6 months (2.5–4.5)	3.9 months (3.3–4.9)	<i>p</i> = 0.550
<i>Pembrolizumab setting</i>						
Cohort A (progressed after first-line chemotherapy)	7.2 months (5.9–9.7)	12.6 months (9.2–15.8)	<i>p</i> < 0.001	4.3 months (3.5–6.1)	6.4 months (4.8–8.4)	<i>p</i> = 0.008
Cohort B (recurred within < 1 years from adjuvant/neoadjuvant therapy)	16.0 months (8.8–21.2)	17.0 months (10.3–45.9)	<i>p</i> = 0.262	6.9 months (3.5–14.9)	8.6 months (6.2–25.4)	<i>p</i> = 0.150

ECOG-PS = Eastern Cooperative Oncology Group-Performance Status; UC = Urothelial Carcinoma, CI = Confidence interval, y = years; statistically significant *p* values are in bold

Fig. 7 Forest plots showing hazard ratios (HR) and 95% confidence intervals (CI) for overall survival (OS) and progression-free survival (PFS) according to the PPI use in all patient subgroups



population, which allowed us to perform a detailed subgroup analysis.

In conclusion, our data show that the concomitant use of PPIs may adversely affect the outcome of mUC patients treated with second-line pembrolizumab. Further studies investigating the biological and immunological background of this interaction are warranted in order to optimize the outcome of patients receiving immunotherapy in this setting.

Acknowledgements The authors would like to thank all patients voluntarily taking part in the study.

Author contributions Manuscript: Use of concomitant proton pump inhibitors, statins, or metformin in patients treated with pembrolizumab

for advanced urothelial carcinoma: Data from the ARON-2 retrospective study. Fiala et al. O. Fiala: Investigation, Writing - Original Draft M. Santoni: Conceptualization, Methodology, Investigation, Formal analysis, Writing - Original Draft S. Buti: Conceptualization, Methodology, Investigation, Writing - Original Draft All authors: Investigation

Funding None to declare.

Declarations

Conflict of interest O. Fiala received honoraria from Roche, Janssen, GSK and Pfizer for consultations and lectures unrelated to this project. S. Buti received honoraria as speaker at scientific events and advisory role by BMS, Pfizer, MSD, Ipsen, AstraZeneca, Merck, all unrelated to this project. M. Santoni has received research support and honoraria from Janssen, Bristol Myers Squibb, Ipsen, MSD, Astellas and

Bayer, all unrelated to this project. R. Kanesvaran has received fees for speaker bureau and advisory board activities from the following companies; Pfizer, MSD, BMS, Eisai, Ipsen, Johnson and Johnson, Merck, Amgen, Astellas and Bayer, all unrelated to this project. E. Grande has received honoraria for speaker engagements, advisory roles or funding of continuous medical education from Adacap, AMGEN, Angelini, Astellas, Astra Zeneca, Bayer, Blueprint, Bristol Myers Squibb, Caris Life Sciences, Celgene, Clovis-Oncology, Eisai, Eusa Pharma, Genetracer, Guardant Health, HRA-Pharma, IPSEN, ITM-Radiopharma, Janssen, Lexicon, Lilly, Merck KGaA, MSD, Nanostring Technologies, Natera, Novartis, ONCODNA (Biosequence), Palex, Pharmamar, Pierre Fabre, Pfizer, Roche, Sanofi-Genzyme, Servier, Taiho, and Thermo Fisher Scientific and has received research grants from Pfizer, Astra Zeneca, Astellas, and Lexicon Pharmaceuticals, all unrelated to this project. F. S. M. Monteiro has received research support from Janssen, Merck Sharp Dome and honoraria from Janssen, Ipsen, Bristol Myers Squibb and Merck Sharp Dome, all unrelated to this project. C. Porta has received honoraria from Angelini Pharma, Astra Zeneca, BMS, Eisai, General Electric, Ipsen and MSD and acted as a Protocol Steering Committee Member for BMS, Eisai and MSD, all unrelated to this project. Z. Myint has received research support from Merck unrelated to this project. J. Molina-Cerrillo declares consultant, advisory or speaker roles for IPSEN, Roche, Pfizer, Sanofi, Janssen, and BMS and has received research grants from Pfizer, IPSEN and Roche, all unrelated to this project. P. Giannatempo has received research support from Ipsen, Astra Zeneca, MSD and honoraria for speaker engagements, advisory roles from Astellas, MSD, Janssen, Pfizer, all unrelated to this project. E. T. Lam has received institutional research funding from Merck Sharp & Dohme Corp., a subsidiary of Merck & Co., Inc., Kenilworth, NJ, USA, Bristol-Myers Squibb, Pfizer, and F. Hoffmann-La Roche Ltd. The other authors declare to have no conflicts of interest.

References

- Taguchi S, Kawai T, Nakagawa T, Miyakawa J, Kishitani K, Sugimoto K et al (2022) Improved survival in real-world patients with advanced urothelial carcinoma: a multicenter propensity score-matched cohort study comparing a period before the introduction of pembrolizumab (2003–2011) and a more recent period (2016–2020). *Int J Urol* 29(12):1462–1469. <https://doi.org/10.1111/iju.15014>
- Bellmunt J, de Wit R, Vaughn DJ, Fradet Y, Lee JL, Fong L et al (2017) KEYNOTE-045 investigators. Pembrolizumab as second-line therapy for advanced urothelial carcinoma. *N Engl J Med* 376(11):1015–1026. <https://doi.org/10.1056/NEJMoa1613683>
- Peters S, Dziadziuszko R, Morabito A, Felip E, Gadgeel SM, Cheema P et al (2018) Atezolizumab versus chemotherapy in patients with platinum treated locally advanced or metastatic urothelial carcinoma (IMvigor211): a multicentre, open-label, phase 3 randomised controlled trial. *Lancet* 28(9):1831–1839. <https://doi.org/10.1038/s41591-022-01933-w>
- Sharma P, Retz M, Siefker-Radtke A, Baron A, Necchi A, Bedke J et al (2017) Nivolumab in metastatic urothelial carcinoma after platinum therapy (CheckMate 275): a multicenter, single-arm, phase 2 trial. *Lancet Oncol* 18(3):312–322. [https://doi.org/10.1016/S1470-2045\(17\)30065-7](https://doi.org/10.1016/S1470-2045(17)30065-7)
- Patel MR, Ellerton J, Infante JR, Agrawal M, Gordon M, Aljumailyet R et al (2018) Avelumab in metastatic urothelial carcinoma after platinum failure (JAVELIN Solid Tumor): pooled results from two expansion cohorts of an open-label, phase 1 trial. *Lancet Oncol* 19(1):51–64. [https://doi.org/10.1016/S1470-2045\(17\)30900-2](https://doi.org/10.1016/S1470-2045(17)30900-2)
- Buti S, Bersanelli M, Perrone F, Tiseo M, Tucci M, Adamo V et al (2021) Effect of concomitant medications with immune-modulatory properties on the outcomes of patients with advanced cancer treated with immune checkpoint inhibitors: development and validation of a novel prognostic index. *Eur J Cancer* 142:18–28. <https://doi.org/10.1016/j.ejca.2020.09.033>
- Buti S, Bersanelli M, Perrone F, Bracarda S, Di Maio M, Giusti R et al (2021) Predictive ability of a drug-based score in patients with advanced non-small-cell lung cancer receiving first-line immunotherapy. *Eur J Cancer* 150:224–231. <https://doi.org/10.1016/j.ejca.2021.03.041>
- Takada K, Buti S, Bersanelli M, Shimokawa M, Takamori S, Matsubara T et al (2022) Antibiotic-dependent effect of probiotics in patients with non-small cell lung cancer treated with PD-1 checkpoint blockade. *Eur J Cancer* 172:199–208. <https://doi.org/10.1016/j.ejca.2022.06.002>
- Hopkins AM, Kichenadasse G, Karapetis CS, Rowland A, Sorich MJ (2020) Concomitant antibiotic use and survival in urothelial carcinoma treated with atezolizumab. *Eur Urol* 78(4):540–543. <https://doi.org/10.1016/j.eururo.2020.06.061>
- Hopkins AM, Kichenadasse G, Karapetis CS, Rowland A, Sorich MJ (2020) Concomitant proton pump inhibitor use and survival in urothelial carcinoma treated with atezolizumab. *Clin Cancer Res* 26:5487–5493
- Ruiz-Bañobre J M-DA, Fernández-Calvo O, Fernández-Núñez N, Medina-Colmenero A, Santomé L et al (2021) Rethinking prognostic factors in locally advanced or metastatic urothelial carcinoma in the immune checkpoint blockade era: a multicenter retrospective study. *ESMO Open* 6(2):100090. <https://doi.org/10.1016/j.esmoop.2021.100090>
- Ishiyama Y, Kondo T, Nemoto Y, Kobari Y, Ishihara H, Tachibana H et al (2021) Antibiotic use and survival of patients receiving pembrolizumab for chemotherapy-resistant metastatic urothelial carcinoma. *Urol Oncol* 39(12):834.e21–834.e28. <https://doi.org/10.1016/j.urolonc.2021.05.033>
- Fukuokaya W, Kimura T, Komura K, Uchimoto T, Nishimura K, Yanagisawa T et al (2022) Effectiveness of pembrolizumab in patients with urothelial carcinoma receiving proton pump inhibitors. *Urol Oncol* 40(7):346.e1–346.e8. <https://doi.org/10.1016/j.urolonc.2022.02.020>
- Tomisaki I, Harada M, Minato A, Nagata Y, Kimuro R, Higashijima K et al (2022) Impact of the use of proton pump inhibitors on pembrolizumab effectiveness for advanced urothelial carcinoma. *Anticancer Res* 42(3):1629–1634. <https://doi.org/10.21873/anticancer.15638>
- Kunimitsu Y, Morio K, Hirata S, Yamamoto K, Yamamoto K, Omura T, Hara T et al (2022) Effects of proton pump inhibitors on survival outcomes in patients with metastatic or unresectable urothelial carcinoma treated with pembrolizumab. *Biol Pharm Bull* 45(5):590–595. <https://doi.org/10.1248/bpb.b21-00939>
- Okuyama Y, Hatakeyama S, Numakura K, Narita T, Tanaka T, Miura Y et al (2021) Prognostic impact of proton pump inhibitors for immunotherapy in advanced urothelial carcinoma. *BJUI Compass* 3:154–161. <https://doi.org/10.1002/bco.2.118>
- Rizzo A, Santoni M, Mollica V, Ricci AD, Calabrò C, Cusmai A et al (2022) The impact of concomitant proton pump inhibitors on immunotherapy efficacy among patients with urothelial carcinoma: a meta-analysis. *J Pers Med* 12(5):842. <https://doi.org/10.3390/jpm12050842>
- Triadafilopoulos G, Roorda AK, Akiyama J (2013) Indications and safety of proton pump inhibitor drug use in patients with cancer. *Expert Opin Drug Saf* 12(5):659–672. <https://doi.org/10.1517/14740338.2013.797961>
- Chalabi M, Cardona A, Nagarkar DR, Scala AD, Gandara DR, Rittmeyer A et al (2020) Efficacy of chemotherapy and atezolizumab in patients with non-small-cell lung cancer receiving

- antibiotics and proton pump inhibitors: pooled post hoc analyses of the OAK and POPLAR trials. *Ann Oncol* 31(4):525–531. <https://doi.org/10.1016/j.annonc.2020.01.006>
20. Jackson MA, Goodrich JK, Maxan ME, Freedberg DE, Abrams JA, Poole AC et al (2016) Proton pump inhibitors alter the composition of the gut microbiota. *Gut* 65(5):749–756. <https://doi.org/10.1136/gutjnl-2015-310861>
 21. Deng R, Zhang H, Li Y, Shi Y (2022) Effect of antacid use on immune checkpoint inhibitors in advanced solid cancer patients: a systematic review and meta-analysis. *J Immunother* 46(2):43–55. <https://doi.org/10.1097/CJI.0000000000000442>
 22. Ciccarese C, Iacovelli R, Buti S, Primi F, Astore S, Massari F et al (2022) Concurrent nivolumab and metformin in diabetic cancer patients: is it safe and more active? *Anticancer Res* 42(3):1487–1493. <https://doi.org/10.21873/anticancer.15620>
 23. Santoni M, Molina-Cerrillo J, Myint ZW, Massari F, Buchler T, Buti S et al (2022) Concomitant use of statins, metformin, or proton pump inhibitors in patients with advanced renal cell carcinoma treated with first-line combination therapies. *Target Oncol* 17(5):571–581. <https://doi.org/10.1007/s11523-022-00907-9>
 24. Santoni M, Massari F, Matrana MR, Basso U, De Giorgi U, Aurilio G et al (2022) Statin use improves the efficacy of nivolumab in patients with advanced renal cell carcinoma. *Eur J Cancer* 172:191–198. <https://doi.org/10.1016/j.ejca.2022.04.035>
 25. Perrone F, Minari R, Bersanelli M, Bordi P, Tiseo M, Favari E et al (2020) The prognostic role of high blood cholesterol in advanced cancer patients treated with immune checkpoint inhibitors. *J Immunother* 43(6):196–203. <https://doi.org/10.1097/CJI.0000000000000321>
 26. Cantini L, Pecci F, Hurkmans DP, Belderbos RA, Lanese A, Copparoni C et al (2021) High-intensity statins are associated with improved clinical activity of PD-1 inhibitors in malignant pleural mesothelioma and advanced non-small cell lung cancer patients. *Eur J Cancer* 144:41–48. <https://doi.org/10.1016/j.ejca.2020.10.031>
 27. Schwartz LH, Litière S, de Vries E, Ford R, Gwyther S, Mandrekas S et al (2016) RECIST 1.1-update and clarification: from the RECIST committee. *Eur J Cancer* 62:132–137. <https://doi.org/10.1016/j.ejca.2016.03.081>
 28. Mollica V, Rizzo A, Montironi R, Cheng L, Giunchi F, Schiavina R et al (2020) Current strategies and novel therapeutic approaches for metastatic urothelial carcinoma. *Cancers (Basel)* 12(6):1449. <https://doi.org/10.3390/cancers12061449>
 29. Routy B, Le Chatelier E, Derosa L, Duong CPM, Alou MT, Daillère R et al (2018) Gut microbiome influences efficacy of PD-1-based immunotherapy against epithelial tumors. *Science* 359(6371):91–97. <https://doi.org/10.1126/science.aan3706>
 30. Seto CT, Jeraldo P, Orenstein R, Chia N, DiBaise JK (2014) Prolonged use of a proton pump inhibitor reduces microbial diversity: implications for *Clostridium difficile* susceptibility. *Microbiome* 2:42. <https://doi.org/10.1186/2049-2618-2-42>
 31. Clooney AG, Bernstein CN, Leslie WD, Vagianos K, Sargent M, Laserna-Mendieta EJ et al (2016) A comparison of the gut microbiome between long-term users and non-users of proton pump inhibitors. *Aliment Pharmacol Ther* 43(9):974–984. <https://doi.org/10.1111/apt.13568>
 32. Tsuda A, Suda W, Morita H, Takanashi K, Takagi A, Koga Y et al (2015) Influence of proton-pump inhibitors on the luminal microbiota in the gastrointestinal tract. *Clin Transl Gastroenterol* 6(6):e89. <https://doi.org/10.1038/ctg.2015.20>
 33. Imhann F, Bonder MJ, Vich Vila A, Fu J, Mujagic Z, Vork L et al (2016) Proton pump inhibitors affect the gut microbiome. *Gut* 65(5):740–748. <https://doi.org/10.1136/gutjnl-2015-310376>
 34. Thouvenin J, Martínez Chanzá N, Alhalabi O, Lang H, Tannir NM, Barthélémy P et al (2021) Efficacy of immune checkpoint inhibitors in upper tract urothelial carcinomas: current knowledge and future directions. *Cancers (Basel)* 13(17):4341. <https://doi.org/10.3390/cancers13174341>
 35. Su X, Lu X, Bazal SK, Compérat E, Mouawad R, Yao H et al (2021) Comprehensive integrative profiling of upper tract urothelial carcinomas. *Genome Biol* 22(1):7. <https://doi.org/10.1186/s13059-020-02230-w>
 36. Taguchi S, Kawai T, Buti S, Bersanelli M, Uemura Y, Kishitani K, et al (2023) Validation of a drug-based score in advanced urothelial carcinoma treated with pembrolizumab. <https://doi.org/10.2217/imt-2023-0028>.
 37. Chan KK, Oza AM, Siu LL (2003) The statins as anticancer agents. *Clin Cancer Res* 9:10–19
 38. Keyomarsi K, Sandoval L, Band V (2003) Synchronization of tumor and normal cells from G1 to multiple cell cycles by lovastatin. *Cancer Res* 9:10–19
 39. Wong WW, Dimitroulakos J, Minden MD, Penn LZ (2002) HMG-CoA reductase inhibitors and the malignant cell: the statin family of drugs as triggers of tumor-specific apoptosis. *Leukemia* 16:508–519
 40. Ferro M, Marchioni M, Lucarelli G, Vartolomei MD, Soria F, Terracciano D et al (2021) Association of statin use and oncological outcomes in patients with first diagnosis of T1 high grade non-muscle invasive urothelial bladder cancer: results from a multicenter study. *Minerva Urol Nephrol* 73(6):796–802. <https://doi.org/10.23736/S2724-6051.20.04076-X>
 41. Haimerl L, Strobach D, Mannell H, Stief CG, Buchner A, Karl A et al (2022) Retrospective evaluation of the impact of non-oncologic chronic drug therapy on the survival in patients with bladder cancer. *Int J Clin Pharm* 44(2):339–347. <https://doi.org/10.1007/s11096-021-01343-x>
 42. Samuel SM, Varghese E, Kubatka P, Triggler CR, Büsselberg D et al (2019) Metformin: the answer to cancer in a flower? Current knowledge and future prospects of metformin as an anti-cancer agent in breast cancer. *Biomolecules* 9(12):846. <https://doi.org/10.3390/biom9120846>
 43. Chen YC, Li H, Wang J (2020) Mechanisms of metformin inhibiting cancer invasion and migration. *Am J Transl Res* 12(9):4885–4901

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Springer Nature or its licensor (e.g. a society or other partner) holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.

Authors and Affiliations

Ondřej Fiala^{1,2} · Sebastiano Buti^{3,4} · Hideki Takeshita⁵ · Yohei Okada⁵ · Francesco Massari⁶ · Georgia Anguera Palacios⁷ · Michele Dionese⁸ · Sarah Scagliarini⁹ · Thomas Büttner¹⁰ · Giuseppe Fornarini¹¹ · Zin W. Myint¹² · Luca Galli¹³ · Vinicius Carrera Souza^{14,15} · Renate Pichler¹⁶ · Ugo De Giorgi¹⁷ · María Natalia Gandur Quiroga^{15,18} · Danielle Gilbert¹⁹ · Lazar Popovic²⁰ · Enrique Grande²¹ · Giulia Mammone²² · Rossana Berardi²³ · Simon J. Crabb²⁴ · Javier Molina-Cerrillo²⁵ · Marcelo Freitas^{15,26} · Murilo Luz^{15,27} · Roberto Iacovelli²⁸ · Fabio Calabrò²⁹ · Deniz Tural³⁰ · Francesco Atzori³¹ · Zsófia Küronya³² · Rita Chiari³³ · Saul Campos^{15,34} · Orazio Caffo³⁵ · André P. Fay^{15,36} · Jakub Kucharz³⁷ · Paolo Andrea Zucali^{38,39} · José Augusto Rinck^{15,40} · Annalisa Zeppellini⁴¹ · Diogo Assed Bastos^{15,42} · Gaetano Aurilio⁴³ · Augusto Mota^{15,44} · Karine Trindade^{15,45} · Cinzia Ortega⁴⁶ · Juan Pablo Sade⁴⁷ · Mimma Rizzo⁴⁸ · Nuno Vau⁴⁹ · Patrizia Giannatempo⁵⁰ · Allan Barillas^{15,51} · Fernando Sabino Marques Monteiro^{15,52} · Breno Dauster^{15,53} · Carlo Cattrini⁵⁴ · Lucas Nogueira^{15,55} · Roni de Carvalho Fernandes⁵⁶ · Emmanuel Seront⁵⁷ · Luís Garcia Aceituno^{15,58} · Francesco Grillone⁵⁹ · Hernan Javier Cutuli⁶⁰ · Mauricio Fernandez^{15,61} · Maria Bassanelli⁶² · Giandomenico Roviello⁶³ · Halima Abahssain⁶⁴ · Giuseppe Procopio^{50,65} · Michele Milella⁶⁶ · Jindrich Kopecky⁶⁷ · Angelo Martignetti⁶⁸ · Carlo Messina⁶⁹ · Manuel Caitano^{15,70} · Eva Inman^{15,71} · Ravindran Kanesvaran⁷² · Daniel Herchenhorn^{15,73} · Daniele Santini⁷⁴ · Ray Manneh⁷⁵ · Renato Bisonni⁷⁶ · Roubini Zakopoulou⁷⁷ · Alessandra Mosca⁷⁸ · Franco Morelli⁷⁹ · Fernando Maluf^{80,81} · Andrey Soares^{15,81,82} · Fernando Nunes^{15,83} · Alvaro Pinto⁸⁴ · Anca Zgura⁸⁵ · Lorena Incorvaia⁸⁶ · Jawaher Ansari⁸⁷ · Ignacio Ortego Zabalza²¹ · Johannes Landmesser⁸⁸ · Alessandro Rizzo⁸⁹ · Veronica Mollica⁶ · Giulia Sorgentoni⁹⁰ · Nicola Battelli⁹⁰ · Camillo Porta⁹¹ · Joaquim Bellmunt⁹² · Matteo Santoni⁹⁰

✉ Ondřej Fiala
fialao@fnplzen.cz

¹ Department of Oncology and Radiotherapeutics, Faculty of Medicine and University Hospital Pilsen, Charles University, Alej Svobody 80, 304 60 Pilsen, Czech Republic

² Biomedical Center, Faculty of Medicine in Pilsen, Charles University, Pilsen, Czech Republic

³ Department of Medicine and Surgery, University Hospital of Parma, Parma, Italy

⁴ Medical Oncology Unit, University Hospital of Parma, Parma, Italy

⁵ Department of Urology, Saitama Medical Center, Saitama Medical University, Saitama, Japan

⁶ Medical Oncology, IRCCS Azienda Ospedaliero-Universitaria Di Bologna, Via Albertoni - 15, Bologna, Italy

⁷ Department of Medical Oncology, Institutd' Investigació Biomèdica Sant Pau, Hospital de La Santa Creu I Sant Pau, Barcelona, Spain

⁸ Oncology 1 Unit, Department of Oncology, Istituto Oncologico Veneto IOV IRCCS, 35128 Padua, Italy

⁹ UOC Di Oncologia, Azienda Ospedaliera Di Rilievo Nazionale Cardarelli Di Napoli, Naples, Italy

¹⁰ Department of Urology, University Hospital Bonn (UKB), 53127 Bonn, Germany

¹¹ IRCCS Ospedale Policlinico San Martino, Genoa, Italy

¹² Markey Cancer Center, University of Kentucky, Lexington, KY 40536-0293, USA

¹³ Oncology Unit 2, University Hospital of Pisa, 56126 Pisa, Italy

¹⁴ Hospital São Rafael Oncologia D'Or, Salvador, BA, Brazil

¹⁵ Latin American Cooperative Oncology Group – LACOG, Porto Alegre, Brazil

¹⁶ Department of Urology, Medical University of Innsbruck, Anichstrasse 35, 6020 Innsbruck, Austria

¹⁷ Department of Medical Oncology, IRCCS Istituto Romagnolo Per Lo Studio Dei Tumori (IRST) “Dino Amadori”, Meldola, Italy

¹⁸ Hospital Angel Roffo, Buenos Aires, CABA, Argentina

¹⁹ University of Colorado Cancer Center, Anschutz Medical Campus, Aurora, CO, USA

²⁰ Oncology Institute of Vojvodina, Faculty of Medicine, University of Novi Sad, Novi Sad, Serbia

²¹ Department of Medical Oncology, MD Anderson Cancer Center Madrid, Madrid, Spain

²² Department of Radiological, Oncological and Anatomopathological Science, Sapienza University of Rome, Viale Regina Elena 324, 00185 Rome, Italy

²³ Department of Medical Oncology, Università Politecnica Delle Marche, AOU Ospedali Riuniti Delle Marche, Ancona, Italy

²⁴ Southampton Experimental Cancer Medicine Centre, University of Southampton, Southampton, UK

²⁵ Department of Medical Oncology, Hospital Ramón Y Cajal, Madrid, Spain

²⁶ Centro de Pesquisas Oncológicas - CEPON, Florianópolis, SC, Brazil

²⁷ Hospital Erasto Gaertner, Curitiba, PR, Brazil

²⁸ Oncologia Medica, Fondazione Policlinico Universitario Agostino Gemelli IRCCS, Rome, Italy

- 29 Department of Oncology, San Camillo Forlanini Hospital, Rome, Italy
- 30 Department of Medical Oncology, Bakirköy Dr. Sadi Konuk Training and Research Hospital, Zuhuratbaba District, Tevfik Saglam St. No: 11, Bakirkoy, Istanbul, Turkey
- 31 Unità Di Oncologia Medica, Azienda Ospedaliero Universitaria Di Cagliari, Cagliari, Italy
- 32 Department of Genitourinary Medical Oncology and Clinical Pharmacology, National Institute of Oncology, Budapest, Hungary
- 33 UOC Oncologia, Azienda Ospedaliera Ospedali Riuniti Marche Nord, Fano, Italy
- 34 Centro Oncologico Estatal “Dr José Luis Barrera Franco” del ISSEMYM, Toluca de Lerdo, Mexico
- 35 Medical Oncology Unit, Santa Chiara Hospital, Trento, Italy
- 36 Pontificia Universidade Católica Do Rio Grande Do Sul - PUCRS, Porto Alegre, RS, Brazil
- 37 Department of Uro-Oncology, Maria Sklodowska-Curie National Research Institute of Oncology, Warsaw, Poland
- 38 Department of Oncology, IRCCS Humanitas Research Hospital, Rozzano, Milan, Italy
- 39 Department of Biomedical Sciences, Humanitas University, Pieve Emanuele, Milan, Italy
- 40 Hospital AC Camargo, São Paulo, SP, Brazil
- 41 Niguarda Cancer Center, Grande Ospedale Metropolitano Niguarda, Milan, Italy
- 42 Department of Oncology, Hospital Sírio-Libanês, São Paulo, SP, Brazil
- 43 Medical Oncology Division of Urogenital and Head and Neck Tumours, IEO, European Institute of Oncology IRCCS, Milan, Italy
- 44 Clínica AMO, Salvador, BA, Brazil
- 45 Oncologia D’Or, Fortaleza, CE, Brazil
- 46 Division of Oncology, Institute for Cancer Research and Treatment, Asl Cn2 Alba-Brà, 12051 Alba-Brà, Italy
- 47 Instituto Alexander Fleming, Buenos Aires, CABA, Argentina
- 48 Division of Medical Oncology, A.O.U. Consorziata Policlinico Di Bari, Piazza G. Cesare 11, 70124 Bari, Italy
- 49 Urologic Oncology, Champalimaud Clinical Center, 1400-038 Lisbon, Portugal
- 50 Dipartimento Di Oncologia Medica, Fondazione IRCCS Istituto Nazionale Dei Tumori, Milan, Italy
- 51 Clinicas Medicas Especializadas NUCARE, Guatemala City, Guatemala
- 52 Oncology and Hematology Department, Hospital Santa Lucia, SHLS 716 Cj. C, Brasília, DF 70390-700, Brazil
- 53 Hospital Sao Rafael, Salvador, BA, Brazil
- 54 Department of Medical Oncology, “Maggiore Della Carità” University Hospital, 28100 Novara, Italy
- 55 Universidade Federal de Minas Gerais - UFMG, Belo Horizonte, MG, Brazil
- 56 Hospital Santa Casa de Sao Paulo, São Paulo, SP, Brazil
- 57 Department of Medical Oncology, Centre Hospitalier de Jolimont, Haine Saint Paul, Belgium
- 58 Clinica Medica Especializada en Oncologia Medica, Guatemala City, Guatemala
- 59 SOC Oncologia Medica, Azienda Ospedaliera “Pugliese – Ciaccio”, Catanzaro, Italy
- 60 Hospital Sirio Libanes, Buenos Aires, CABA, Argentina
- 61 Fundacion Centro Oncologico de Integracion Regional - COIR, Mendoza, Argentina
- 62 Medical Oncology, I-IRCCS Regina Elena National Cancer Institute, Rome, Italy
- 63 Department of Health Sciences, Section of Clinical Pharmacology and Oncology, University of Florence, Viale Pieraccini 6, 50139 Florence, Italy
- 64 Medical Oncology Unit, National Institute of Oncology, Medicine and Pharmacy Faculty, Mohammed V University, Rabat, Morocco
- 65 Oncologia Medica, Ospedale Maggiore Di Cremona, Cremona, Italy
- 66 Section of Oncology, Department of Medicine, School of Medicine and Verona University Hospital Trust, University of Verona, Verona, Italy
- 67 Department of Clinical Oncology and Radiotherapy, University Hospital Hradec Kralove, Hradec Kralove, Czech Republic
- 68 UOC Oncologia Medica, Ospedale Dell’alta Val D’Elsa – Usl sud est Toscana Area Senese, Poggibonsi, Italy
- 69 Oncology Unit, A.R.N.A.S. Civico, Palermo, Italy
- 70 Hospital do Câncer Porto Dias – Rede Mater Dei de Saúde, Belém, PA, Brazil
- 71 ONCOR Life Medical Center, Saltillo, Mexico
- 72 National Cancer Centre Singapore, Singapore, Republic of Singapore
- 73 Instituto D’Or de Ensino E Pesquisa, Rio de Janeiro, RJ, Brazil
- 74 Department of Radiological, Oncological and Pathological Sciences, Policlinico Umberto I, Sapienza University of Rome, Rome, Italy
- 75 Clinical Oncology, Sociedad de Oncología Y Hematología del Cesar, Valledupar, Colombia
- 76 UOC Oncologia Medica, Ospedale A. Murri, Fermo, Italy
- 77 2nd Propaedeutic Department of Internal Medicine, Attikon University Hospital, National and Kapodistrian University of Athens, Athens, Greece
- 78 Department of Oncology Candiolo Cancer Institute, FPO-IRCCS, Candiolo, Turin, Italy
- 79 Medical Oncology Unit, Gemelli Molise Hospital, Università Cattolica del Sacro Cuore, Campobasso, Italy
- 80 Hospital Beneficencia Portuguesa de São Paulo, São Paulo, SP, Brazil
- 81 Hospital Israelita Albert Einstein, São Paulo, SP, Brazil
- 82 Centro Paulista de Oncologia/Oncoclinicas, Sao Paulo, Brazil

- ⁸³ Clinica de Oncologia - Clion, Salvador, BA, Brazil
- ⁸⁴ Medical Oncology Department, La Paz University Hospital, Madrid, Spain
- ⁸⁵ Department of Oncology-Radiotherapy, Prof. Dr. Alexandru Trestioreanu Institute of Oncology, University of Medicine and Pharmacy, Bucharest, Romania
- ⁸⁶ Department of Surgical, Oncological and Oral Sciences, Section of Medical Oncology, University of Palermo, Palermo, Italy
- ⁸⁷ Medical Oncology, Tawam Hospital, Al Ain, United Arab Emirates
- ⁸⁸ Klinik Für Urologie, Ratzeburger Allee 160, 23538 Lübeck, Germany
- ⁸⁹ Struttura Semplice Dipartimentale Di Oncologia Medica Per La Presa in Carico Globale del Paziente Oncologico “Don Tonino Bello”, I.R.C.C.S. Istituto Tumori “Giovanni Paolo II”, Viale Orazio Flacco 65, 70124 Bari, Italy
- ⁹⁰ Oncology Unit, Macerata Hospital, Via Santa Lucia 2, 62100 Macerata, Italy
- ⁹¹ Interdisciplinary Department of Medicine, University of Bari “Aldo Moro”, Bari, Italy
- ⁹² Dana Farber Cancer Institute, Harvard Medical School, Boston, MA, USA