Segala *et al. BMC Public Health* (2024) 24:2271
https://doi.org/10.1186/s12889-024-19763-5

BMC Public Health

RESEARCH Open Access

Measuring preparedness to infectious diseases among people exposed to climate disasters in Cabo Delgado, Mozambique: a cross-sectional study

Francesco Vladimiro Segala[1*], Edoardo Occa[2], Elsa Chambisse[2], Francesco Cavallin[3], Agostinho Nanomba[2], Rumela Cobre[2], Giorgia Gelfi[2], Vitor Laguessa[2], Simone Cadorin[2], Abdul Chorai[2], Ketan Chitnis[4], Claudia Marotta[5], Inusso Chuau[4], Angelo Ghelardi[4], Aida Mahomed[4], Giovanni Putoto[5], Manuel Aly Mussa[6], Annalisa Saracino[1] and Francesco Di Gennaro[1]

Abstract

Introduction Climate change is contributing to increase the frequency and severity of climate disasters in Mozambique, leading, since 2019, to extensive damage to infrastructure and displacement 1.3 million people. Aim of this study is to evaluate baseline preparedness to vector-borne and water-borne infections among households and internally displaced people exposed to climate disasters in Mozambique.

Methods This was a cross-sectional, community-based survey assessing the preparedness to infectious diseases outbreaks among people exposed to climate disasters in six districts in Mozambique. Structured form was delivered via face-to-face between October 15th and November 7th, 2022. Study outcome was defined as a seven-point score of preparedness to infectious disease outbreaks. Multivariable analysis of the score was conducted using Conway-Maxwell-Poisson regression.

Results This study included 2,140 households and 11,239 people, with IDPs accounting for 30% of them. Overall, 1,186 (55.4%) households were overcrowded. Median score of preparedness was 3 points (IQR 2–4). At multivariable analyses, districts with low preparedness were Montepuez and Mueda. Higher preparedness was associated with family planning ($p < 0.0001$), access to primary education for all children living in the household ($p < 0.001$) and possession of a birth certificate for all children aged < 5y ($p < 0.0001$), while preparedness was heterogeneous among the districts ($p < 0.05$). Households composed by IDPs were not associated with a lower preparedness score.

Conclusions In climate-vulnerable communities in Mozambique, households practicing family planning, providing access to primary education and birth certificate for all children were less vulnerable to water-borne and vector-borne infectious disease outbreaks. Being family planning and childcare primarily performed by women, our findings can

*Correspondence:
Francesco Vladimiro Segala
fvsegala@gmail.com

Full list of author information is available at the end of the article

© The Author(s) 2024. **Open Access** This article is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License, which permits any non-commercial use, sharing, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if you modified the licensed material. You do not have permission under this licence to share adapted material derived from this article or parts of it. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by-nc-nd/4.0/.

Content courtesy of Springer Nature, terms of use apply. Rights reserved.

inform policymakers and practitioners on the importance of addressing women to mitigate the impact of climate disasters and reduce the risk of infectious disease outbreaks.

Keywords Climate change, WASH, Cholera, Malaria, Internally displaced people, Climate-sensitive infectious diseases, Low- and middle-income countries, Sub-saharan africa, Mozambique

Background

Human-induced climate change is exacerbating the frequency and severity of climate disasters in Mozambique [1]. Since 2019, extremely vulnerable communities in Mozambique were hit by Ciclone Idai, Cyclone Kenneth, Tropical Storm Ana, Cyclone Guambe and Cyclone Freddy [2]. These disasters have impacted over 3.1 million people, resulting in the destruction of approximately 318,000 houses and displacing up to 1.3 million people [3–6]. Internally displaced people (IDPs) are especially vulnerable to infectious diseases due to their profoundly under-resourced living conditions [7].

Frequent floods, droughts, and cyclones have caused extensive damage to infrastructure, including water and sanitation systems. These disruptions in hygiene and prevention practices have escalated the risk of waterborne [8] and vector-borne diseases [9]. Malaria and cholera – two highly climate-sensitive diseases tracked by the Lancet Countdown [10] – frequently lead to epidemics following cyclones and floods.

Only Cyclone Freddy, which made landfall in Mozambique twice in less than a month (24 February and 11 March 2023) [11], lead to nearly 30,000 cholera cases [6], while cyclone Idai led to nearly 26,000 cumulative cases of malaria [3]. Risk factors for cholera include poverty, overcrowding, inadequate access to clean water and poor hygiene conditions such as the impossibility to wash hands, dispose waste and access to proper sanitation facilities. At present, Cabo Delgado province is facing the deadliest cholera outbreak in the country in the past 25 years. Since September 2022, the outbreak has resulted in over 6,000 reported cases, with 801 (3.1%) of these cases originating from Cabo Delgado itself [12]. On the other hand, prevention against vector-borne diseases if heavily affected by humanitarian emergencies, which disrupt access to health facilities and public health programs such as mass insecticide-treated nets (ITN) distribution, or indoor residual spraying [13]. Cabo Delgado, this exacerbates the disease burden in the province with the highest malaria prevalence, which reaches 57.3% in children aged 0–5 years. The situation is further exacerbated by severe malnutrition and ongoing armed conflicts, resulting in the displacement of up to one million individuals. These conflicts are fueled by the socioeconomic exclusion experienced by northern Mozambique, alongside political and religious beliefs, and are intensified by the inequalities stemming from mineral and natural gas discoveries in the area [14].

In response to the recurrent outbreaks exacerbated by climate disasters, Doctors with Africa CUAMM (University College for Aspiring Missionary Doctors) and UNICEF (United Nations Children's Fund) Mozambique, conducted a multi-stakeholder, Social Behavioural Change program titled "Support adoption of key behaviors to prevent Covid-19, cholera, malaria, malnutrition and improve access to services in Cabo Delgado". Implemented in six districts of the Cabo Delgado province, the program focuses on piloting the national strategy for preventing communicable diseases, primarily cholera. Its objective is to encourage family units to adopt fourteen observable, low-cost behaviors that immediately reduce vulnerability to climate-sensitive outbreaks.

Aim of this study is to evaluate baseline preparedness to vector-borne and water-borne infections among family units and internally displaced people exposed to climate disasters in Mozambique.

Methods

Study design

This was a cross-sectional, community-based survey assessing the preparedness to vector-borne and water-borne infectious diseases outbreaks among people exposed to climate disasters in Mozambique. The survey was conducted between October 15th and November 7th, 2022, with face-to-face interviews. This reports follows the STROBE statement for cross-sectional studies [15].

Study setting

In Cabo Delgado province, resettlement started as early as March 2019, when both Cyclone Idai and non-state armed group (NSAG) violence hit local communities and continued through COVID-19 pandemic until mid-2022. As of today, roughly one million IDPs are living in the Cabo Delgado province. The present study was conducted in six of the seventeen districts in Cabo Delgado, which were supported by both CUAMM and UNICEF: Ancuabe, Balama, Chiure, Ibo, Montepuez, Mueda. The study involved both IDP sites and villages hosting sedentary population. At the time the study was conducted, all districts were affected by conflicts involving the Islamist armed group Ahlu Sunna Wal Jama [16].

Data collection and sampling

Data collection was conducted by trained community health workers who administered structured

Content courtesy of Springer Nature, terms of use apply. Rights reserved.

questionnaires through face-to-face interviews. Responses were standardized as either "available" or "not available". All variables included in the questionnaire were initially derived from the 2028 Mozambique Malaria indicator survey [17] and from the annual Global World Health Organization report on water, sanitation, and hygiene [18]. Subsequently, they were adjusted to suit the local context with the assistance of community leaders. A convenience, non-probability sampling method was employed, relying on the availability of family units and approval from local authorities. Prior to study implementation, residents in target resettlement sites and villages were extensively informed through dedicated meetings led by community leaders, who also supervised and assisted in the fieldwork.

Study outcome

To assess the household preparedness for waterborne and vector-borne infectious disease, seven variables were identified as indicators of adaptation to climate-sensitive outbreaks. These included (i) presence of insecticide treated mosquito nets (ITNs) [13]; (ii) presence of a functional latrine; (iii) presence of a sanitary landfill to dispose waste; (iv) capacity to dry food; (v) possibility to protect aliments from domestic animal contamination (i.e.: availability of a pylon stick); (vi) possibility of handwashing and (vii) disposal of safe water treatment and water storage system [19, 20]. Each of these practices was assigned a score of 1 point if present, resulting in a final score ranging from 0 (indicating complete unpreparedness) to 7 (indicating high preparedness).

Statistical analysis

Continuous data were summarized as median and interquartile range (IQR), and categorical data as number and percentage. Multivariable analysis of the score of preparedness to infection disease outbreaks was conducted using Conway-Maxwell-Poisson regression, which was preferred to Poisson regression due to the underdispersion of the dependent variable in the model [21]. Multivariable analysis was performed in different sets of family units, as follows. In the whole sample, the model included the following independent variables: district, family planning, overcrowding (at least 5 people in the family unit), and the presence of IDP, children aged 0–5 months, children aged 6–60 months, children aged 6–14 years, pregnant women, lactating women, adults with disability and children with disability. In the subset of family units with children of any age, a variable about nutrition (all children eating at least three times a day) was added to the original list of independent variables. In the subset of family units with children aged ≥ 6 years, a variable about nutrition (all children eating at least three times a day) and a variable about school (all children aged ≥ 6 years go to school) were added to the original list of independent variables, while the variables about child age (presence of children aged 0–5 months, children aged 6–60 months, and children aged 6–14 years) were removed. In the subset of family units with children aged 0–60 months, a variable about nutrition (all children eating at least three times a day) and having the birth certificate were added to the original list of independent variables, while one variable about child age (presence of children aged 6–14 years) was removed. All tests were 2-sided and a p-value < 0.05 was considered statistically significant. Statistical analysis was performed using R 4.3 (R Foundation for Statistical Computing, Vienna, Austria) [22].

Ethical approval

Data collection that informs this study served as a pilot activity for the program implementation mentioned earlier. The project operates under a framework agreement between CUAMM and the government of Mozambique, and an agreement with Cabo Delgado province, which authorizes all study activities. Participation was voluntary, anonymous, and without compensation. Data were collected anonymously at the family-unit level and aggregated for analysis.

Results

This study included a total of 2,140 households and 11,239 people. Characteristics of the study participants are summarized in Table1. Most households lived in the Mueda district (n=803, 37.5%), followed by Ancuabe (n=416, 19.5%) and Montepuez (n=371, 17.3%). Overall, IDPs accounted for 30% (n=3,375/11,239) of the participants and 37.5% (n=803/2,140) of all households. Displacements were non-uniformly geographically distributed, ranging from 74.1% (n=1100) of all people living in Chiure to 10.8% (n=1,215) of the people living in Balama district. Also, 979/1,857 (52.7%) households with children were able to provide at least 3 meals a day and 484/1,496 (32.4%) households with school-aged children were able to provide primary education.

Information regarding adaptation to climate-sensitive outbreaks is reported in Fig. 1. The median score of preparedness to infection disease outbreaks was 3 out of maximum 7 points (IQR 2–4). Overll, the most available adaptation strategy to climate-sensitive outbreaks was insecticide-treated mosquito net (n=1,697, 79.3%), while the least available ones were the pylon stick (n=203, 9.5%) – a tool used to protect food from domestic animal contamination – and the availability of handwashing facilities (n=282, 13.2%).

Table2 presents the results from multivariable analyses of the preparedness score for infectious disease outbreaks. In synthesis, family planning was associated with higher score of preparedness (p<0.0001) across

Content courtesy of Springer Nature, terms of use apply. Rights reserved.

Table 1 Characteristics of the study participants, overall and stratified per study district

N of family units	**Overall 2140**	**Ancuabe 416 (19.5)**	**Balama 251 (11.7)**	**Chiure 319 (14.9)**	**Ibo 306 (14.3)**	**Montepuez 371 (17.3)**	**Mueda 477 (22.3)**
Households with IDP [b]	803 (37.5)	129 (31.0)	16 (6.4)	246 (77.1)	183 (59.8)	94 (25.3)	135 (28.3)
Households with at least one child aged 0–5 months	395 (18.5)	62 (14.9)	82 (32.7)	57 (18.2)	60 (19.6)	61 (16.4)	72 (15.1)
Households with at least one child aged 6–60 months	1201 (56.1)	255 (61.3)	151 (60.2)	180 (56.4)	201 (65.7)	211 (56.9)	203 (42.6)
Households with at least one child aged 6–14 years	1496 (69.9)	286 (68.8)	167 (66.5)	219 (68.7)	222 (72.5)	267 (72.0)	335 (70.2)
Households with at least one pregnant woman	275 (12.9)	53 (12.7)	52 (20.7)	46 (14.4)	42 (13.7)	33 (8.9)	49 (10.3)
Households with at least one lactating woman	684 (32.0)	144 (34.6)	106 (42.2)	104 (32.6)	96 (31.4)	116 (31.3)	118 (24.7)
Households with at least one adult with disability	110 (5.1)	21 (5.0)	0 (0)	15 (4.7)	24 (7.8)	9 (2.4)	41 (8.6)
Households with at least one child with disability	65 (3.0)	14 (3.4)	1 (0.4)	12 (3.8)	18 (5.9)	3 (0.8)	17 (3.6)
Households with overcrowding (at least 5 people)	1186 (55.4)	251 (60.3)	129 (51.4)	153 (48.0)	183 (59.8)	225 (60.6)	245 (51.4)
Households doing family planning (contraception)	712/2139 (33.3)	127 (30.5)	88 (35.1)	108 (33.9)	139/307 (45.6)	111 (29.9)	139 (29.1)
Households with all children eating at least 3 times a day	979/1859 (52.7)	175/362 (43.3)	181/232 (78.0)	93/283 (32.9)	137/272 (50.4)	207/316 (65.5)	186/394 (47.2)
Households in which all children aged ≥ 6 years go to school	1115/1496 (74.5)	221/286 (77.3)	131/167 (78.4)	180/219 (82.2)	197/222 (88.7)	163/267 (61.0)	223/335 (66.6)
Households in which all children aged ≥ 6 years concluded primary school	484/1496 (32.4)	74/286 (25.9)	78/167 (46.7)	34/219 (15.5)	63/222 (28.4)	101/267 (37.8)	134/335 (40.0)
Households where all children aged 0–5 years have a birth certificate	867/1367 (63.4)	195/282 (69.1)	135/190 (71.1)	92/208 (44.2)	190/215 (88.4)	132/230 (57.4)	123/242 (50.8)
Total n of people	**Overall 11,239**	**Ancuabe 2285 (20.3)**	**Balama 1215 (10.8)**	**Chiure 1485 (13.2)**	**Ibo 1775 (15.8)**	**Montepuez 2071 (18.4)**	**Mueda 2408 (21.4)**
N of IDP	3375 (30.0)	553 (24.2)	56 (4.6)	1103 (74.3)	718 (40.5)	508 (24.5)	437 (18.1)
N of children aged 0–5 months	430 (3.8)	66 (2.9)	85 (7)	63 (4.2)	63 (3.5)	74 (3.6)	79 (3.3)
N of children aged 6–60 months	1660 (5.9)	379 (16.6)	210 (17.3)	231 (15.6)	296 (16.7)	271 (13.1)	273 (11.3)
N of children aged 6–14 years	2863 (24.4)	594 (26)	298 (24.5)	401 (27)	421 (23.7)	520 (25.1)	629 (26.1)
N of people aged > 14 years	6286 (55.9)	1246 (54.5)	622 (51.2)	790 (53.2)	995 (56.1)	1206 (58.2)	1427 (59.3)
N of pregnant women	291 (2.6)	59 (2.6)	54 (4.4)	50 (3.4)	44 (2.5)	34 (1.6)	50 (2.1)
N of lactating mothers	739 (6.6)	160 (7)	108 (8.9)	109 (7.3)	103 (5.8)	130 (6.3)	129 (5.4)
N of children aged 0–14 years with disabilities	73 (0.6)	21 (0.9)	1 (0.1)	13 (0.9)	18 (1)	3 (0.1)	17 (0.7)
N of people aged > 14 years with disabilities	128 (1.1)	29 (1.3)	0 (0)	17 (1.1)	24 (1.4)	10 (0.5)	48 (2)

Data summarized as n (%) or [a] median (IQR). [b] The median proportion of IDP within the household was 100% (IQR 57–100%). IDP: internally displaced people

the entire sample and in each subset. Additionally, the analyses indicated varying levels of adaptation to water-borne and vector-borne outbreaks among the six study districts ($p<0.001$), with Mueda and Montepuez showing the lowest preparedness. Among households with at least one child aged ≥ 6 years, the variable related to school attendance ("all children aged ≥ 6 years go to school") ($p<0.001$) and the presence of adults with disabilities were associated with a higher preparedness ($p<0.05$). Moreover, in family units with children aged 0–60 months, possessing birth certificates for all children was associated with higher preparedness for infectious disease outbreaks ($p<0.0001$). Notably, the variable regarding child nutrition ("all children eating at least three times a day") was not associated with the preparedness score in households with children.

Discussion

In this study, we observed varying levels of vulnerability against infectious diseases among people exposed to climate disasters in Mozambique, with factors such as district of residence, access to family planning, and key child development indicators influencing preparedness for these outbreaks. Following cyclones, tropical storms and floods, the risk for water-borne [20] and vector-borne [23] outbreaks increases substantially, with both cholera and malaria monitored by The Lancet Countdown for climate change's impact on human health [10]. Since, under current trends of fossil fuel extraction, global temperatures and extreme weather events are expected to rise [24], assessing adaptation strategies for climate-sensitive diseases imperative for public health in low- and middle-income countries (LMICs). Mozambique, despite producing roughly 0.02% of the annual share of global CO2

Content courtesy of Springer Nature, terms of use apply. Rights reserved.




Fig. 1 Adaptation to climate-sensitive outbreaks (**A**), score of preparedness to climate-sensitive outbreaks in the overall population (**B**) and score result distribution stratified per study district (**C**)

emissions [25], is profoundly affected by human-induced climate change [26].

In this study, we assessed vulnerability to climate-sensitive outbreaks using a validated score comprising seven low-cost prevention strategies applicable during humanitarian emergencies, including insecticide-treated nets [13], and practices related to water, sanitation, and hygiene (WASH) [27]. A peculiar finding of our study was that, even afted adjusting for other variables, household with internally displaced people did not did not exhibit significantly different preparedness levels for water- and vector-borne infectious diseases. This could be attributed to the equitable access to sanitation resources among both internally displaced people and residents, along with comparable participation in public health initiatives such as insecticide-treated net distributions. Additionally, our survey targeted areas where people were displaced since early 2019, when Cyclone Idai made landfall, potentially equalizing access to sanitation services over time. Furthermore, despite the lack of association between outbreak preparedness and the provision of three meals per day for all children, this finding may be explained by the region's status as the most food insecure province in Mozambique during the study period, as reported by The United Nations World Food Programme. [28].

Interestingly, our multivariable analysis showed that indicators positively associated with high preparedness against climate-sensitive infectious diseases were were predominantly tasks associated with women: family planning and childcare [29]. Qualitative studies conducted in Mozambique [30, 31] and other sub-Saharan countries have consistently indicated low male participation in family planning, and in some cases, obstruction' [32]. This assumption is supported by by the lack of association between other indicators, such as overcrowding and presence of IDP in the household, and preparedness score. While acknowledging that the correlation found in our analysis does not imply causation – we highlight that family planning serves as an indicator of socio-economic wellbeing and positive female personal agency. It is associated with low child mortality [33], fewer unsafe abortions [34] and overall lower maternal mortality rates [35]. Thus, family planning may serve as a proxy for household resilience against climate-sensitive infectious diseases.

Content courtesy of Springer Nature, terms of use apply. Rights reserved.

Table 2 Multivariable analysis of the score of preparedness to infection disease outbreaks (as proxy of household adaptation to climate-sensitive outbreaks)

	All family units		Family units with children of any age		Family units with children aged ≥ 6 years		Family units with children aged 0–60 months	
	Regression coefficient (95% confidence interval)	*p*-value	Regression coefficient (95% confidence interval)	*p*-value	Regression coefficient (95% confidence interval)	*p*-value	Regression coefficient (95% confidence interval)	*p*-value
Districts:								
Ancuabe	Reference	-	Reference	-	Reference	-	Reference	-
Balama	-0.10 (-0.22 to 0.01)	0.08	-0.07 (-0.20 to 0.05)	0.23	-0.22 (-0.45 to 0.01)	0.06	-0.07 (-0.21 to 0.07)	0.34
Chiure	0.25 (0.14 to 0 0.36)	**< 0.0001**	0.31 (0.19 to 0.43)	**< 0.0001**	0.18 (-0.01 to 0.36)	0.06	0.33 (0.18 to 0.47)	**< 0.0001**
Ibo	0.41 (0.30 to 0.52)	**< 0.0001**	0.44 (0.32 to 0.56)	**< 0.0001**	0.35 (0.16 to 0.53)	**0.0002**	0.40 (0.26 to 0.53)	**< 0.0001**
Montepuez	-0.21 (-0.32 to -0.10)	**0.0001**	-0.18 (-0.30 to -0.07)	**0.001**	-0.31 (-0.49 to -0.13)	**0.0005**	-0.13 (-0.26 to 0.01)	0.06
Mueda	-0.33 (-0.44 to -0.23)	**< 0.0001**	-0.32 (-0.44 to -0.21)	**< 0.0001**	-0.35 (-0.52 to -0.19)	**< 0.0001**	-0.31 (-0.45 to -0.16)	**< 0.0001**
Household with IDP	0.01 (-0.05 to 0.09)	0.61	0.00 (-0.07 to 0.07)	0.97	-0.02 (-0.14 to 0.09)	0.72	0.06 (-0.02 to 0.15)	0.16
Households with at least one child aged 0–5 months	0.06 (-0.04 to 0.17)	0.22	0.06 (-0.04 to 0.17)	0.25	-	-	0.08 (-0.04 to 0.20)	0.18
Households with at least one child aged 6–60 months	-0.06 (-0.14 to 0.01)	0.06	-0.06 (-0.14 to 0.01)	0.10	-	-	-0.04 (-0.19 to 0.10)	0.57
Households with at least one child aged 6–14 years	0.04 (-0.03 to 0.12)	0.26	0.05 (-0.04 to 0.15)	0.27	-	-	-	-
Household with at least one pregnant woman	0.02 (-0.06 to 0.12)	0.53	0.02 (-0.07 to 0.12)	0.66	-0.02 (-0.19 to 0.14)	0.79	0.03 (-0.07 to 0.15)	0.52
Households with at least one lactating woman	-0.01 (-0.10 to 0.08)	0.84	0.00 (-0.09 to 0.08)	0.91	0.02 (-0.33 to 0.38)	0.88	0.00 (-0.09 to 0.09)	0.95
Households with at least one adult with disability	0.04 (-0.10 to 0.18)	0.56	0.00 (-0.15 to 0.16)	0.94	0.22 (0.01 to 0.45)	**0.04**	-0.09 (-0.28 to 0.10)	0.34
Households with at least one child with disability	-0.05 (-0.23 to 0.12)	0.55	-0.02 (-0.20 to 0.16)	0.83	0.00 (-0.39 to 0.37)	0.97	-0.05 (-0.26 to 0.15)	0.62
Households with overcrowding (at least 5 people)	-0.02 (-0.10 to 0.06)	0.61	-0.01 (-0.09 to 0.06)	0.72	-0.10 (-0.22 to 0.01)	0.09	0.00 (-0.09 to 0.09)	0.95
Households doing family planning (contraception)	0.22 (0.16 to 0.29)	**< 0.0001**	0.20 (0.13 to 0.28)	**< 0.0001**	0.22 (0.10 to 0.35)	**0.0003**	0.21 (0.12 to 0.29)	**< 0.0001**
Households with all children eating at least 3 times a day	-	-	0.02 (-0.04 to 0.09)	0.51	-0.04 (-0.16 to 0.07)	0.47	0.03 (-0.04 to 0.12)	0.40
Households where all children aged ≥ 6 years go to school	-	-	-	-	0.19 (0.08 to 0.62)	**0.0007**	-	-
Households where all children aged 0–5 years have a birth certificate	-	-	-	-	-	-	0.22 (0.12 to 0.31)	**< 0.0001**

Each model did not include variables that could pose restrictions to the family units or could be incompatible with the subset of family units

p values < 0.05 are reported in bold

Content courtesy of Springer Nature, terms of use apply. Rights reserved.

Recognizing the pivotal role of women during humanitarian crises in LMICs is essential in the fight against climate change and climate-sensitive infectious diseases. In sub-Saharan Africa, prioritizing women empowerment and agency through comprehensive strategies is essential for building resilient communities [36]. Beyond improving WASH behaviors, empowering women can lead to a better access to healthcare, increased education and awareness, enhanced economic growth, and contribute to long-term community resilience [37]. Despite facing disproportionate burden from the impacts of climate change – in terms of reduction of agency, increased unpaid workload and exacerbated gender-based violence [38] – women are pivotal in both adaptation and mitigation to the climate crisis [39]. The results of our study suggest that, by empowering women to lead community-based interventions, vulnerable people in Mozambique can enhance their capacity to combat cholera, malaria, and other infectious diseases that will be fostered by the ongoing climate crisis.

This study has some limitations that need to be considered. Firstly, our study was limited to six district in the Cabo Delgado province followed by CUAMM and UNICEF, thus limiting the generalizability of our findings to the broader Mozambican population. While our sample included data from more than two-thousands households, variations may exist between other provinces heavily impacted by cyclones and floods, such as Zambezia, Sofala, Nampula and Tete), which were not included in our study. Secondly, the unpredictable dynamics of extreme weather events, as well as the unique characteristics of a population both affected by climate disasters and conflicts, may further restrict the generalizability of our results. Thirdly, due to the challenging-to-reach nature of the population under study, we used convenience sampling, potentially affecting the representativeness of our findings. Fourthly, the outcome selected for this study served as a proxy of the overall prevention measures against post-disaster epidemics available to households at the time of the survey was conducted. Finally, we did not collect any clinical or microbiological data, precluding inferences regarding the true impact of water-borne and vector-borne outbreaks among the included population.

Among climate vulnerable people and IPD living in Mozambique, microbiological and syndromic surveillance of vector-borne and water-borne diseases is urgently needed. Further evidence is required to assess effective preventive strategies in reducing the burden of water-borne and vector-borne outbreaks in post-disaster humanitarian emergencies.

Conclusions

In climate-vulnerable communities in Mozambique, households practicing family planning and providing access to primary education and birth certificate to all children were foud to be less vulnerable to water-borne and vector-borne outbreaks. Our findings can inform policymakers and practitioners on the importance of addressing women to mitigate the impact of climate disasters and reduce the risk of infectious disease outbreaks. Also, our study clearly identifies two districts in which people are more vulnerable to climate-sensitive epidemics, which should be prioritized by local policymakers: Montepuez and Mueda.

Abbreviations

IDP	Internally displaces people
ITN	Insecticide-treated nets
LMIC	Low- and middle-income countries
NSAG	Non-state armed group

Acknowledgements

The authors thank all the population involved for the collaboration, the village leaders who contributed to creating a context of trust and interest, NIOP for the technical and institutional support, UNICEF Mozambique colleagues for their constant work. This research was partially supported by EU funding within the NextGenerationEU-MUR PNRR Extended Partnership initiative on Emerging Infectious Diseases (Project no. PE00000007, INF-ACT).

Author contributions

FVS, EO, FDG and CM contributed to study concept and design. EC, AN, RC, GG, VL, SC, AC, IC, AG, AM, MAM contributed to patient inclusion and data collection. FC and FVS cleaned the data and FC contributed to the statistical analysis. FVS, EO, FDG, CM and FC interpreted the data. FVS, FC, GP and EO drafted the manuscript. FDG, AS and KC critically revised the manuscript. All authors approved the final manuscript.
The study was carried out with the financial support of UNICEF Mozambique..

Data availability

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Declarations

Ethics approval and consent to participate

Data collection that informs this study was performed within a framework agreement between CUAMM and the government of Mozambique. The study was reviewed and approved by ethics committee of MINEC Ministerio dos Negocios Estrangeiros e Cooperaçao, Nota 1883/MINEC.DAJC.DONG/2022. Participation was voluntary, anonymous, and without compensation, and informed consent to participate in the study was obtained from all included households. This study has been conducted in accordance with the Declaration of Helsinki.

Consent for publication

Not applicable.

Competing interests

The authors declare no competing interests.

Author details

[1]Clinic of Infectious Diseases, Department of Precision and Regenerative Medicine and Ionian Area - (DiMePRe-J), University of Bari Aldo Moro, Bari, Italy
[2]Doctors with Africa CUAMM, Maputo, Mozambique
[3]Independent Statistician, Solagna, Italy
[4]UNICEF Mozambique, Maputo, Mozambique
[5]Operational Research Unit, Doctors with Africa CUAMM, Padua, Italy

Content courtesy of Springer Nature, terms of use apply. Rights reserved.

[6]NIOP Nucleo de Investigaçao Operacional del Pemba, Pemba, Mozambique

Received: 30 June 2023 / Accepted: 12 August 2024
Published online: 21 August 2024

References

1. Otto FEL, Zachariah M, Wolski P, Pinto I, Nhamtumbo B, Bonnet R et al. Climate change increased rainfall associated with tropical cyclones hitting highly vulnerable communities in Madagascar, Mozambique & Malawi.
2. Worlddata.info [Internet]. [cited 2023 May 7]. Most recent cyclones in Mozambique. https://www.worlddata.info/africa/mozambique/cyclones.php
3. Tropical Cyclones Idai and Kenneth, Mozambique National Situation Report 1. 10 May - Mozambique | ReliefWeb [Internet]. 2019 [cited 2023 May 7]. https://reliefweb.int/report/mozambique/tropical-cyclones-idai-and-kenneth-mozambique-national-situation-report-1-10-may
4. Mozambique. Tropical Storm Ana Flash Update No.6 (As of 30 January 2022) - Mozambique | ReliefWeb [Internet]. 2022 [cited 2023 May 7]. https://reliefweb.int/report/mozambique/mozambique-tropical-storm-ana-flash-update-no6-30-january-2022
5. WFP Mozambique Situation Report #1 Cyclone Gombe. (15 March 2022) - Mozambique | ReliefWeb [Internet]. 2022 [cited 2023 May 7]. https://reliefweb.int/report/mozambique/wfp-mozambique-situation-report-1-cyclone-gombe-15-march-2022
6. Mozambique TC. Freddy, Floods and Cholera - Situation Report No.3 - Mozambique | ReliefWeb [Internet]. 2023 [cited 2023 May 7]. https://reliefweb.int/report/mozambique/mozambique-tropical-cyclone-freddy-floods-and-cholera-situation-report-no3
7. Desai AN, Ramatowski JW, Marano N, Madoff LC, Lassmann B. Infectious disease outbreaks among forcibly displaced persons: an analysis of ProMED reports 1996–2016. Confl Health. 2020;14(1):49.
8. Di Gennaro F, Occa E, Chitnis K, Guelfi G, Canini A, Chuau I, et al. Knowledge, attitudes and practices on Cholera and Water, Sanitation, and Hygiene among internally displaced persons in Cabo Delgado Province, Mozambique. Am J Trop Med Hyg. 2023;108(1):195–9.
9. Gennaro FD, Occa E, Ramirez L, Marotta C, Segala FV, Santana J, et al. Malaria, HIV and Malnutrition among internally displaced people in Mozambique during COVID-19 pandemic: results from a community-based intervention. Annals Global Health. 2022;88(1):106.
10. Romanello M, Napoli CD, Drummond P, Green C, Kennard H, Lampard P, et al. The 2022 report of the Lancet countdown on health and climate change: health at the mercy of fossil fuels. Lancet. 2022;400(10363):1619–54.
11. Tropical Cyclone Freddy. may set new record | World Meteorological Organization [Internet]. [cited 2023 May 7]. https://public.wmo.int/en/media/news/tropical-cyclone-freddy-may-set-new-record
12. Occa E. Outlining a 'Semantic Anthropology of Data Science': the Humanitarian response to the Cholera epidemic of doctors with Africa CUAMM in Mozambique. Anthropol Action. 2023;30(3):40–8.
13. Messenger LA, Furnival-Adams J, Chan K, Pelloquin B, Paris L, Rowland M. Vector control for malaria prevention during humanitarian emergencies: a systematic review and meta-analysis. Lancet Global Health. 2023;11(4):e534–45.
14. Mozambique, Acute Malnutrition Situation for May - September. 2023 and Projection for October 2023 - March 2024 | IPC - Integrated Food Security Phase Classification [Internet]. [cited 2024 Feb 17]. https://www.ipcinfo.org/ipc-country-analysis/details-map/en/c/1156718/
15. von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP. The strengthening the reporting of Observational studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. Lancet. 2007;370(9596):1453–7.
16. ACAPS. ACAPS Thematic report - Mozambique: Impact of the five-year conflict in Cabo Delgado (07 July 2023) - Mozambique | ReliefWeb [Internet]. 2023 [cited 2024 Feb 17]. https://reliefweb.int/report/mozambique/acaps-thematic-report-mozambique-impact-five-year-conflict-cabo-delgado-07-july-2023
17. de Saúde/INS. Estatistica/INE IN de, Malária/PNCM PN de C da, ICF. Mozambique Inquérito Nacional sobre Indicadores de Malária (IIM)2018. 2019 Apr 1 [cited 2024 Feb 17]; https://dhsprogram.com/publications/publication-mis33-mis-final-reports.cfm
18. World Health Organization. WHO Global water, sanitation and hygiene: Annual report 2022 [Internet]. 2023 [cited 2024 Feb 17]. https://www.who.int/publications-detail-redirect/9789240076297
19. World Health Organization. Water, sanitation and hygiene (WASH) [Internet]. [cited 2023 May 23]. https://www.who.int/health-topics/water-sanitation-and-hygiene-wash
20. Lequechane JD, Mahumane A, Chale F, Nhabomba C, Salomão C, Lameira C, et al. Mozambique's response to cyclone Idai: how collaboration and surveillance with water, sanitation and hygiene (WASH) interventions were used to control a cholera epidemic. Infect Dis Poverty. 2020;9(1):68.
21. Shmueli G, Minka TP, Kadane JB, Borle S, Boatwright P. A useful distribution for Fitting Discrete Data: Revival of the Conway-Maxwell-Poisson distribution. J Royal Stat Soc Ser C (Applied Statistics). 2005;54(1):127–42.
22. R Core Team. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. [Internet]. 2023 [cited 2023 May 20]. https://www.r-project.org/
23. Nearly. 15,000 cases of malaria reported in areas of Mozambique affected by Cyclone Idai – UNICEF [Internet]. [cited 2023 May 23]. https://www.unicef.org/press-releases/nearly-15000-cases-malaria-reported-areas-mozambique-affected-cyclone-idai-unicef
24. AR6 Synthesis Report. Climate Change 2023 [Internet]. [cited 2023 May 23]. https://www.ipcc.ch/report/ar6/syr/
25. Ritchie H, Roser M, Rosado P, CO_2. and Greenhouse Gas Emissions. Our World in Data [Internet]. 2020 May 11 [cited 2023 May 23]; https://ourworldindata.org/co2-emissions
26. Increasing the resilience of Mozambique'. s health system to adverse climate change impacts [Internet]. [cited 2023 May 23]. https://www.who.int/about/accountability/results/who-results-report-2020-mtr/country-story/2021/mozambique
27. Humanitarian emergencies [Internet]. [cited 2023 May 23]. https://www.who.int/teams/environment-climate-change-and-health/water-sanitation-and-health/environmental-health-in-emergencies/humanitarian-emergencies
28. The United Nations World Food Programme. Funds dry up as hunger looms in Mozambique's north | World Food Programme [Internet]. 2022 [cited 2023 May 24]. https://www.wfp.org/news/funds-dry-hunger-looms-mozambiques-north
29. Osinuga A, Janssen B, Fethke NB, Story WT, Imaledo JA, Baker KK. Understanding Rural women's domestic work experiences (DWE) in Ibadan, Nigeria: development of a Measurement Tool using confirmatory factor analysis. Int J Environ Res Public Health. 2021;18(21):11043.
30. Pires P, Mupueleque M, Macaringue C, Zakus D, Siemens R, Belo C. Users' perspectives on the quality of family planning services in Mozambique: a case study. Pan Afr Med J. 2022;42:174.
31. S CL DC, Nb SF. O, J H. There Are Things We Can Do and There Are Things We Cannot Do. A Qualitative Study About Women's Perceptions on Empowerment in Relation to Fertility Intentions and Family Planning Practices in Mozambique. Frontiers in global women's health [Internet]. 2022 Mar 23 [cited 2023 May 23];3. https://pubmed.ncbi.nlm.nih.gov/35400131/
32. Gahungu J, Vahdaninia M, Regmi PR. The unmet needs for modern family planning methods among postpartum women in Sub-saharan Africa: a systematic review of the literature. Reproductive Health. 2021;18(1):35.
33. PRB [Internet]. [cited 2023 May 30]. Family Planning Saves Lives. https://www.prb.org/resources/family-planning-saves-lives/
34. UNICEF. [Benefits of family planning]. Bol Asoc Chil Prot Fam. 1992;28(7–12):3–5.
35. Save The Children S. Save the Children's Resource Centre. [cited 2023 May 30]. The Key to Achieving the Millennium Development Goals: Universal Access to Family Planning and Reproductive Health. https://resourcecentre.savethechildren.net/document/key-achieving-millennium-development-goals-universal-access-family-planning-and-reproductive/
36. Oyebanji O, Okereke E. Empowering African women leaders is key for health equity. Nat Hum Behav. 2023;1–3.
37. WHO | Regional Office for Africa [Internet]. 2023 [cited 2023 May 23]. Report: Addressing the Challenge of Women's Health in Africa. https://www.afro.who.int/publications/report-addressing-challenge-womens-health-africa
38. Dimensions. and examples of the gender-differentiated impacts of climate change, the role of women as agents of change and opportunities for women. Synthesis report by the secretariat | UNFCCC [Internet]. [cited 2023 May 24]. https://unfccc.int/documents/494455

Content courtesy of Springer Nature, terms of use apply. Rights reserved.

39. The UN Internet Gateway on Gender Equality and Empowerment of Women. Women, Gender Equality and Climate Chang [Internet]. United Nations; https://www.un.org/womenwatch/feature/climate_change/

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Content courtesy of Springer Nature, terms of use apply. Rights reserved.

Terms and Conditions

Springer Nature journal content, brought to you courtesy of Springer Nature Customer Service Center GmbH (“Springer Nature”).
Springer Nature supports a reasonable amount of sharing of research papers by authors, subscribers and authorised users (“Users”), for small-scale personal, non-commercial use provided that all copyright, trade and service marks and other proprietary notices are maintained. By accessing, sharing, receiving or otherwise using the Springer Nature journal content you agree to these terms of use (“Terms”). For these purposes, Springer Nature considers academic use (by researchers and students) to be non-commercial.
These Terms are supplementary and will apply in addition to any applicable website terms and conditions, a relevant site licence or a personal subscription. These Terms will prevail over any conflict or ambiguity with regards to the relevant terms, a site licence or a personal subscription (to the extent of the conflict or ambiguity only). For Creative Commons-licensed articles, the terms of the Creative Commons license used will apply.
We collect and use personal data to provide access to the Springer Nature journal content. We may also use these personal data internally within ResearchGate and Springer Nature and as agreed share it, in an anonymised way, for purposes of tracking, analysis and reporting. We will not otherwise disclose your personal data outside the ResearchGate or the Springer Nature group of companies unless we have your permission as detailed in the Privacy Policy.
While Users may use the Springer Nature journal content for small scale, personal non-commercial use, it is important to note that Users may not:

1. use such content for the purpose of providing other users with access on a regular or large scale basis or as a means to circumvent access control;
2. use such content where to do so would be considered a criminal or statutory offence in any jurisdiction, or gives rise to civil liability, or is otherwise unlawful;
3. falsely or misleadingly imply or suggest endorsement, approval , sponsorship, or association unless explicitly agreed to by Springer Nature in writing;
4. use bots or other automated methods to access the content or redirect messages
5. override any security feature or exclusionary protocol; or
6. share the content in order to create substitute for Springer Nature products or services or a systematic database of Springer Nature journal content.

In line with the restriction against commercial use, Springer Nature does not permit the creation of a product or service that creates revenue, royalties, rent or income from our content or its inclusion as part of a paid for service or for other commercial gain. Springer Nature journal content cannot be used for inter-library loans and librarians may not upload Springer Nature journal content on a large scale into their, or any other, institutional repository.
These terms of use are reviewed regularly and may be amended at any time. Springer Nature is not obligated to publish any information or content on this website and may remove it or features or functionality at our sole discretion, at any time with or without notice. Springer Nature may revoke this licence to you at any time and remove access to any copies of the Springer Nature journal content which have been saved.
To the fullest extent permitted by law, Springer Nature makes no warranties, representations or guarantees to Users, either express or implied with respect to the Springer nature journal content and all parties disclaim and waive any implied warranties or warranties imposed by law, including merchantability or fitness for any particular purpose.
Please note that these rights do not automatically extend to content, data or other material published by Springer Nature that may be licensed from third parties.
If you would like to use or distribute our Springer Nature journal content to a wider audience or on a regular basis or in any other manner not expressly permitted by these Terms, please contact Springer Nature at

onlineservice@springernature.com