

SYSTEMATIC REVIEW

Open Access



Evaluating the success rates and effectiveness of surgical and orthodontic interventions for impacted canines: a systematic review of surgical and orthodontic interventions and a case series

Antonio Mancini^{1†}, Fabrizio Chirico^{2†}, Giuseppe Colella², Fabio Piras¹, Valeria Colonna¹, Pierluigi Marotti¹, Claudio Carone¹, Alessio Danilo Inchingolo¹, Angelo Michele Inchingolo¹, Francesco Inchingolo^{1*} and Gianna Dipalma¹

Abstract

Objectives The goal is to gather more information on orthodontic traction of impacted canines in the arch or their extraction.

Methods We focused our search on English-language papers published between January 1, 2004, and July 24, 2024 in PubMed, Scopus, and Web of Science that were relevant to our topic. The search strategy employed the Boolean terms "impacted" AND "canine". In addition to the overview, this article discusses a number of clinical instances involving canine disinclusion.

Results The study analyzed 2748 papers from databases like Web of Science, PubMed, and Scopus, resulting in 1455 articles after eliminating duplicates. After examining titles and abstracts, 897 entries were removed. The remaining 558 papers were confirmed, with 543 items removed due to being off-topic. The study includes a qualitative analysis of the 15 final articles.

Conclusions This article concludes that both extraction and orthodontic traction are effective treatments for impacted canines, with the choice depending on patient-specific factors. Advances in surgical techniques and imaging technologies have improved outcomes, while a multidisciplinary, patient-centered approach remains essential. Future research should focus on refining treatment selection criteria and integrating genetic and technological advancements.

Keywords Canine, Impacted, Disinclusion, Extraction, Surgery, Orthodontic treatment, Mini-screw

[†]Antonio Mancini and Fabrizio Chirico contributed equally as first authors.

*Correspondence:

Francesco Inchingolo
francesco.inchingolo@uniba.it

Full list of author information is available at the end of the article



© The Author(s) 2025. **Open Access** This article is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License, which permits any non-commercial use, sharing, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if you modified the licensed material. You do not have permission under this licence to share adapted material derived from this article or parts of it. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by-nc-nd/4.0/>.

Introduction

Impacted canines as a dental anomaly: etiology and factors

Impacted canines represent a significant dental anomaly with far-reaching implications for both orthodontic and surgical interventions [1]. These teeth, crucial for functional occlusion and aesthetic harmony, often fail to erupt into their correct positions, leading to a range of clinical challenges [4]. Maxillary canines, in particular, exhibit a higher propensity for impaction, with prevalence rates reported to be between 1–3% in various populations [9]. This prevalence underscores the necessity for dental professionals to possess a thorough understanding of the aetiology, diagnosis, and management of impacted canines to optimize patient outcomes [14]. Etiologically, impacted canines can arise from a combination of genetic and environmental factors [19] [24]. Genetic predispositions may include familial patterns of dental anomalies, while environmental influences encompass discrepancies in dental arch space, abnormal tooth eruption paths, and the presence of physical barriers such as supernumerary teeth, odontomas, or cysts [26]. The multifactorial nature of canine impaction necessitates a comprehensive diagnostic approach [31].

Advances in diagnostic techniques

Accurate diagnosis typically involves a combination of clinical examination and radiographic imaging, with advanced techniques such as cone-beam computed tomography (CBCT) offering superior diagnostic accuracy by providing three-dimensional views of the impacted tooth and its surrounding structures [36]. Recent researches have advanced our understanding of the genetic components contributing to canine impaction [44]. Studies have identified specific genetic markers associated with increased risk, suggesting that early genetic screening could play a role in future diagnostic protocols [49]. Additionally, innovations in imaging technologies have enhanced our ability to detect and assess impacted canines with greater precision [54]. CBCT, in particular, has become a valuable tool in pre-surgical planning, allowing for detailed visualization of the impacted tooth's position relative to adjacent structures, which is critical for minimizing surgical risks [59].

Multidisciplinary treatment approaches

The management of impacted canines often necessitates a multidisciplinary approach, integrating orthodontic strategies to create space and guide eruption, alongside surgical procedures to expose or remove the impacted tooth [64]. Orthodontic techniques such as the use of palatal expanders and fixed appliances can be employed to create the necessary space for eruption, while surgical

options may include surgical exposure with orthodontic guidance, or in some cases, extraction [69]. The timing of intervention is critical, as early detection and treatment can significantly reduce the complexity and duration of therapy [77]. Early intervention, particularly in patients with a familial history of dental anomalies, can facilitate more conservative management options and improve prognostic outcomes [82].

Surgical management and post-operative considerations

Hospitalization for the surgical management of impacted canines is generally minimal, with most procedures performed on an outpatient basis [87]. However, the complexity of the impaction and the presence of complicating factors such as cyst formation or severe crowding may necessitate more extensive surgical interventions, potentially requiring short-term hospitalization [92]. Post-operative care is crucial to prevent complications such as infection or excessive bleeding and to ensure proper healing and integration of orthodontic devices [97].

Comparing treatment modalities

This study carefully examines the results of different treatment modalities, such as early genetic screening, CBCT-guided surgical planning, and integrated orthodontic-surgical treatment techniques, in order to improve our understanding of the most successful therapies for impacted canines [102]. Key variables like treatment length, hospitalization rates, post-operative complications, and overall patient satisfaction will be the focus of the study's evaluation of patient outcomes [107]. The importance of biological factors such as: age of the patient, position of the canine, presence or absence of crowding, and iatrogenic factors as the ultimate goal of treatment are critical to the choice between extraction of the impacted tooth and guided traction in the arch of the latter [112]. Each approach has distinct advantages, limitations, and potential complications that must be considered to optimize patient outcomes [117]. By analysing these factors, the study seeks to provide evidence-based guidance for clinical decision-making, enhancing the quality of care for patients with impacted canines and determining the most effective and patient-centred treatment approach [122].

Extraction as primary option

When canines are severely dislocated, surrounding structures are affected, or there is not enough arch space to allow for guided traction, extractive treatment is thought to be the best course of action [127]. Its main advantage is the immediate resolution of the problem, reducing the risk of infection, cyst formation, and further dental damage, and potentially shortening the overall treatment time

[132]. However, extraction can lead to aesthetic concerns and functional issues, necessitating prosthetic replacements to maintain the appearance and occlusal function [137].

Traction and its benefits

Traction involves surgically exposing the impacted canine and using orthodontic techniques to guide it into position, preserving the natural tooth and maintaining aesthetic and functional integrity [142]. This method, while beneficial for retaining a natural and appealing smile, and ensuring proper occlusion, is complex, time-consuming, and requires meticulous orthodontic management and patient compliance. Surgical exposure also carries risks such as infection and potential damage to surrounding structures [152]. These randomized controlled trials aim to comprehensively compare extraction and traction by evaluating outcomes like treatment duration, hospitalization rates, post-operative complications, patient satisfaction, and long-term functional and aesthetic results. By analysing these factors, the study seeks to provide evidence-based guidance for clinical decision-making, enhancing the quality of care for patients with impacted canines and determining the most effective and patient-centred treatment approach [157].

Study objectives

This article outlines the methodology and presents the initial findings of our study on the management of impacted canines. By synthesizing recent research and clinical practices, we aim to provide evidence-based recommendations to enhance the clinician's ability to effectively address this common yet complex dental issue. Through an exploration of genetic insights, advancements in imaging technology, and multidisciplinary treatment strategies, we hope to contribute to the ongoing efforts to improve patient outcomes and quality of care in the management of impacted canines.

Materials and methods

Protocol and registration

This review was carried out in compliance with PRISMA (Preferred Reporting Items for Systematic Reviews and

Meta-Analyses) criteria, and it was filed with the number CRD 577493 on PROSPERO (The International Prospective Register of Systematic Reviews) [163]. The primary research question of this review is to evaluate and compare the effectiveness of different surgical and orthodontic interventions for the management of impacted canines. Specifically, the review aims to assess the outcomes of extraction versus orthodontic traction in terms of treatment duration, patient satisfaction, post-operative complications, and overall success. In addition to the review, this article also deals with a series of clinical cases of canine disinclusion. Three patients between the ages of 9–14 (mean 11.5 years) with unerupted canines were selected and examined at the Department of Orthodontics, University of Bari. The participants and parents provided written informed consent to be involved in the study. Exclusion criteria included open and deep-bite, oral or systemic diseases, previous orthodontic treatment. The panoramic radiograph (a single-cut tomogram of the entire Maxilla and jaw) is the first radiographic record needed. The pre-treatment variables on the panoramic radiograph as alpha-angle (α), distance (d) and sector (s) were often evaluated [164]. The same method for worsening of mandibular canine was used.

Search processing

We limited our search to English-language papers published between January 1, 2004, and July 24, 2024, in PubMed, Scopus and Web of Science that were relevant to our topic. In the search approach, the Boolean keywords "impacted" AND "canine" were used. We chose these terms because they most correctly described our investigation's goal, which was to acquire greater knowledge into orthodontic traction of impacted canines in the arch or their extraction (Table 1).

Inclusion criteria

Three reviewers reviewed all relevant publications using the following criteria: (1) only human subjects research, (2) full text, and (3) scientific studies evaluating canine disengagement. The PICO model was created using the following steps:

- Criteria: Application in the present study;
- Population: Human subjects with impacted canine;

Table 1 Indicators for database searches

Articles screening strategy

KEYWORDS: "A": impacted; "B": canine;

Boolean Indicators: "A" AND "B"

Timespan: January 1, 2004, to July 24, 2024

Electronic databases: Pubmed; Scopus; Web of Science

- Intervention: Orthodontic treatment for to disinclude canines to bring them into the arch or extract them;
- Comparison: Control group;
- Outcome: Evaluation of successful canine disclusion;
- Study design: Randomized Controlled Trials, observational study

Exclusion criteria

Exclusion criteria included non-English-language articles, ineligible research designs, ineligible outcome measures, ineligible populations, case studies, reviews, and animal studies.

Data processing

Author conflicts about the choice of articles were addressed and resolved.

Article identification procedure

Two reviewers, E.I. and F.P., independently evaluated appropriateness. An extra manual search was performed to boost the number of articles available for full-text analysis. English-language articles that fit the inclusion requirements were considered, and duplicates and non-qualifying items were identified with a rationale for exclusion.

Study evaluation

The reviewers separately evaluated the article data using a specific electronic form established according to the following categories: authors, year of study, aim of the study, materials and methods, and results.

Quality assessment

Two reviewers, F.P. and E.I., utilized the ROBINS-I tool to evaluate the quality of the included articles. ROBINS-I was developed to assess the risk of bias in non-randomized studies that compare the health effects of two or more medications. Each of the seven assessed points was given a bias degree. E.I., the third reviewer, was contacted in the case of a disagreement until a resolution was reached. To improve the impartiality and uniformity of the assessments, any differences or disagreements among reviewers were handled via discussion and consensus-building. When there was no consensus, a third reviewer made the final choice. It helped to improve the overall assessment of the quality and dependability of the results by identifying the strengths and shortcomings of the evidence base. By accounting for the likelihood of bias, the authors of this review were able to draw more informed interpretations and conclusions based on the evidence presented.



Fig. 1 First panoramic radiograph



Fig. 2 Panoramic radiograph after odontoma removal

Case series

Case 1

FA, a nine-year-old girl, presented an unerupted mandibular canine. After six months of observation lower-right-canine deciduous was still present. Panoramic radiograph was prescribed. Odontome was present in the area of permanent lower-right canine that was submerged in bone, far from its eruptive path. Deciduous canine has not root resorption. So, surgical enucleation of composite odontome was performed, and eruption of canine was expected. On panoramic radiograph canine presented a α angle of 19° a distance (d) of 18 mm and was located in sector (s) 3. After two months a panoramic radiograph was prescribed (Fig. 1).

Position of permanent canine was worsened. Its transposition was observed, canine was near the mandibular midline. On second panoramic radiograph canine presented a α angle of 38° a distance (d) of 17 mm and was located in sector (s) 3 (Fig. 2).

So orthodontic treatment with fixed appliance was chosen to start orthodontic traction of the impacted cuspid. The treatment of the canine was concluded with its extraction (Fig. 3).

Case 2

DS, an eight-year-old girl, came in Department of Orthodontics, University of Bari for dental malocclusion

evaluation. Patients presented many caries on deciduous teeth so the panoramic and cranium radiographies were prescribed. Panoramic evaluation showed that both maxillary canines presented an alpha angle unfavourable to physiologic eruption. Upper left canine presented an alpha angle worse than the right: canine 1.3 presented an α angle of 23° a distance (d) of 22 mm and was located in sector (s) 3, while 2.3 presented an α angle of 25° a distance (d) of 24 mm and was located in sector (s) 3. On both maxillary sides premolars and canine lost the eruptive line (Fig. 4). The patient's parents decided to postpone the treatment for several months. Eight months later patients returned in Orthodontics Department and new radiographs were prescribed. Both canines worsen their position. An orthodontic treatment with a fixed appliance was proposed and accepted by the patient. Treatment started in upper arch bonding the brackets on four upper incisors and positioning two orthodontic bands on upper molars. After three months of treatment a radiographic control was done. Right canine further worsen its position. The left canine remained in its position. Radiograph evaluation showed the follow values for both canines: 1.3 α angle of 25° a distance (d) of 24 mm



Fig. 3 Operating sequence of mandibular right canine extraction



Fig. 4 First panoramic radiograph

and was located in sector (s) 3; 2.3 α angle of 54° a distance (d) of 24 mm and was located in sector (s) 3 (Fig. 5).

Case 3

AC, a twelve-year-old girl, came in Department of Orthodontics, University of for dental malocclusion evaluation. Radiographic evaluation was done. On panoramic radiography mandibular second premolars and two incisors agenesis was evaluated. The 2.3 showed a α angle of 53° a distance (d) of 25 mm and was located in sector (s) 2. The 1.3 presented a α angle of 49° a distance (d) of 29 mm and was located in sector (s) 1. In maxillary arch no agenesis was evaluated (Fig. 6).

On radiograph bilateral canines inclusion was present. Left canine had a worse position than right canine. Extraction of deciduous molars and fixed appliances on both arches (American Orthodontics, Sheboygan, WI, USA) was estimate. After three months of orthodontic treatment with fixed oral appliance a panoramic radiograph was prescribed before surgical treatment to hook both canines. On radiograph a worsening of left canine was evaluated. The 2.3 presented a α angle of 92° a distance (d) of 29 mm and was located in sector (s) 1; 1.3 presented an α angle of 32° a distance (d) of 11 mm and was located in sector (s) 2 (Fig. 7).



Fig. 5 Panoramic radiograph after three months of treatment



Fig. 6 First panoramic radiograph



Fig. 7 Panoramic radiograph where one can see the worsening of the position of the left canine

When imaging young patients, low-dose CBCT protocols should be implemented to minimize radiation exposure, in accordance with ALARA and ALADAIP principles, while still providing sufficient detail for procedures such as assessing impacted canines, as the latest CBCT devices offer effective low-dose imaging solutions [31].

Results

A total of 2748 papers were obtained from the databases Web of Science (796), PubMed (740), and Scopus (1212). This resulted in 1455 articles after eliminating duplicates (1293). 897 entries were eliminated after their titles and abstracts were examined. The writers were able to successfully obtain the remaining 558 papers and confirm their eligibility. 543 items were eliminated as a result of this process because they were off topic. The qualitative analysis of the 15 final articles is included in this study (Fig. 8). Each study's findings were presented in Table 2.

Quality assessment and risk of bias of included articles

Figures 9 and 10, showing the risk of bias in the included studies assessed using RoB 2 for randomized trials and ROBINS-I for the observational study. Figure 9 highlights the risk of bias in the included randomized controlled

trials (RCTs). Most studies exhibit a high risk of bias due to issues in the randomization process and deviations from intended interventions. Measurement bias generally shows a low risk, indicating reliable outcome measurements. Many studies also show minimal bias from missing data. However, bias in the selection of reported outcomes is prevalent across most RCTs, impacting the robustness of their findings. Overall, the findings indicate that few RCTs maintain a low risk of bias, while several exhibit moderate to high risks.

Figure 10, illustrating the risk of bias in the observational study, shows that there is a serious risk of bias due to confounding and measurement of outcomes. Selection bias and deviations from intended interventions generally display low-risk levels, indicating adequate participant selection and adherence to intervention protocols. The observational study reveals moderate to serious bias levels overall, with a significant risk due to confounding and measurement inaccuracies, which may impact the interpretation of findings.

Discussion

Impacted canine control is an integral part of orthodontic treatment and involves a delicate balance between effectiveness, safety, and comfort for the patient. Impacted canines are relatively common dental anomalies that create a variety of diagnostic and treatment challenges for an orthodontist [180]. Recent advances in technology and treatment philosophies have altered the approach to managing impacted canines and added new and improved solutions to the clinician's armamentarium [185].

The most common forms of treating impacted canines include extraction and traction. This review compares the two approaches based on the most recent and best research and available clinical outcomes about their relative effectiveness and impact on patient care.

Technological advances in orthodontics

Computer-aided techniques significantly improved the safety and accuracy of orthodontic surgical intervention. Kivovics et al. (2024) confirmed the advanced presurgical planning process with individual presurgical guides to increase precision and safety, especially in complex cases [166]. Moreover, Björksved et al. (2021) stated that three-D imaging with pc-guided planning has better results in cases of hard anatomical systems since it reduces complications and gives better surgical effects [179]. The minimally invasive strategies in orthodontics have also advanced and involve laser technology [195]. According to Impellizzeri et al. in 2023, laser strategies have been found to lower bleeding and post-operative ache by approximately 30% and 20%, correspondingly, compared




Fig. 8 PRISMA flowchart of the literature search and article inclusion process

to traditional techniques [40]. These strategies are also found to promote faster recovery and decrease the requirement of analgesia, thereby improving the comfort and recovery of patients [200]. This, therefore, becomes part of less invasive approaches in the best interest of patient well-being and satisfaction [204]. Corticotomy-assisted treatments were evaluated for their ability to increase tooth motion [207]. Mousa et al. observed that corticotomy could decrease orthodontic treatment time by approximately 40% [14]. However, Verma et al. found that this technique initiates regional acceleratory phenomenon, resulting in increased tooth movement with no evidence of periodontal health impairment [170]. These studies exemplify the changing face of orthodontic treatments, with a search toward more patient-friendly options.

Comparison of extraction and traction techniques

Impacted canines are managed by several techniques, including extraction and traction methods. A study by Hadler-Olsen et al. (2020) on the management of the palatally displaced canine compared the single and double extraction of primary teeth [174]. No significant difference in eruption rates was noted between these

two methods. Both techniques proved useful, but critical factors toward success were initial canine angulation and space conditions [212]. Naoumova et al. (2015) reviewed deciduous canine extraction and noted that eruption rates in the extraction group were significantly greater than in the control group at 80% compared with 50% after 12 months, thereby supporting deciduous canine extraction as an effective interceptive measure [52]. Antunes et al. (2018) evaluated the effect of alveolar bone grafting on the eruption of impacted canines in patients with cleft lip and palate [176]. According to this study, while bone grafting usually provided for spontaneous eruption, 33.33% of them still needed orthodontic traction, emphasizing the need for individualized treatment plans. In a 2019 study, Parkin et al. compared the aesthetic and functional outcomes using closed and open surgical techniques for impacted canines [110]. While minor differences in the aesthetics of both closed and open surgical techniques have been noted, closed techniques involve less postoperative pain and promote faster recovery, making them more favorable for reducing discomfort [220]. Björksved et al. (2021) supported this by indicating that open surgical techniques demonstrated increased postoperative pain and functional impairment,

Table 2 A descriptive item selection summary

Authors	Type of study	Aim of the study	Materials and Methods	Results	Sample Sizes	Study Groups	Clinical Parameters for Effectiveness
M. Kivovics et al. (2024) [166]	randomized clinical trial	To assess the feasibility of computer-assisted open exposure of palatally impacted canines (PICs)	Patients aged 11–30 years who required orthodontic eruption for the full palatal impaction of their canines	Computer-assisted surgery is a feasible method for open exposure of PICs, which decreases the duration of surgery compared to the free-hand method	34 participants	Orthodontic patients requiring exposure of palatally impacted canines	Surgery duration compared between computer-assisted and free-hand methods; shorter times indicated higher feasibility
M. Björksved et al. (2021) [167]	randomized clinical trial	To evaluate the costs of open and closed surgical exposure and subsequent orthodontic treatment for the correction of palatally displaced canines (PDCs)	One hundred twenty adolescents between 9 and 16 years of age. Cost analysis was performed including costs for surgery, orthodontic treatment, emergency visits and material, as well as costs for transports and time spent in connection with every appointment	The two different surgical exposures and following orthodontic treatments did not differ significantly in terms of costs	120 participants	Randomized into two groups: open vs. closed surgical exposure	Cost parameters included surgery, treatment, and other related expenses to assess overall cost-effectiveness of both methods
A. Impellizzeri et al. (2023) [168]	randomized clinical trial	To show the effectiveness of laser technology for the exposure of PICs, using a CO2 or diode laser, and to evaluate the possible bio-stimulation effect of the laser on the spontaneous eruption of the canine	27 patients, divided randomly into three groups: treated with a CO2 laser (Group A), treated with a diode laser (Group B), and treated with a cold blade (Group C). Monitoring was performed at 1, 8, and 16 weeks after surgery, through photo and digital scans performed with a CS3500 intraoral scanner	The average total eruptions are 4.55 mm for Group A, 5.36 mm for Group B, and 3.01 mm for Group C. The difference in eruption between groups A and B is not significant. Comparing the laser groups with the control group, it has emerged that the difference in eruption is statistically significant	27 participants (3 groups)	Group A: CO2 laser, Group B: Diode laser, Group C: Cold blade	Canine eruption measured in millimeters; significantly higher eruption in laser groups compared to the control (cold blade group)

Table 2 (continued)

Authors	Type of study	Aim of the study	Materials and Methods	Results	Sample Sizes	Study Groups	Clinical Parameters for Effectiveness
M. R. Mousa et al. (2023) [169]	randomized clinical trial	To evaluate the effectiveness of a minimally-invasive corticotomy-assisted (MICA) treatment of PICs compared with the traditional method by evaluating the velocity of movement, and the associated dentoalveolar changes	Forty-six patients with upper PICs were distributed into two groups: the corticotomy-assisted traction group and the traditional treatment group. The closed surgical approach was used in both study groups	The traction movement velocity of the PICs can be increased using MICA orthodontic treatment. The side effects of the acceleration procedure were minimal and almost similar to those of the traditional technique	46 participants	Group 1: MICA-assisted traction, Group 2: Traditional treatment	Velocity of canine movement measured; faster movement with MICA technique indicated higher effectiveness. Minimal side effects noted
S. Verma et al. (2023) [170]	randomized clinical trial	To evaluate the success rate, treatment duration, and pain perceived during forced eruption of maxillary PICs using the K9 and Ballista springs	30 unilateral PICs of moderate and difficult category were enrolled. Group 1 comprised canine impacted springs, and Group 2 comprised Ballista springs	The IC of moderate and difficult category were erupted with a 100% success rate and similar treatment duration with both interventions. The pain scores of Ballista springs were greater after 24 h of force application	30 patients	Group 1 (K9 springs) vs Group 2 (Ballista springs)	Success measured by eruption rate and pain perception, with Ballista springs causing higher pain scores while both methods achieved full eruption
M. Björksved et al. (2018) [171]	randomized clinical trial	To compare treatment time, patients' perceptions during orthodontic treatment, dental fear and side effects, between open and closed surgical exposures in patients with PDCs	125 patients from three different orthodontic centers were randomized into one of the two intervention arms, open or closed surgical exposure	The closed exposure group experienced more pain and discomfort mostly during active orthodontic traction	125 patients	Two groups: open surgical exposure vs closed surgical exposure	Patient perception and discomfort during orthodontic traction were primary parameters, with the closed exposure group reporting higher pain levels
J. Cunha-Cruz (2015) [172]	randomized clinical trial	To determine the best course of action for an 11-year-old girl with suspected impacted using the five steps of Evidence-based practice: Ask, Acquire, Appraise, Apply, and Assess	The orthodontist selected three studies: a case report, a cohort study, and a randomized controlled trial (RCT). The RCT was chosen for its lower risk of bias, as it randomly allocated patients to treatment and control groups	Extraction of DC increased the likelihood of permanent canine eruption by 80%	Single case study	N/A	Efficacy measured by eruption probability, with an 80% increase in likelihood of permanent canine eruption following deciduous tooth extraction

Table 2 (continued)

Authors	Type of study	Aim of the study	Materials and Methods	Results	Sample Sizes	Study Groups	Clinical Parameters for Effectiveness
H. Jang et al. (2021) [173]	randomized clinical trial	To investigate the effects of palatal stents in patients who underwent supernumerary tooth removal through a palatal approach and to suggest the optimal stent thickness and material	144 patients were divided into a control group and compressive palatal stent group each distinguished by the thickness and material of the thermoplastic acrylic stent used	Compressive palatal stents reduce discomfort by decreasing pain and alleviating swelling. Although a stent is effective regardless of its thickness and material, 2-mm hard stents maximized such positive effects with minimal discomfort	144 patients	Control group vs compressive palatal stent group	Effectiveness measured by reduction in discomfort, pain, and swelling. The 2-mm hard stents were found to provide optimal results
Hadler-Olsen et al., (2020) [174]	Randomized Controlled Trial	Compare effectiveness of double vs single primary tooth extraction for PDCs	32 children with 48 PDCs; PDCs were split into two groups at random: the double-extraction group (DEG) and the single-extraction group (SEG). Until the canine emerged or orthodontic treatment was initiated, logistic regression was used to examine the factors impacting PDC emergence at baseline and six-month intervals. Clinical and radiographic exams emergence (yes/no), emergence into a favorable position (yes/no), and MC positional shift (angulation and sector) were the outcome measures	No significant difference in PDC emergence; both groups showed significant distal movement. In order to assist PDC eruption into the oral cavity and into a desirable position in the dental arch, single or double primary tooth extraction methods are similar. Early evaluations of canine angulation and spacing may be utilized to forecast the success of PDC eruption	32 children (48 PDCs)	DEG (double extraction) vs SEG (single extraction)	Canine angulation and spacing were key clinical parameters determining the success of canine eruption following primary tooth extraction

Table 2 (continued)

Authors	Type of study	Aim of the study	Materials and Methods	Results	Sample Sizes	Study Groups	Clinical Parameters for Effectiveness
Parkin et al., (2013) [175]	Randomized Controlled Trial	Evaluate esthetic outcomes of closed vs open surgical exposure of PDCs	The study included 67 participants, with laypeople identifying the operated tooth only 49.7% of the time, and orthodontists identifying the treated canine with certainty only 60.7%. Both panels assessed the unoperated canine's appearance better than the contralateral operated canine, but no differences were found	Aligning a PDC has an aesthetic impact, but it is largely subtle and unlikely to be seen by the general public. Regardless of whether the canine was made visible through an open or closed surgical procedure, the aesthetic impression was the same	67 participants	Closed vs open surgical exposure	Aesthetic outcomes were judged by laypeople and professionals. No significant differences were observed between closed and open procedures in terms of aesthetics
Antunes C. et al. (2018) [176]	Observational Study	Investigate eruption patterns of impacted teeth after alveolar bone graft in cleft lip and/or palate (CLP) patients	Between 2004 and 2011, 42 patients were treated at a facility, with the most impacted being 9–11 year olds. The most common cleft was unilateral (85.71%). The most common impacted tooth was the MC, and orthodontic traction was required for 64.3 and 35.7% respectively	Higher incidence of CLP in males; Following the implantation of the bone graft, all impacted teeth in the preforamen incisor cleft spontaneously erupted. The canine teeth were the most common, primarily associated with the TIC. IC spontaneously erupted during secondary alveolar bone grafting in CLP	42 CLP patients	Single group (CLP patients with impacted teeth)	Effectiveness measured by spontaneous eruption following alveolar bone grafting, with maxillary canines being the most commonly impacted and responsive to treatment

Table 2 (continued)

Authors	Type of study	Aim of the study	Materials and Methods	Results	Sample Sizes	Study Groups	Clinical Parameters for Effectiveness
Naoumova et al., (2015) [177]	Prospective Randomized Clinical Longitudinal Trial	Evaluate effect of DC extraction on PDC eruption	67 children aged 10–13 with maxillary unilateral or bilateral periodontal disease, persisting DC and no previous experience of orthodontic treatment. The patients were randomly assigned to extraction or non-extraction, underwent clinical examinations and cone beam computed tomography at baseline, 6 months, and 12 months. Outcome measures included eruption, positional changes, eruption duration, and root resorption. Extraction group vs control group; monitored radiographic changes over time	The extraction group experienced more spontaneous eruptions of PDCs than the control group, with rates of 69% and 39%, respectively. Differences were found in all variables except for the sagittal angle, with larger changes in canine cusp-tip distances and apex changes	67 children	Extraction group vs non-extraction control group	Spontaneous eruption rate and positional changes in the canine cusp and apex were measured to determine effectiveness of DC extraction for PDC management
Mousa et al., (2023) [14]	Randomized Controlled Trial	Compare periodontal outcomes of MICA technique vs conventional traction for PDCs	Individuals with unilaterally affected canines participated in a randomized controlled experiment to examine periodontal characteristics. Accelerated therapy (CAT) and conventional treatment (CT) groups were created for the participants. Two weeks following the fixed orthodontic appliance's removal, post-treatment measurements were obtained	The study involved 46 patients in either the CT or CAT groups for expedited therapy. The differences in periodontal outcomes were not statistically significant. The combined PPD values for neighboring teeth were almost identical, and the two study groups had very low gingival recession level (GRL) values	46 patients	Group 1 (MICA) vs Group 2 (conventional traction)	Periodontal outcomes, specifically probing pocket depth (PPD) and gingival recession level (GRL), were measured to assess treatment effectiveness, with no significant differences noted between groups

Table 2 (continued)

Authors	Type of study	Aim of the study	Matherials and Methods	Results	Sample Sizes	Study Groups	Clinical Parameters for Effectiveness
Moussa et al., (2022) [178]	Randomized Controlled Trial	Assess patient-centered outcomes of conventional vs accelerated minimally invasive corticotomy-assisted orthodontic treatment for PDCs	A study of 52 patients with unilateral PICs, divided into two groups, measured pain, discomfort, and functional issues using a visual analog scale 24 h, 4 days, 7 days, 14 days, and 28 days after the surgical exposure procedure	The study did not find any statistically significant differences between the two therapy groups. On the first day following surgery, pain and discomfort were somewhat greater in the traditional group compared to the corticotomy-assisted group. Four days were needed for recovery from edema, mastication issues, swallowing issues, and speech abnormalities; whereas seven days were needed for other outcomes	52 patients	Group 1 (minimally invasive corticotomy-assisted) vs Group 2 (conventional treatment)	Patient-centered outcomes, including pain and discomfort levels, were tracked post-surgery. Minimally invasive corticotomy-assisted therapy resulted in marginally better pain management early on
Biörksved et al., (2021) [179]	Multicenter Randomized Controlled Trial	Compare postoperative outcomes of closed vs open surgical exposure of PDCs	119 consecutive patients from three orthodontic centers who underwent surgical exposure. The procedures involved elevating a full-thickness mucoperiosteal flap, removing bone covering, and suturing the flap back. To prevent gingival overgrowth, a tissue window was excised and glass ionomer cement was applied. Patient perceptions were assessed using questionnaires	Open technique associated with more postoperative pain and impairment; no significant difference in surgery time	119 patients	Group 1 (closed surgical exposure) vs Group 2 (open surgical exposure)	Postoperative pain and impairment were the primary clinical parameters, with the open surgical exposure group reporting more pain and functional limitations compared to the closed exposure group




Fig. 9 Bias assessment by RoB 2 for randomized trials




Fig. 10 Bias assessment by ROBINS-I for the observational study

especially in bilateral cases, and thus closed technique had to be preferred since it causes less pain and has fewer major complications [167]. Mousa et al. (2023) reviewed the periodontal outcomes of corticotomy-assisted versus conventional closed-flap traction, finding no significant differences in periodontal health parameters between the

two modalities, with minimal negative effects on periodontal tissues [169]. These findings have been supported by Verma et al. (2023), which stated that corticotomy-assisted techniques surely accelerate tooth movement significantly and positively, with the maintenance of good periodontal health, and therefore turn into a feasible

alternative to conventional techniques of traction [170]. Cunha-Cruz et al. in 2015 and Jang et al. in 2021 further emphasized the role of evidence-based practice in orthodontics, advocating for the integration of available evidence into clinical practice to make informed treatment decisions and improve patient outcomes [172]. Björksved et al. (2021) highlighted the significance of patient-centered care, stating that treatment should be based on patient preference and clinical concerns [179]. From a perspective of individualized care, this care should be oriented toward fulfilling individual needs and expectations for improved results and satisfaction [226].

Key insights and clinical implications

Studies have shown that extraction of primary teeth, especially the deciduous canines, as discussed in Naoumova et al., greatly facilitates the eruption of impacted canines [76]. The simplicity and minimal invasiveness of the technique associated with high success rates and a low complication rate make it an option whose efficiency may differ in different patients; therefore, treatment must be planned on an individual basis [231]. On the contrary, traction methods, particularly corticotomy-assisted techniques following surgical intervention, are potent alternatives that do not seem to compromise periodontal health. These are useful in patients in whom simple extraction may prove inadequate. According to Mousa et al., conventional and accelerated traction methods are comparable in terms of comfort and periodontal outcomes, with no significant differences in pain or functional impairment [178]. The results from Björksved et al. study, on closed vs. open surgical exposure techniques, outline the benefits of the former in terms of less postoperative pain and fewer functional impairments [171]. This is important information to have at hand when being faced with clinical decisions that involve patient comfort and recovery as major concerns. Parkin et al. commented on the aesthetics-related outcomes of the techniques involved in their study, outlining that both methods have similar effects concerning aesthetics, with slight advantages of the closed techniques being possible in the clinical setting [113]. This reiterates the incorporation of variables specific to the patient, such as aesthetic priorities and general comfort, in order to decide upon which approach is best for every patient. Timing of intervention also had its importance, from the study by Antunes et al., wherein effectiveness in alveolar bone grafting for patients with cleft lip and palate indicated benefits toward improved eruption outcomes [176]. This suggests that in order to maximize patient outcomes, orthodontic treatment planning must be done in a timely and planned manner.

In conclusion, the comparison between extraction and traction techniques in orthodontics delineates the fact

that several factors, like patient preferences, clinical conditions, and evidence-based practice, have to be considered in order to arrive at the most appropriate and effective treatment plan. The trend toward less invasive and more patient-friendly techniques, such as corticotomy-assisted treatments and closed surgical methods, is one of improvement in patient comfort and recovery with successful outcomes.

Limitations and future directions

The majority of orthodontic research is restricted by a number of critical limitations that significantly compromise the reliability and practicality of findings [236]. First, many studies have small sample sizes, with short follow-up, similar to those conducted by Hadler-Olsen et al. in 2020 and Naoumova et al. in 2015, that could compromise the ability to detect long-term problems and generalize findings [174]. Additionally, there is significant variability in study designs and protocols, making it difficult to compare results across different studies, as evidenced by the discrepancies between the methods utilized by Parkin et al. (2013) and Björksved et al. (2021) [175]. The subjectivity in measurements of outcomes, mainly related to aesthetic assessments, implies possible bias and inconsistency in the results obtained, which was actually confirmed by Parkin et al. (2013) [175]. Moreover, there is a lack of literature regarding the effectiveness of various techniques and related complications in the long term. For instance, Mousa et al. (2023) present short-term results [169]. In this context, the response of each patient to, for example, bone grafting or application of traction is very different, as stated by Antunes et al. (2018), hence too subjective to predict the outcomes, and thus requires a specific and unique treatment plan in each case [176].

Future studies should give priority to the extension of works involving larger samples with diversified populations to increase the generalizability of findings [239]. Long-term, multi-center trials with an extension of follow-up may explore durability and efficacy accruable to various treatment techniques [242]. Also, standardization of the research protocols is needed regarding uniformity in standards for outcome assessments with respect to aesthetic and functional measures and standardized pain and discomfort rating scales. Future prospects include developments in imaging technology that presently incorporate 3D CBCT. Patient-centered outcomes, such as quality of life, satisfaction with treatment, and functional and aesthetic results in the long term, are parameters that should be monitored to keep the patient close to the treatment plan [146]. Investigating combined treatment techniques may also reveal new insights into optimizing care, and developing predictive models using data

analysis and machine learning could lead to more personalized and efficient treatment planning. Addressing these limitations and future directions will drive progress in orthodontics, ultimately enhancing treatment options for impacted canines and improving patient care.

Conclusions

This systematic review highlights the critical factors in the management of impacted canines, emphasizing the importance of tailored treatment approaches. Both extraction and orthodontic traction are effective options, each with specific advantages depending on the severity of impaction and patient-specific factors. While extraction provides a faster solution and reduces risks of complications, it may lead to aesthetic and functional issues requiring prosthetic solutions. On the other hand, traction, though more complex and time-consuming, preserves the natural tooth and aesthetic integrity. Advances in surgical techniques, such as the use of laser technology and minimally invasive corticotomy, have improved outcomes by reducing surgical time and enhancing the speed of tooth eruption, with minimal side effects. The integration of advanced imaging technologies like CBCT has also enhanced diagnostic accuracy and surgical planning, reducing risks and improving patient outcomes. The review underscores the necessity of a patient-centered, multidisciplinary approach to treatment, considering the multifactorial etiology of canine impaction, which includes both genetic and environmental factors. Early genetic screening may play a future role in predicting impaction risks and guiding timely interventions. The findings advocate for further research, particularly randomized controlled trials, to refine treatment selection criteria and incorporate emerging genetic and technological advancements into clinical practice.

Abbreviations

α	Alpha-angle
CBCT	Cone-beam computed tomography
CLP	Cleft lip and/or palate
IC	Impacted canines
d	Distance
DC	Deciduous canine
DEG	Double-extraction group
GRL	Gingival recession level
PIC	Palatally impacted canines
PDC	Palatally displaced canines
PPD	Periodontal pocket depth
PRISMA	Preferred Reporting Items for Systematic Reviews and Meta-Analyses
PROSPERO	The International Prospective Register of Systematic Reviews
RCT	Randomized controlled trial
s	Sector
SEG	Single-extraction group

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s12903-025-05635-w>.

Supplementary Material 1.

Acknowledgements

Not applicable.

Authors' contributions

Conceptualization, A.M. and F.C.; methodology, A.M.I. and C.C.; software, V.C.; validation, V.C., P.M. and C.C.; formal analysis, A.D.I. and P.M.; investigation, F.I. and F.P.; resources, G.D. and G.C.; data curation, F.I.; writing—original draft preparation, G.D. and G.C.; writing—review and editing, F.P.; visualization, A.D.I.; supervision, A.M.I.; project administration, F.C.; funding acquisition, A.M.. All authors have read and agreed to the published version of the manuscript.

Funding

This research received no external funding.

Data availability

No datasets were generated or analysed during the current study.

Declarations

Ethics approval and consent to participate

This study was approved by Ethics Committee of "Azienda Ospedaliero Universitaria Consorziale Policlinico", Bari, Italy (0015987, 17 February 2022).

Competing interests

The authors declare no competing interests.

Author details

¹Department of Interdisciplinary Medicine, University of Bari "Aldo Moro", 70124 Bari, Italy. ²Multidisciplinary Department of Medical-Surgical and Dental Specialties, Oral and Maxillofacial Surgery Unit, University of Campania "Luigi Vanvitelli", 80138 Naples, Italy.

Received: 14 August 2024 Accepted: 11 February 2025

Published online: 24 February 2025

References

- Kwon HK, Owen DG, Davidson WM. A case of supernumerary teeth in the premaxilla, maxillary cuspid, and mandibular premolar regions. *MSDA J*. 1998;41:105–8.
- Lempesi E, Pandis N, Fleming PS, Mavragani M. A Comparison of apical root resorption after orthodontic treatment with surgical exposure and traction of maxillary impacted canines versus that without impactions. *Eur J Orthod*. 2014;36:690–7. <https://doi.org/10.1093/ejo/cjt099>.
- Baccetti T. A controlled study of associated dental anomalies. *Angle Orthod*. 1998;68:267–74. [https://doi.org/10.1043/0003-3219\(1998\)068%3c0267:ACSOAD%3e2.3.CO;2](https://doi.org/10.1043/0003-3219(1998)068%3c0267:ACSOAD%3e2.3.CO;2).
- Koo TK, Li MY. A guideline of selecting and reporting intraclass correlation coefficients for reliability research. *J Chiropr Med*. 2016;15:155–63. <https://doi.org/10.1016/j.jcm.2016.02.012>.
- Brown LH, Berkman S, Cohen D, Kaplan AL, Rosenberg M. A radiological study of the frequency and distribution of impacted teeth. *J Dent Assoc S Afr*. 1982;37:627–30.
- Mason C, Azam N, Holt RD, Rule DC. A Retrospective study of unerupted maxillary incisors associated with supernumerary teeth. *Br J Oral Maxillofac Surg*. 2000;38:62–5. <https://doi.org/10.1054/bjom.1999.0210>.

7. Litsas G, Acar A. A review of early displaced maxillary canines: etiology, diagnosis and interceptive treatment. *Open Dent J*. 2011;5:39–47. <https://doi.org/10.2174/1874210601105010039>.
8. Richardson G, Russell KA. A review of impacted permanent maxillary cuspids-diagnosis and prevention. *J Can Dent Assoc*. 2000;66:497–501.
9. Bedoya MM, Park JH. A review of the diagnosis and management of impacted maxillary canines. *J Am Dent Assoc*. 2009;140:1485–93. <https://doi.org/10.14219/jada.archive.2009.0099>.
10. Dachi SF, Howell FV. A survey of 3,874 routine full-mouth radiographs. I. A study of retained roots and teeth. *Oral Surg Oral Med Oral Pathol*. 1961;14:916–24. [https://doi.org/10.1016/0030-4220\(61\)90003-2](https://doi.org/10.1016/0030-4220(61)90003-2).
11. Naoumova J, Kuroi J, Kjellberg H. A systematic review of the interceptive treatment of palatally displaced maxillary canines. *Eur J Orthod*. 2011;33:143–9. <https://doi.org/10.1093/ejo/cjq045>.
12. Zeitoun IM, Dhanrajani PJ, Mosadomi HA. Adenomatoid odontogenic tumor arising in a calcifying odontogenic cyst. *J Oral Maxillofac Surg*. 1996;54:634–7. [https://doi.org/10.1016/s0278-2391\(96\)90650-3](https://doi.org/10.1016/s0278-2391(96)90650-3).
13. Bravo M, White D, Miles L, Cotton R. Adenomatoid odontogenic tumor mimicking a dentigerous cyst. *Int J Pediatr Otorhinolaryngol*. 2005;69:1685–8. <https://doi.org/10.1016/j.ijporl.2005.03.055>.
14. Mousa MR, Hajeer MY, Burhan AS, Heshmeh O. Adult Periodontal comparison after treatment of palatally impacted canines aligned by conventional or accelerated minimally-invasive corticotomy-assisted orthodontic traction: a randomized controlled trial. *Int Orthod*. 2023;21:100785. <https://doi.org/10.1016/j.ortho.2023.100785>.
15. Beckmann SH, Kuitert RB, Prahli-Andersen B, Segner D, The RP, Tuinzing DB. Alveolar and skeletal dimensions associated with lower face height. *Am J Orthod Dentofacial Orthop*. 1998;113:498–506. [https://doi.org/10.1016/s0889-5406\(98\)70260-4](https://doi.org/10.1016/s0889-5406(98)70260-4).
16. Kim Y, Hyun HK, Jang KT. The position of maxillary canine impactions and the influenced factors to adjacent root resorption in the Korean population. *Eur J Orthod*. 2012;34:302–6. <https://doi.org/10.1093/ejo/cjr002>.
17. Kulldorff M, Huang L, Pickle L, Duczmal L. An elliptical spatial scan statistic. *Stat Med*. 2006;25:3929–43. <https://doi.org/10.1002/sim.2490>.
18. Inchingolo F, Dipalma G, Azzollini D, Trilli I, Carpentiere V, Hazbala D, Bordea IR, Palermo A, Inchingolo AD, Inchingolo AM. Advances in preventive and therapeutic approaches for dental erosion: a systematic review. *Dent J (Basel)*. 2023;11:274. <https://doi.org/10.3390/dj11120274>.
19. Philipsen HP, Reichart PA, Siar CH, Ng KH, Lau SH, Zhang X, Dhanuthai K, Swadison S, Jankittivong A, Meer S, et al. An updated clinical and epidemiological profile of the adenomatoid odontogenic tumour: a collaborative retrospective study. *J Oral Pathol Med*. 2007;36:383–93. <https://doi.org/10.1111/j.1600-0714.2007.00536.x>.
20. Tenzin, null; Dhand, N.K.; Ward, M.P. Anthropogenic and Environmental Risk Factors for Rabies Occurrence in Bhutan. *Prev Vet Med* 2012, 107, 21–26. <https://doi.org/10.1016/j.prevetmed.2012.05.003>.
21. Lund H, Gröndahl K, Hansen K, Gröndahl HG. Apical root resorption during orthodontic treatment. A prospective study using cone beam CT. *Angle Orthod*. 2012;82:480–7. <https://doi.org/10.2319/061311-390.1>.
22. Brusveen EMG, Brudevik P, Bøe OE, Mavragani M. Apical root resorption of incisors after orthodontic treatment of impacted maxillary canines: a radiographic study. *Am J Orthod Dentofacial Orthop*. 2012;141:427–35. <https://doi.org/10.1016/j.ajodo.2011.10.022>.
23. Inchingolo F, Inchingolo AM, Latini G, Palmieri G, Di Pede C, Trilli I, Ferrante L, Inchingolo AD, Palermo A, Lorusso F, et al. Application of graphene oxide in oral surgery: a systematic review. *Materials (Basel)*. 2023;16:6293. <https://doi.org/10.3390/ma16186293>.
24. Di Stasio D, Romano A, Paparella RS, Gentile C, Serpico R, Minervini G, Candotto V, Laino L. How social media meet patients' questions: YouTube™ review for mouth sores in children. *J Biol Regul Homeost Agents*. 2018;32:117–21.
25. Di Stasio D, Lauritano D, Romano A, Salerno C, Minervini G, Minervini G, Gentile E, Serpico R, Lucchese A. In vivo characterization of oral pemphigus vulgaris by optical coherence tomography. *J Biol Regul Homeost Agents*. 2015;29:39–41.
26. Mousa M, Hajeer M, Burhan A, Heshmeh O, Darwich K. Assessment of patient-centered outcomes when treating palatally impacted canines using conventional versus accelerated minimally invasive corticotomy-assisted orthodontic treatment: a randomized controlled trial. *Cureus*. 2022;14:e30392. <https://doi.org/10.7759/cureus.30392>.
27. Kolokitha O-E, Balli D, Zarkadi A-E, Gizani S. Association between maxillary canine impaction and other dental anomalies: radiological study of a mixed dentition children's cohort from an orthodontic clinic. *Eur Arch Paediatr Dent*. 2023;24:401–7. <https://doi.org/10.1007/s40368-023-00798-y>.
28. Malcangi G, Patano A, Ciocia AM, Netti A, Viapiano F, Palumbo I, Trilli I, Guglielmo M, Inchingolo AD, Dipalma G, et al. Benefits of natural antioxidants on oral health. *Antioxidants (Basel)*. 2023;12:1309. <https://doi.org/10.3390/antiox12061309>.
29. Inchingolo AM, Patano A, De Santis M, Del Vecchio G, Ferrante L, Morolla R, Pezzolla C, Sardano R, Dongiovanni L, Inchingolo F, et al. Comparison of different types of palatal expanders: scoping review. *Children (Basel)*. 2023;10:1258. <https://doi.org/10.3390/bambini10071258>.
30. Bozkurt SB, Hakki EE, Kayis SA, Dundar N, Hakki SS. Biostimulation with diode laser positively regulates cementoblast functions, in vitro. *Lasers Med Sci*. 2017;32:911–9. <https://doi.org/10.1007/s10103-017-2192-z>.
31. Scarfe WC, Farman AG, Sukovic P. Clinical applications of cone-beam computed tomography in dental practice. *J Can Dent Assoc*. 2006;72:75–80.
32. Umemura S, Yamaguchi T, Suginome K, Nishiguchi S. Clinical investigation of orthodontic treatment of palatally impacted canines, using ballista spring appliance. *Nippon Kyosei Shika Gakkai Zasshi*. 1990;49:369–78.
33. Bishara SE. Clinical management of impacted maxillary canines. *Semin Orthod*. 1998;4:87–98. [https://doi.org/10.1016/s1073-8746\(98\)80006-6](https://doi.org/10.1016/s1073-8746(98)80006-6).
34. Keiko M, Echigo S, Kimizuka S, Takahashi M, Chiba M. Clinical study on eruption of permanent canines after secondary alveolar bone grafting. *Cleft Palate Craniofacial J*. 2005;42:309–13. <https://doi.org/10.1597/03-113.1>.
35. Inchingolo AD, Dipalma G, Ferrara I, Viapiano F, Netti A, Ciocia AM, Mancini A, Malcangi G, Palermo A, Inchingolo AM, et al. Clear aligners in the growing patient: a systematic review. *Children (Basel)*. 2024;11:385. <https://doi.org/10.3390/children11040385>.
36. Di Stasio D, Lauritano D, Gritti P, Migliozi R, Maio C, Minervini G, Petruzzi M, Serpico R, Candotto V, Lucchese A. Psychiatric disorders in oral lichen planus: a preliminary case control study. *J Biol Regul Homeost Agents*. 2018;32:97–100.
37. Di Stasio D, Romano A, Gentile C, Maio C, Lucchese A, Serpico R, Paparella R, Minervini G, Candotto V, Laino L. Systemic and Topical Photodynamic Therapy (PDT) on oral mucosa lesions: an overview. *J Biol Regul Homeost Agents*. 2018;32:123–6.
38. Lucchese A, Dolci A, Minervini G, Salerno C, Di Stasio D, Minervini G, Laino L, Silvestre F, Serpico R. Vulvovaginal Gingival Lichen Planus: Report of Two Cases and Review of Literature. *Oral Implantsol (Rome)* 2016, 9, 54–60. <https://doi.org/10.11138/orl/2016.9.2.054>.
39. Leon JE, Mata GM, Fregnani ER, Carlos-Bregini R, de Almeida OP, Mosqueda-Taylor A, Vargas PA. Clinicopathological and immunohistochemical study of 39 cases of adenomatoid odontogenic tumour: a multicentric study. *Oral Oncol*. 2005;41:835–42. <https://doi.org/10.1016/j.oraloncology.2005.04.008>.
40. Impellizzeri A, Horodyski M, Serritella E, Palaia G, De Stefano A, Polimeni A, Galluccio G. Uncovering and autonomous eruption of palatally impacted canines—a case report. *Dent J (Basel)*. 2021;9:66. <https://doi.org/10.3390/dj9060066>.
41. Cavuoti S, Matarese G, Isola G, Abdolreza J, Femiano F, Perillo L. Combined orthodontic-surgical management of a transmigrated mandibular canine. *Angle Orthod*. 2016;86:681–91. <https://doi.org/10.2319/050615-309.1>.
42. Janson G, Bombonatti R, Brandão AG, Henriques JFC, de Freitas MR. Comparative radiographic evaluation of the alveolar bone crest after orthodontic treatment. *Am J Orthod Dentofacial Orthop*. 2003;124:157–64. [https://doi.org/10.1016/s0889-5406\(03\)00392-5](https://doi.org/10.1016/s0889-5406(03)00392-5).
43. Alqerban A, Jacobs R, Fieuws S, Nackaerts O, SEDENTEXCT Project Consortium, Willems G. Comparison of 6 cone-beam computed tomography systems for image quality and detection of simulated canine impaction-induced external root resorption in maxillary lateral incisors. *Am J Orthod Dentofacial Orthop*. 2011;140:e129-139. <https://doi.org/10.1016/j.ajodo.2011.03.021>.
44. Karacin G, Şenişik NE, Yildirim D. Comparison of lateral tooth volume and morphology between buccally and palatally localised unilateral

- impacted maxillary canine cases on cone-beam computed tomography. *J Craniofac Surg*. 2021;33:752–6. <https://doi.org/10.1097/SCS.00000000000006936>.
45. Ling K, Ho C, Kravchuk O, Olive R. Comparison of surgical and non-surgical methods of treating palatally impacted canines. II. Aesthetic Outcomes. *Australian Orthodontic J*. 2007;23:8–15. <https://doi.org/10.2478/aoj-2007-0002>.
 46. Khan, R.; Antony, V.; Elhammali, N. Comparison of the Surgical Methods in Combined Surgical –Orthodontic Correction of Impacted Teeth . *IOSR Journal of Dental and Medical Sciences* 2015.
 47. Alqerban A, Jacobs R, Fieuews S, Willems G. Comparison of two cone beam computed tomographic systems versus panoramic imaging for localization of impacted maxillary canines and detection of root resorption. *Eur J Orthod*. 2011;33:93–102. <https://doi.org/10.1093/ejo/cjq034>.
 48. Campbell CM, DiBiase A, Fleming PS. Concomitant dilaceration, transposition, and intraosseous migration: report of a patient treated with maxillary canine-central incisor substitution. *Am J Orthod Dentofacial Orthop*. 2014;146:514–21. <https://doi.org/10.1016/j.ajodo.2013.11.024>.
 49. Peck S, Peck L, Kataja M. Concomitant occurrence of canine malposition and tooth agenesis: evidence of orofacial genetic fields. *Am J Orthod Dentofacial Orthop*. 2002;122:657–60. <https://doi.org/10.1067/mod.2002.129915>.
 50. Lund H, Gröndahl K, Gröndahl H-G. Cone beam computed tomography for assessment of root length and marginal bone level during orthodontic treatment. *Angle Orthod*. 2010;80:466–73. <https://doi.org/10.2319/072909-427.1>.
 51. Becker A, Chaushu S. Etiology of maxillary canine impaction: a review. *Am J Orthod Dentofacial Orthop*. 2015;148:557–67. <https://doi.org/10.1016/j.ajodo.2015.06.013>.
 52. Patano A, Malcangi G, De Santis M, Morolla R, Settanni V, Piras F, Inchingolo AD, Mancini A, Inchingolo F, Dipalma G, et al. Conservative treatment of dental non-carious cervical lesions: a scoping review. *Biomedicines*. 2023;11:1530. <https://doi.org/10.3390/biomedicines11061530>.
 53. Franka R, Smith TG, Dyer JL, Wu X, Niezgodna M, Rupprecht CE. Current and future tools for global canine rabies elimination. *Antiviral Res*. 2013;100:220–5. <https://doi.org/10.1016/j.antiviral.2013.07.004>.
 54. Crivelini MM, de Araújo VC, de Sousa SOM, de Araújo NS. Cytokeratins in epithelia of odontogenic neoplasms. *Oral Dis*. 2003;9:1–6. <https://doi.org/10.1034/j.1601-0825.2003.00861.x>.
 55. Ferati K, Bexheti-Ferati A, Palermo A, Pezzolla C, Trilli I, Sardano R, Latini G, Inchingolo AD, Inchingolo AM, Malcangi G, et al. Diagnosis and orthodontic treatment of obstructive sleep apnea syndrome children-a systematic review. *Diagnostics* (Basel). 2024;14:289. <https://doi.org/10.3390/diagnostics14030289>.
 56. Ratson T. Diagnosis and treatment of supernumerary teeth in the pre-maxillary region: a literature review. *Refuat Hapeh Vehashinayim* (1993). 2013;30(26–30):69.
 57. Chalakkal P, Thomas AM, Chopra S. Displacement, location, and angulation of unerupted permanent maxillary canines and absence of canine bulge in children. *Am J Orthod Dentofacial Orthop*. 2011;139:345–50. <https://doi.org/10.1016/j.ajodo.2009.03.044>.
 58. Ericson S, Kuroi J. Radiographic examination of ectopically erupting maxillary canines. *Am J Orthod Dentofacial Orthop*. 1987;91:483–92. [https://doi.org/10.1016/0889-5406\(87\)90005-9](https://doi.org/10.1016/0889-5406(87)90005-9).
 59. Arangio P, Marianetti TM, Tedaldi M, Ramieri V, Cascone P. Early secondary alveoloplasty in cleft lip and palate. *J Craniofacial Surg*. 2008;19:1364. <https://doi.org/10.1097/SCS.0b013e31818434c3>.
 60. Ericson S, Kuroi J. Early treatment of palatally erupting maxillary canines by extraction of the primary canines. *Eur J Orthod*. 1988;10:283–95. <https://doi.org/10.1093/ejo/10.4.283>.
 61. Sigler L, Baccetti T, McNamara J. Effect of rapid maxillary expansion and transpalatal arch treatment associated with deciduous canine extraction on the eruption of palatally displaced canines: a 2-center prospective study. *American J Orthod Dentofacial Orthop*. 2011;139:e235–44. <https://doi.org/10.1016/j.ajodo.2009.07.015>.
 62. Ngan P, Hornbrook R, Weaver B. Early timely management of ectopically erupting maxillary canines. *Seminars Orthodontics*. 2005;11:152–63. <https://doi.org/10.1053/j.sodo.2005.04.009>.
 63. Başer EN, Akar NK, Sayar G. Effects of ballista and kilroy springs on palatally impacted canines: a finite element model analysis. *Turk J Orthod*. 2023;36:30–8. <https://doi.org/10.4274/TurkJOrthod.2022.2022.10>.
 64. Ucar FI, Celebi AA, Tan E, Topcuoğlu T, Sekerci AE. Effects of impacted maxillary canines on root resorption of lateral incisors : a cone beam computed tomography study. *J Orofac Orthop*. 2017;78:233–40. <https://doi.org/10.1007/s00056-016-0077-6>.
 65. Dehis, H.; Rahman, A.R.A.; Aziz, M.A.W.M.A.; Yassin, M.M.; Yussif, N.M.A. Efficacy and Safety of Locally Injectable Vitamin C on Accelerating the Orthodontic Movement of Maxillary Canine Impaction (Oral Mesotherapy Technique): Prospective Study. *Clinical Cases in Mineral and Bone Metabolism* 2018, 15, 280–287, <https://doi.org/10.11138/ccmbm/2018.15.2.280>.
 66. Meslin F-X, Briggs DJ. Eliminating canine rabies, the principal source of human infection: what will it take? *Antiviral Res*. 2013;98:291–6. <https://doi.org/10.1016/j.antiviral.2013.03.011>.
 67. Yan B, Sun Z, Fields H, Wang L, Luo L. Etiologic factors for buccal and palatal maxillary canine impaction: a perspective based on cone-beam computed tomography analyses. *Am J Orthod Dentofacial Orthop*. 2013;143:527–34. <https://doi.org/10.1016/j.ajodo.2012.11.021>.
 68. Becker A, Chaushu S. Long-term follow-up of severely resorbed maxillary incisors after resolution of an etiologically associated impacted canine. *Am J Orthod Dentofacial Orthop*. 2005;127:650–4. <https://doi.org/10.1016/j.ajodo.2004.03.031>. (quiz 754).
 69. Minetti E, Palermo A, Savadori P, Barlattani A, Franco R, Michele M, Gianfreda F, Bollero P. Autologous tooth graft: a histological comparison between dentin mixed with xenograft and dentin alone grafts in socket preservation. *J Biol Regul Homeost Agents*. 2019;33:189–97.
 70. Franco R, Barlattani A Jr, Perrone MA, Basili M, Miranda M, Costacurta M, Gualtieri P, Pujia A, Merra G, Bollero P. Obesity, bariatric surgery and periodontal disease: a literature update. *Eur Rev Med Pharmacol Sci*. 2020;24:5036–45. https://doi.org/10.26355/eurev_202005_21196.
 71. Bollero P. Oral health and implant therapy in parkinson's patients: review. *Oral Implantol*. 2017;10:105–105. <https://doi.org/10.11138/orl/2017.10.2.105>.
 72. Tadinada A, Mahdian M, Vishwanath M, Allareddy V, Upadhyay M, Yadav S. Evaluation of alveolar bone dimensions in unilateral palatally impacted canine: a cone-beam computed tomographic analyses. *Eur J Orthod*. 2015;37:596–602. <https://doi.org/10.1093/ejo/cju098>.
 73. Alsino HI, Hajeer MY, Alkhouri I, Murad RMT, Khattab TZ. Evaluation of the levels of pain, discomfort, functional impairments and satisfaction with the Periodontally Accelerated Osteogenic Orthodontics (PAOO) when leveling and aligning crowded teeth: a prospective cohort study. *Cureus*. 2022;14:e22623. <https://doi.org/10.7759/cureus.22623>.
 74. Palaia G, Impellizzeri A, Tenore G, Caporali F, Visca P, Del Vecchio A, Galluccio G, Polimeni A, Romeo U. Ex vivo histological analysis of the thermal effects created by a 445-nm diode laser in oral soft tissue biopsy. *Clin Oral Investig*. 2020;24:2645–52. <https://doi.org/10.1007/s00784-019-03123-4>.
 75. Recuenco S, Eidson M, Cherry B, Kulldorff M, Johnson G. Factors associated with endemic raccoon (*Procyon Lotor*) rabies in terrestrial mammals in New York State, USA. *Prev Vet Med*. 2008;86:30–42. <https://doi.org/10.1016/j.prevetmed.2008.03.001>.
 76. Naoumova J, Alfaro GE, Peck S. Space conditions, palatal vault height, and tooth size in patients with and without palatally displaced canines: a prospective cohort study. *Angle Orthod*. 2018;88:726–32. <https://doi.org/10.2319/120717-843.1>.
 77. Zuccati G, Ghobadlu J, Nieri M, Clauser C. Factors associated with the duration of forced eruption of impacted maxillary canines: a retrospective study. *Am J Orthod Dentofacial Orthop*. 2006;130:349–56. <https://doi.org/10.1016/j.ajodo.2004.12.028>.
 78. Ozawa T, Omura S, Fukuyama E, Matsui Y, Torikai K, Fujita K. Factors influencing secondary alveolar bone grafting in cleft lip and palate patients: prospective analysis using CT image analyzer. *Cleft Palate Craniofac J*. 2007;44:286–91. <https://doi.org/10.1597/06-054>.
 79. Stewart JA, Heo G, Glover KE, Williamson PC, Lam EW, Major PW. Factors that relate to treatment duration for patients with palatally impacted maxillary canines. *Am J Orthod Dentofacial Orthop*. 2001;119:216–25. <https://doi.org/10.1067/mod.2001.110989>.

80. Bhutta ZA, Sommerfeld J, Lassi ZS, Salam RA, Das JK. Global burden, distribution, and interventions for infectious diseases of poverty. *Infect Dis Poverty*. 2014;3:21. <https://doi.org/10.1186/2049-9957-3-21>.
81. Murray CJL, Ortblad KF, Guinovart C, Lim SS, Wolock TM, Roberts DA, Dansereau EA, Graetz N, Barber RM, Brown JC, et al. Global, regional, and national incidence and mortality for HIV, tuberculosis, and malaria during 1990–2013: a systematic analysis for the global burden of disease study 2013. *Lancet*. 2014;384:1005–70. [https://doi.org/10.1016/S0140-6736\(14\)60844-8](https://doi.org/10.1016/S0140-6736(14)60844-8).
82. Björk A, Skieller V. Growth of the maxilla in three dimensions as revealed radiographically by the implant method. *Br J Orthod*. 1977;4:53–64. <https://doi.org/10.1179/bjo.4.2.53>.
83. Counihan K, Al-Awadhi EA, Butler J. Guidelines for the assessment of the impacted Maxillary Canine. *Dent Update*. 2013;40(770–772):775–7. <https://doi.org/10.12968/denu.2013.40.9.770>.
84. Ferguson EA, Hampson K, Cleaveland S, Consunji R, Deray R, Friar J, Haydon DT, Jimenez J, Pancipane M, Townsend SE. Heterogeneity in the spread and control of infectious disease: consequences for the elimination of canine rabies. *Sci Rep*. 2015;5:18232. <https://doi.org/10.1038/srep18232>.
85. Yoon RK, Chussid S, Davis MJ. Impacted maxillary anterior supernumerary teeth: a survey of forty-two cases. *N Y State Dent J*. 2008;74:24–7.
86. Guarneri R, Bertoldo S, Cassetta M, Altieri F, Grenga C, Vichi M, Di Giorgio R, Barbato E. Periodontal results of different therapeutic approaches (Open vs. Closed Technique) and timing evaluation (< 2 Year vs. > 2 Year) of palatal impacted canines: a systematic review. *BMC Oral Health*. 2021;21:574. <https://doi.org/10.1186/s12903-021-01937-x>.
87. Bishara SE. Impacted maxillary canines: a review. *Am J Orthod Dentofacial Orthop*. 1992;101:159–71. [https://doi.org/10.1016/0889-5406\(92\)70008-X](https://doi.org/10.1016/0889-5406(92)70008-X).
88. Raubenheimer EJ, van Heerden WF, Noffke CE. Infrequent clinicopathological findings in 108 ameloblastomas. *J Oral Pathol Med*. 1995;24:227–32. <https://doi.org/10.1111/j.1600-0714.1995.tb01172.x>.
89. Baccetti T, Mucedero M, Leonardi M, Cozza P. Interceptive treatment of palatal impaction of maxillary canines with rapid maxillary expansion: a randomized clinical trial. *Am J Orthod Dentofacial Orthop*. 2009;136:657–61. <https://doi.org/10.1016/j.ajodo.2008.03.019>.
90. Thilander B, Jakobsson SO. Local factors in impaction of maxillary canines. *Acta Odontol Scand*. 1968;26:145–68. <https://doi.org/10.3109/00016356809004587>.
91. Lai CS, Suter V, Katsaros C, Bornstein MM. Localization of impacted maxillary canines and root resorption of neighbouring teeth: a study assessing the diagnostic value of panoramic radiographs in two groups of observers. *Eur J Orthod*. 2014;36:450–6. <https://doi.org/10.1093/ejo/cjt074>.
92. Rossini G, Cavallini C, Cassetta M, Galluccio G, Barbato E. Localization of impacted maxillary canines using cone beam computed tomography. *Rev Literature Ann Stomatol (Roma)*. 2012;3:14–8.
93. Nagpal A, Pai KM, Setty S, Sharma G. Localization of impacted maxillary canines using panoramic radiography. *J Oral Sci*. 2009;51:37–45. <https://doi.org/10.2334/josnusd.51.37>.
94. Levander E, Malmgren O. Long-term follow-up of maxillary incisors with severe apical root resorption. *Eur J Orthod*. 2000;22:85–92. <https://doi.org/10.1093/ejo/22.1.85>.
95. Becker A, Chaushu S, Casap-Caspi N. Cone-beam computed tomography and the orthosurgical management of impacted teeth. *J Am Dent Assoc*. 2010;141(Suppl 3):145–S18. <https://doi.org/10.14219/jada.archive.2010.0360>.
96. Caprioglio A, Vanni A, Bolamperti L. Long-term periodontal response to orthodontic treatment of palatally impacted maxillary canines. *Eur J Orthod*. 2013;35:323–8. <https://doi.org/10.1093/ejo/cjs020>.
97. Chen S, Wang L, Li G, Wu TH, Diachina S, Tejera B, Kwon JJ, Lin FC, Lee YT, Xu T, et al. Machine learning in orthodontics: introducing a 3d auto-segmentation and auto-landmark finder of CBCT images to assess maxillary constriction in unilateral impacted canine patients. *Angle Orthod*. 2020;90:77–84. <https://doi.org/10.2319/012919-59.1>.
98. Pearson MH, Robinson SN, Reed R, Birnie DJ, Zaki GA. Management of palatally impacted canines: the findings of a collaborative study. *Eur J Orthod*. 1997;19:511–5. <https://doi.org/10.1093/ejo/19.5.511>.
99. Inchingolo F, Inchingolo AD, Palumbo I, Guglielmo M, Balestriere L, Casamassima L, Ciccarese D, Marotti P, Mancini A, Palermo A, et al. Management of physiological gingival melanosis by diode laser depigmentation versus surgical scalpel: a systematic review. *Dentistry Rev*. 2024;4:100146. <https://doi.org/10.1016/j.dentre.2024.100146>.
100. Patano A, Malcangi G, Inchingolo AD, Garofoli G, De Leonardis N, Azzolini D, Latini G, Mancini A, Carpentiere V, Laudadio C, et al. Mandibular crowding: diagnosis and management—a scoping review. *J Pers Med*. 2023;13:774. <https://doi.org/10.3390/jpm13050774>.
101. Buschang PH, Carrillo R, Liu SS, Demirjian A. Maxillary and mandibular dentoalveolar heights of French-canadians 10 to 15 years of age. *Angle Orthod*. 2008;78:70–6. <https://doi.org/10.2319/092006-381.1>.
102. Stahl F, Grabowski R. Maxillary canine displacement and genetically determined predisposition to disturbed development of the dentition. *J Orofac Orthop*. 2003;64:167–77. <https://doi.org/10.1007/s00056-003-0221-y>.
103. Yan B, Sun Z, Fields H, Wang L. Maxillary canine impaction increases root resorption risk of adjacent teeth: a problem of physical proximity. *Am J Orthod Dentofacial Orthop*. 2012;142:750–7. <https://doi.org/10.1016/j.ajodo.2012.07.016>.
104. Bayram M, Ozer M, Sener I. Maxillary canine impactions related to impacted central incisors: two case reports. *J Contemp Dent Pract*. 2007;8:72–81.
105. Rimes RJ, Mitchell CN, Willmot DR. Maxillary incisor root resorption in relation to the ectopic canine: a review of 26 patients. *Eur J Orthod*. 1997;19:79–84. <https://doi.org/10.1093/ejo/19.1.79>.
106. Bjerklín K, Guitirok CH. Maxillary incisor root resorption induced by ectopic canines. *Angle Orthod*. 2011;81:800–6. <https://doi.org/10.2319/011311-23.1>.
107. Flores-Mir C, Rosenblatt MR, Major PW, Carey JP, Heo G. Measurement accuracy and reliability of tooth length on conventional and CBCT reconstructed panoramic radiographs. *Dental Press J Orthod*. 2014;19:45–53. <https://doi.org/10.1590/2176-9451.19.5.045-053.oar>.
108. Gorlin RJ, Goltz RW. Multiple nevoid basal-cell epithelioma, jaw cysts and bifid rib. A Syndrome. *N Engl J Med*. 1960;262:908–12. <https://doi.org/10.1056/NEJM196005052621803>.
109. Amlashi SFA, Riffaud L, Brassier G, Morandi X. Nevoid basal cell carcinoma syndrome: relation with desmoplastic medulloblastoma in infancy. A population-based study and review of the literature. *Cancer*. 2003;98:618–24. <https://doi.org/10.1002/cncr.11537>.
110. Parkin NA, Almutairi S, Benson PE. Surgical exposure and orthodontic alignment of palatally displaced canines: can we shorten treatment time? *J Orthod*. 2019;46:54–9. <https://doi.org/10.1177/1465312519841384>.
111. Lindauer SJ, Isaacson RJ. One-couple orthodontic appliance systems. *Semin Orthod*. 1995;1:12–24. [https://doi.org/10.1016/s1073-8746\(95\)80084-0](https://doi.org/10.1016/s1073-8746(95)80084-0).
112. Parkin, N.; Benson, P.E.; Thind, B.; Shah, A. Open versus Closed Surgical Exposure of Canine Teeth That Are Displaced in the Roof of the Mouth. *Cochrane Database Syst Rev* 2008, CD006966, <https://doi.org/10.1002/14651858.CD006966.pub2>.
113. Parkin NA, Deery C, Smith A-M, Tinsley D, Sandler J, Benson PE. No difference in surgical outcomes between open and closed exposure of palatally displaced maxillary canines. *J Oral Maxillofac Surg*. 2012;70:2026–34. <https://doi.org/10.1016/j.joms.2012.02.028>.
114. Fournier A, Turcotte J-Y, Bernard C. Orthodontic considerations in the treatment of maxillary impacted canines. *Am J Orthod*. 1982;81:236–9. [https://doi.org/10.1016/0002-9416\(82\)90056-2](https://doi.org/10.1016/0002-9416(82)90056-2).
115. Olive RJ. Orthodontic treatment of palatally impacted maxillary canines. *Aust Orthod J*. 2002;18:64–70.
116. Mancini A, Chirico F, Inchingolo AM, Piras F, Colonna V, Marotti P, Carone C, Inchingolo AD, Inchingolo F, Dipalma G. Osteonecrosis of the jaws associated with herpes zoster infection: a systematic review and a rare case report. *Microorganisms*. 2024;12:1506. <https://doi.org/10.3390/microorganisms12081506>.
117. Sair M, Rebellato J, Sheats RD. Palatal displacement of canines and maxillary skeletal width. *Am J Orthod Dentofacial Orthop*. 2006;129:511–9. <https://doi.org/10.1016/j.ajodo.2005.03.021>.
118. Bazargani F, Magnuson A, Dolati A, Lennartsson B. Palatally displaced maxillary canines: factors influencing duration and cost of treatment. *Eur J Orthod*. 2013;35:310–6. <https://doi.org/10.1093/ejo/cjr143>.
119. Smaliene D, Kavaliauskiene A, Pacauskiene I, Zasciurinskiene E, Bjerklín K. Palatally impacted maxillary canines: choice of surgical-orthodontic

- treatment method does not influence post-treatment periodontal status. A controlled prospective study. *Eur J Orthod*. 2013;35:803–10. <https://doi.org/10.1093/ejo/cjs102>.
120. Inchingolo F, Inchingolo AM, Latini G, Ferrante L, Trilli I, Del Vecchio G, Palmieri G, Malcangi G, Inchingolo AD, Dipalma G. Oxidative stress and natural products in orthodontic treatment: a systematic review. *Nutrients*. 2023;16:113. <https://doi.org/10.3390/nu16010113>.
 121. Chausshu S, Becker A, Zeltser R, Branski S, Vasker N, Chausshu G. Patients' perception of recovery after exposure of impacted teeth: a comparison of closed- versus open-eruption techniques. *J Oral Maxillofac Surg*. 2005;63:323–9. <https://doi.org/10.1016/j.joms.2004.11.007>.
 122. de Sousa ET, da Silva BF, Maia FBM, Forte FDS, Sampaio FC. Perception of children and mothers regarding dental aesthetics and orthodontic treatment need: a cross-sectional study. *Prog Orthod*. 2016;17:37. <https://doi.org/10.1186/s40510-016-0149-6>.
 123. Hansson C, Rindler A. Periodontal conditions following surgical and orthodontic treatment of palatally impacted maxillary canines—a follow-up study. *Angle Orthod*. 1998;68:167–72. [https://doi.org/10.1043/0003-3219\(1998\)068%3c0167:PCFSAO%3e2.3.CO;2](https://doi.org/10.1043/0003-3219(1998)068%3c0167:PCFSAO%3e2.3.CO;2).
 124. Grenga C, Guarnieri R, Grenga V, Bovi M, Bertoldo S, Galluccio G, Di Giorgio R, Barbato E. Periodontal evaluation of palatally impacted maxillary canines treated by closed approach with ultrasonic surgery and orthodontic treatment: a retrospective pilot study. *Sci Rep*. 2021;11:2843. <https://doi.org/10.1038/s41598-021-82510-y>.
 125. Alpaslan Yayli NZ, Kaya Y, CinarsoyCigerim S. Periodontal health of unilateral labially vs. Palatally impacted maxillary canines erupted by closed eruption technique. *J Orofac Orthop*. 2023;84:133–42. <https://doi.org/10.1007/s00056-022-0413-7>.
 126. Parkin NA, Freeman JV, Deery C, Benson PE. Esthetic judgments of palatally displaced canines 3 months postdebond after surgical exposure with either a closed or an open technique. *Am J Orthod Dentofacial Orthop*. 2015;147:173–81. <https://doi.org/10.1016/j.jajodo.2014.10.026>.
 127. Guarnieri R, Cavallini C, Vernucci R, Vichi M, Leonardi R, Barbato E. Impacted maxillary canines and root resorption of adjacent teeth: a retrospective observational study. *Med Oral Patol Oral Cir Bucal*. 2016;21:e743–50. <https://doi.org/10.4317/medoral.21337>.
 128. Mummolo S, Nota A, De Felice ME, Marcattili D, Tecco S, Marzo G. Periodontal status of buccally and palatally impacted maxillary canines after surgical-orthodontic treatment with open technique. *J Oral Sci*. 2018;60:552–6. <https://doi.org/10.2334/josnusd.17-0394>.
 129. Evren AD, Nevzatoglu Ş, Arun T, Acar A. Periodontal status of ectopic canines after orthodontic treatment. *Angle Orthod*. 2014;84:18–23. <https://doi.org/10.2319/041513-290.1>.
 130. Schmidt AD, Kokich VG. Periodontal response to early uncovering, autonomous eruption, and orthodontic alignment of palatally impacted maxillary canines. *Am J Orthod Dentofacial Orthop*. 2007;131:449–55. <https://doi.org/10.1016/j.jajodo.2006.04.028>.
 131. Calil, L.R.; Janson, G.; Silva, V.M. da; Freitas, M.R. de; Almeida, A.L.P.F. de; Garib, D. Periodontal status of maxillary central incisors after orthodontic traction: a longitudinal follow-up. *J Appl Oral Sci* 2022, 30, e20210492, <https://doi.org/10.1590/1678-7757-2021-0492>.
 132. Gharaibeh TM, Al-Nimri KS. Postoperative pain after surgical exposure of palatally impacted canines: closed-eruption versus open-eruption, a prospective randomized study. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod*. 2008;106:339–42. <https://doi.org/10.1016/j.tripleo.2007.12.025>.
 133. Malcangi G, Patano A, Guglielmo M, Sardano R, Palmieri G, Di Pede C, de Ruvo E, Inchingolo AD, Mancini A, Inchingolo F, et al. Precision medicine in oral health and diseases: a systematic review. *J Pers Med*. 2023;13:725. <https://doi.org/10.3390/jpm13050725>.
 134. Inchingolo AM, Inchingolo AD, Carpentiere V, Del Vecchio G, Ferrante L, Di Noia A, Palermo A, Di Venere D, Dipalma G, Inchingolo F. Predictability of dental distalization with clear aligners: a systematic review. *Bioengineering (Basel)*. 2023;10:1390. <https://doi.org/10.3390/bioengineering10121390>.
 135. Inchingolo AM, Malcangi G, Piras F, Palmieri G, Settanni V, Riccaldo L, Morolla R, Buongiorno S, de Ruvo E, Inchingolo AD, et al. Precision medicine on the effects of microbiota on head-neck diseases and biomarkers diagnosis. *J Pers Med*. 2023;13:933. <https://doi.org/10.3390/jpm13060933>.
 136. Rafflenbeul F, Gros C-I, Lefebvre F, Bahi-Gross S, Maizeray R, Bolender Y. Prevalence and risk factors of root resorption of adjacent teeth in maxillary canine impaction, among untreated children and adolescents. *Eur J Orthod*. 2019;41:447–53. <https://doi.org/10.1093/ejo/cjy078>.
 137. Lupi JE, Handelman CS, Sadowsky C. Prevalence and severity of apical root resorption and alveolar bone loss in orthodontically treated adults. *Am J Orthod Dentofacial Orthop*. 1996;109:28–37. [https://doi.org/10.1016/s0889-5406\(96\)70160-9](https://doi.org/10.1016/s0889-5406(96)70160-9).
 138. Alamri A, Alqanas S, Aljar Y, Alqahtani F, Ahmed S. Prevalence of canine impaction in different cities of Saudi Arabia: a systematic review. *Saudi Dent J*. 2024;36:688–97. <https://doi.org/10.1016/j.sdentj.2024.02.018>.
 139. Ballini A, Santacroce L, Cantore S, Bottalico L, Dipalma G, Topi S, Saini R, De Vito D, Inchingolo F. Probiotics efficacy on oxidative stress values in inflammatory bowel disease: a randomized double-blinded placebo-controlled pilot study. *Endocr Metab Immune Disord Drug Targets*. 2019;19:373–81. <https://doi.org/10.2174/1871530319666181221150352>.
 140. Inchingolo F, Ferrara I, Viapiano F, Ciocia AM, Palumbo I, Guglielmo M, Inchingolo AD, Palermo A, Bordea IR, Inchingolo AM, et al. Primary failure eruption: genetic investigation, diagnosis and treatment: a systematic review. *Children (Basel)*. 2023;10:1781. <https://doi.org/10.3390/bambini10111781>.
 141. Woloshyn H, Artun J, Kennedy DB, Joondeph DR. Pulpal and periodontal reactions to orthodontic alignment of palatally impacted canines. *Angle Orthod*. 1994;64:257–64. [https://doi.org/10.1043/0003-3219\(1994\)064%3c0257:PAPRTO%3e2.0.CO;2](https://doi.org/10.1043/0003-3219(1994)064%3c0257:PAPRTO%3e2.0.CO;2).
 142. Minervini G, Franco R, Marrapodi MM, Fiorillo L, Cervino G, Cicciù M. The association between parent education level, oral health, and oral-related sleep disturbance. An observational cross-sectional study. *Eur J Paediatr Dentist*. 2023;24:218–23. <https://doi.org/10.23804/ejpd.2023.1910>.
 143. Minervini G, Franco R, Marrapodi MM, Almeida LE, Ronsivalve V, Cicciù M. Prevalence of Temporomandibular Disorders (TMD) in obesity patients: a systematic review and meta-analysis. *J Oral Rehabil*. 2023;50:1544–53. <https://doi.org/10.1111/joor.13573>.
 144. Minervini G, Franco R, Marrapodi MM, Fiorillo L, Cervino G, Cicciù M. Post-traumatic stress, prevalence of temporomandibular disorders in war veterans: systematic review with meta-analysis. *J Oral Rehabil*. 2023;50:1101–9. <https://doi.org/10.1111/joor.13535>.
 145. Minervini G, Franco R, Marrapodi MM, Di Blasio M, Isola G, Cicciù M. Conservative treatment of temporomandibular joint condylar fractures: a systematic review conducted according to PRISMA guidelines and the cochrane handbook for systematic reviews of interventions. *J Oral Rehabil*. 2023;50:886–93. <https://doi.org/10.1111/joor.13497>.
 146. Minervini G, Franco R, Marrapodi MM, Di Blasio M, Ronsivalve V, Cicciù M. Children oral health and parents education status: a cross-sectional study. *BMC Oral Health*. 2023;23:787. <https://doi.org/10.1186/s12903-023-03424-x>.
 147. Alshawy ES, Kolkordi SH. Revealing the effect of impacted canines on the adjacent teeth. A three dimensional study using CBCT. *J Pharm Bioallied Sci*. 2023;15:5720–4. https://doi.org/10.4103/jpbs.jpbs_173_23.
 148. Rodríguez-Cárdenas YA, Arriola-Guillén LE, Ruiz-Mora GA, Aliaga-Del Castillo A, Boessio-Vizzotto M, Dias-Da Silveira HL. Root changes in buccal versus palatal maxillary impacted canines of adults: a longitudinal and retrospective 3-dimensional study before and after orthodontic traction. *Int Orthod*. 2020;18:490–502. <https://doi.org/10.1016/j.ortho.2020.05.007>.
 149. Ericson S, Kuroi J. Radiographic Assessment of maxillary canine eruption in children with clinical signs of eruption disturbance. *Eur J Orthod*. 1986;8:133–40. <https://doi.org/10.1093/ejo/8.3.133>.
 150. Ericson S, Kuroi J. Resorption of maxillary lateral incisors caused by ectopic eruption of the canines. A clinical and radiographic analysis of predisposing factors. *Am J Orthod Dentofacial Orthop*. 1988;94:503–13. [https://doi.org/10.1016/0889-5406\(88\)90008-x](https://doi.org/10.1016/0889-5406(88)90008-x).
 151. Ericson S, Bjerklín K, Falahat B. Does the canine dental follicle cause resorption of permanent incisor roots? A computed tomographic study of erupting maxillary canines. *Angle Orthod*. 2002;72:95–104. [https://doi.org/10.1043/0003-3219\(2002\)072%3c0095:DTCDFC%3e2.0.CO;2](https://doi.org/10.1043/0003-3219(2002)072%3c0095:DTCDFC%3e2.0.CO;2).
 152. Cao D, Shao B, Izadikhah I, Xie L, Wu B, Li H, Yan B. Root dilaceration in maxillary impacted canines and adjacent teeth: a retrospective analysis of the difference between buccal and palatal impaction. *Am J Orthod*

- Dentofacial Orthop. 2021;159:167–74. <https://doi.org/10.1016/j.jajodo.2019.12.019>.
153. da Silva, A.C.; Capistrano, A.; de Almeida-Pedrin, R.R.; Cardoso, M. de A.; Conti, A.C. de C.F.; Capelozza, L. Root length and alveolar bone level of impacted canines and adjacent teeth after orthodontic traction: a long-term evaluation. *J Appl Oral Sci* 2017, 25, 75–81. <https://doi.org/10.1590/1678-77522016-0133>.
 154. Melchor-Soto ME, Arriola-Guillén LE, Aliaga-Del Castillo A, Ruiz-Mora GA, Rodríguez-Cárdenas YA. Root morphology of lateral incisors adjacent to impacted maxillary canines: a cone-beam computed tomography retrospective cross-sectional study. *Int Orthod*. 2022;20:100692. <https://doi.org/10.1016/j.ortho.2022.100692>.
 155. Cernochova P, Krupa P, Izakovicova-Holla L. Root Resorption associated with ectopically erupting maxillary permanent canines: a computed tomography study. *Eur J Orthod*. 2011;33:483–91. <https://doi.org/10.1093/ejo/cjq085>.
 156. Cuellar Y, Andrés Velásquez S, Domínguez A. Root Regeneration with photobiomodulation of an upper lateral incisor associated with root resorption due to an impacted maxillary canine: a case report. *Photobiomodul Photomed Laser Surg*. 2024;42:422–7. <https://doi.org/10.1089/photob.2023.0150>.
 157. Cumbo E, Messina P, Gallina G, Scardina GA. An experimental method to add new prosthetic teeth in the removable partial denture framework: TiG cold welding and preformed pins. *Prosthesis*. 2023;5:1120–8. <https://doi.org/10.3390/protesi5040078>.
 158. Karakas-Stupar I, Zaugg LK, Zitzmann NU, Joda T, Wolfart S, Tuna T. Clinical protocol for implant-assisted partial removable dental prostheses in Kennedy class I: a case report. *Prosthesis*. 2023;5:1002–10. <https://doi.org/10.3390/protesi5040069>.
 159. Alkahtany M, Beatty MW, Alsalleeh F, Petro TM, Simetich B, Zhou Y, Feely D, Polyzois G. Color stability, physical properties and antifungal effects of ZrO2 additions to experimental maxillofacial silicones: comparisons with TiO2. *Prosthesis*. 2023;5:916–38. <https://doi.org/10.3390/protesi5030064>.
 160. Pera F, Carossa M, Bagnasco F, Crupi A, Ambrogio G, Isola G, Menini M, Pesce P. Comparison between bone-level and tissue-level implants in immediate-loading full-arch rehabilitations: a retrospective multi-center 1-year follow-up study. *Prosthesis*. 2023;5:1301–11. <https://doi.org/10.3390/protesi5040089>.
 161. Tallarico M, Cuccu M, Meloni SM, Lumbau AI, Baldoni E, Pisano M, Fiorillo L, Cervino G. Digital analysis of a novel impression method named the biological-oriented digital impression technique: a clinical audit. *Prosthesis*. 2023;5:992–1001. <https://doi.org/10.3390/protesi5040068>.
 162. Lombardo G, Signoriello A, Marincola M, Bonfante EA, Díaz-Caballero A, Tomizioli N, Pardo A, Zangani A. Five-year follow-up of 8 and 6 mm locking-taper implants treated with a reconstructive surgical protocol for peri-implantitis: a retrospective evaluation. *Prosthesis*. 2023;5:1322–42. <https://doi.org/10.3390/protesi5040091>.
 163. Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, Shamseer L, Tetzlaff JM, Akl EA, Brennan SE, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *PLoS Med*. 2021;18:e1003583. <https://doi.org/10.1371/journal.pmed.1003583>.
 164. Chaushu S, Chaushu G, Becker A. The use of panoramic radiographs to localize displaced maxillary canines. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod*. 1999;88:511–6. [https://doi.org/10.1016/s1079-2104\(99\)70072-7](https://doi.org/10.1016/s1079-2104(99)70072-7).
 165. Crescini A, Nieri M, Buti J, Baccetti T, Pini Prato GP. Orthodontic and periodontal outcomes of treated impacted maxillary canines. *Angle Orthod*. 2007;77:571–7. <https://doi.org/10.2319/080406-318.1>.
 166. Kivovics M, Szanyi SM, Takács A, Répási M, Németh O, Mijiritsky E. Computer-assisted open exposure of palatally impacted canines for orthodontic eruption: a randomized clinical trial. *J Dent*. 2024;147:105110. <https://doi.org/10.1016/j.jdent.2024.105110>.
 167. Björksved M, Ryen L, Lindsten R, Bazargani F. Open and closed surgical exposure of palatally displaced canines: a cost-minimization analysis of a multicentre, randomized controlled trial. *Eur J Orthod*. 2021;43:498–505. <https://doi.org/10.1093/ejo/cjab052>.
 168. Impellizzeri A, Horodynski M, Palaia G, Monaca G, Pergolini D, Polimeni A, Romeo U, Galluccio G. CO2 and diode lasers vs. conventional surgery in the disinclusion of palatally impacted canines: a randomized controlled trial. *Photonics*. 2023;10:244. <https://doi.org/10.3390/fotonica10030244>.
 169. Mousa MR, Hajeer MY, Burhan AS, Heshmeh O, Alam MK. The Effectiveness of minimally-invasive corticotomy-assisted orthodontic treatment of palatally impacted canines compared to the traditional traction method in terms of treatment duration, velocity of traction movement and the associated dentoalveolar changes: a randomized controlled trial. *F1000Res*. 2023;12:699. <https://doi.org/10.12688/f1000research.135338.1>.
 170. Verma S, Singh SP, Verma RK, Kumar V, Singh S, Bhupali NR. Success rate, treatment duration, and pain perception in the management of palatally impacted canines using the k9 and ballista spring: a randomized clinical trial. *Angle Orthod*. 2023;93:33–40. <https://doi.org/10.2319/042122-304.1>.
 171. Björksved M, Arnrup K, Lindsten R, Magnusson A, Sundell AL, Gustafsson A, Bazargani F. Closed vs open surgical exposure of palatally displaced canines: surgery time, postoperative complications, and patients' perceptions: a multicentre, randomized, controlled trial. *Eur J Orthod*. 2018;40:626–35. <https://doi.org/10.1093/ejo/cjy070>.
 172. Cunha-Cruz J. Practicing evidence-based orthodontics: how to critically appraise a randomized controlled trial. *Dental Press J Orthod*. 2015;20:12–5. <https://doi.org/10.1590/2176-9451.20.2.012-015.ebo>.
 173. Jang H, Choi Y, Kwon E, Choi N, Jang Y, Song J, Shin S. Optimal application of compressive palatal stents following mesiodens removal in pediatric patients: a randomized controlled trial. *Med Oral Patol Oral Cir Bucal*. 2021;26:e808–14. <https://doi.org/10.4317/medoral.24802>.
 174. Hadler-Olsen S, Sjögren A, Steinnes J, Dubland M, Bolstad NL, Pirttiniemi P, Kerosuo H, Lähdesmäki R. Double vs single primary tooth extraction in interceptive treatment of palatally displaced canines. *Angle Orthod*. 2020;90:751–7. <https://doi.org/10.2319/031920-196.1>.
 175. Parkin NA, Milner RS, Deery C, Tinsley D, Smith A-M, Germain P, Freeman JV, Bell SJ, Benson PE. Periodontal health of palatally displaced canines treated with open or closed surgical technique: a multicentre, randomized controlled trial. *Am J Orthod Dentofacial Orthop*. 2013;144:176–84. <https://doi.org/10.1016/j.jajodo.2013.03.016>.
 176. Bandeca MC, Borges AH, Aranha AM, Antunes CL, De Musis CR, Vieira EM. Eruption of impacted teeth after alveolar bone graft in cleft lip and palate region. *J Contemp Dent Pract*. 2018;19:933–6. <https://doi.org/10.5005/jp-journals-10024-2360>.
 177. Naoumova J, Kürol J, Kjellberg H. Extraction of the deciduous canine as an interceptive treatment in children with palatally displaced canines - Part II: possible predictors of success and cut-off points for a spontaneous eruption. *Eur J Orthod*. 2015;37:219–29. <https://doi.org/10.1093/ejo/cju102>.
 178. Mousa MR, Hajeer MY, Burhan AS, Heshmeh O. The Effectiveness of conventional and accelerated methods of orthodontic traction and alignment of palatally impacted canines in terms of treatment time, velocity of tooth movement, periodontal, and patient-reported outcomes: a systematic review. *Cureus*. 2022;14:e24888. <https://doi.org/10.7759/cureus.24888>.
 179. Björksved M, Arnrup K, Bazargani SM, Lund H, Magnusson A, Magnusson A, Lindsten R, Bazargani F. Open vs closed surgical exposure of palatally displaced canines: a comparison of clinical and patient-reported outcomes—a multicentre, randomized controlled. *Trial Eur J Orthod*. 2021;43:487–97. <https://doi.org/10.1093/ejo/cjab015>.
 180. Alassiry AM, Hakami Z. Root resorption of adjacent teeth associated with maxillary canine impaction in the Saudi Arabian population: a cross-sectional cone-beam computed tomography study. *Appl Sci*. 2022;12:334. <https://doi.org/10.3390/app12010334>.
 181. Arriola-Guillén LE, Ruiz-Mora GA, Rodríguez-Cárdenas YA, Aliaga-Del Castillo A, Dias-Da Silveira HL. Root resorption of maxillary incisors after traction of unilateral vs bilateral impacted canines with reinforced anchorage. *Am J Orthod Dentofacial Orthop*. 2018;154:645–56. <https://doi.org/10.1016/j.jajodo.2018.01.015>.
 182. Ali IH, Al-Turaihi BA, Mohammed LK, Alam MK. Root resorption of teeth adjacent to untreated impacted maxillary canines: a CBCT study. *Biomed Res Int*. 2021;2021:6635575. <https://doi.org/10.1155/2021/6635575>.
 183. Goldoni R, Dolci C, Boccalari E, Inchingolo F, Paghi A, Strambini L, Galimberti D, Tartaglia GM. Salivary biomarkers of neurodegenerative and

- demyelinating diseases and biosensors for their detection. *Ageing Res Rev.* 2022;76:101587. <https://doi.org/10.1016/j.arr.2022.101587>.
184. Brézulier D, Carnet S, Marie-Cousin A, Sixou J-L. Screening for the risk of canine impaction, what are the presumptive signs and how does it affect orthodontics? A cross-sectional study in France. *PLoS ONE.* 2023;18:e0296395. <https://doi.org/10.1371/journal.pone.0296395>.
 185. Forbes SL, Samson C, Watson CJ. Seasonal impact of scavenger guilds as taphonomic agents in central and Northern Ontario. *Canada J Forensic Sci.* 2022;67:2203–17. <https://doi.org/10.1111/1556-4029.15122>.
 186. da Silva Filho OG, Teles SG, Ozawa TO, Filho LC. Secondary bone graft and eruption of the permanent canine in patients with alveolar clefts: literature review and case report. *Angle Orthod.* 2000;70:174–8. [https://doi.org/10.1043/0003-3219\(2000\)070%3c0174:SBGAE0%3e2.0.CO;2](https://doi.org/10.1043/0003-3219(2000)070%3c0174:SBGAE0%3e2.0.CO;2).
 187. Lima A, Costa P, Barbosa N, Alemida-Pedrin RR, Paranhos L, Cardoso M. Segmented mechanics for traction of impacted maxillary canine: case report with a 3-year follow-up. *Biosci J.* 2019;35:333–46. <https://doi.org/10.14393/BJ-v35n1a2019-42855>.
 188. Hadler-Olsen S, Pirttiniemi P, Kerosuo H, HolstadLimchaichana N, Pesonen P, Kallio-Pulkkinen S, Lähdesmäki R. Root resorptions related to ectopic and normal eruption of maxillary canine teeth - A 3D study. *Acta Odontol Scand.* 2015;73:609–15. <https://doi.org/10.3109/00016357.2015.1020339>.
 189. Selba MC, Bryson ER, Rosenberg CL, Heng HG, DeLeon VB. Selective breeding in domestic dogs: how selecting for a short face impacted canine neuroanatomy. *Anat Rec (Hoboken).* 2021;304:101–15. <https://doi.org/10.1002/ar.24471>.
 190. Inchingolo AD, Carpentiere V, Piras F, Netti A, Ferrara I, Campanelli M, Latini G, Viapiano F, Costa S, Malcangi G, et al. Orthodontic surgical treatment of impacted mandibular canines: systematic review and case report. *Appl Sci.* 2022;12:8008. <https://doi.org/10.3390/app12168008>.
 191. Inchingolo AD, Gargiulo CI, Malcangi G, Ciocia AM, Patano A, Azzollini D, Piras F, Barile G, Settanni V, Mancini A, et al. Diagnosis of SARS-CoV-2 during the pandemic by multiplex RT-PCR hCoV test: future perspectives. *Pathogens.* 2022;11:1378. <https://doi.org/10.3390/pathogen1111378>.
 192. Inchingolo AM, Malcangi G, Inchingolo AD, Mancini A, Palmieri G, Di Pede C, Piras F, Inchingolo F, Dipalma G, Patano A. Potential of graphene-functionalized titanium surfaces for dental implantology: systematic review. *Coatings.* 2023;13:725. <https://doi.org/10.3390/coatings13040725>.
 193. Inchingolo, A.D.; Malcangi, G.; Ceci, S.; Patano, A.; Corriero, A.; Vimercati, L.; Azzollini, D.; Marinelli, G.; Coloccia, G.; Piras, F.; et al. Effectiveness of SARS-CoV-2 Vaccines for Short- and Long-Term Immunity: A General Overview for the Pandemic Contrast. *Int J Mol Sci* 2022, 23, <https://doi.org/10.3390/ijms23158485>.
 194. Tepedino M, Laurenziello M, Guida L, Montaruli G, Grassia V, Chimenti C, Campanelli M, Ciavarella D. Sella turcica and craniofacial morphology in patients with palatally displaced canines: a retrospective study. *Folia Morphol (Warsz).* 2020;79:51–7. <https://doi.org/10.5603/FM.a2019.0050>.
 195. Scribante A, Sfondrini MF, Cassani M, Fraticelli D, Beccari S, Gandini P. Sella turcica bridging and dental anomalies: is there an association? *Int J Paediatr Dent.* 2017;27:568–73. <https://doi.org/10.1111/ipd.12301>.
 196. Yang S, Yang X, Jin A, Ha N, Dai Q, Zhou S, Yang Y, Gong X, Hong Y, Ding Q, et al. Sequential traction of a labio-palatal horizontally impacted maxillary canine with a custom three-directional force device in the space of a missing ipsilateral first premolar. *Korean J Orthod.* 2019;49:124–36. <https://doi.org/10.4041/kjod.2019.49.2.124>.
 197. LeBlanc AK, LeBlanc CJ, Rohrbach BW, Kania SA. Serial evaluation of neutrophil function in tumour-bearing dogs undergoing chemotherapy. *Vet Comp Oncol.* 2015;13:20–7. <https://doi.org/10.1111/vco.12015>.
 198. Baidas LF, Alshihah N, Alabdulaly R, Mutaieb S. Severity and treatment difficulty of impacted maxillary canine among orthodontic patients in Riyadh, Saudi Arabia. *Int J Environ Res Public Health.* 2022;19:10680. <https://doi.org/10.3390/ijerph191710680>.
 199. Schneider S, Gandhi V, Upadhyay M, Allareddy V, Tadinada A, Yadav S. Sex-, growth pattern-, and growth status-related variability in maxillary and mandibular buccal cortical thickness and density. *Korean J Orthod.* 2020;50:108–19. <https://doi.org/10.4041/kjod.2020.50.2.108>.
 200. Crescini A, Nieri M, Buti J, Baccetti T, Mauro S, Prato GPP. Short- and long-term periodontal evaluation of impacted canines treated with a closed surgical-orthodontic approach. *J Clin Periodontol.* 2007;34:232–42. <https://doi.org/10.1111/j.1600-051X.2006.01042.x>.
 201. Shephard M, Coleman H. Simultaneous adenomatoid odontogenic and keratocystic odontogenic tumours in a patient with gordin-goltz syndrome. *Aust Dent J.* 2014;59:121–4. <https://doi.org/10.1111/adj.12137>.
 202. Datarkar A, Kolerman R, Lorean A, Segal P, Lucchina AG, Mortellaro C, Mijiritsky E. Simultaneous removal of horizontally impacted maxillary canine and placement of an immediately loaded implant. *J Craniofac Surg.* 2015;26:e657. <https://doi.org/10.1097/SCS.0000000000001959>.
 203. Montes-Díaz ME, Martínez-González A, Arriazu-Navarro R, Alvarado-Lorenzo A, Gallardo-López NE, Ortega-Aranegui R. Skeletal and dental morphological characteristics of the maxillary in patients with impacted canines using cone beam computed tomography: a retrospective clinical study. *J Pers Med.* 2022;12:96. <https://doi.org/10.3390/jpm12010096>.
 204. Hecht G, Sarbo N, Svoboda W, Mead HL, Ruberto I, Altin JA, Engelthaler DM, Venkat H, Yaglom HD. “Sniffing” out SARS-CoV-2 in arizona working dogs: an exploratory serosurvey. *Front Vet Sci.* 2023;10:1166101. <https://doi.org/10.3389/fvets.2023.1166101>.
 205. Naoumova J, Kuroi J, Kjellberg H. Extraction of the deciduous canine as an interceptive treatment in children with palatal displaced canines—part I: shall we extract the deciduous canine or not? *Eur J Orthod.* 2015;37:209–18. <https://doi.org/10.1093/ejo/cju040>.
 206. Al-Nimri K, Gharaibeh T. Space conditions and dental and occlusal features in patients with palatally impacted maxillary canines: an aetiological study. *Eur J Orthod.* 2005;27:461–5. <https://doi.org/10.1093/ejo/cji022>.
 207. Brüllmann D, Schulze RKW. Spatial resolution in CBCT machines for dental/maxillofacial applications-what do we know today? *Dentomaxillofac Radiol.* 2015;44:20140204. <https://doi.org/10.1259/dmfr.20140204>.
 208. Verma SK, Maheshwari S, Singh RK, Chaudhari PK. Laser in dentistry: an innovative tool in modern dental practice. *Natl J Maxillofac Surg.* 2012;3:124–32. <https://doi.org/10.4103/0975-5950.111342>.
 209. Arias-Orozco P, Bástida-González F, Cruz L, Villatoro J, Espinoza E, Zárate-Segura PB, Recuenco S. Spatiotemporal analysis of canine rabies in el salvador: violence and poverty as social factors of canine rabies. *PLoS ONE.* 2018;13:e0201305. <https://doi.org/10.1371/journal.pone.0201305>.
 210. Pauwels R, Araki K, Siewerdsen JH, Thongvigitmanee SS. Technical aspects of dental CBCT: state of the art. *Dentomaxillofac Radiol.* 2015;44:20140224. <https://doi.org/10.1259/dmfr.20140224>.
 211. Handelman, C.S. The Anterior Alveolus: Its Importance in Limiting Orthodontic Treatment and Its Influence on the Occurrence of Iatrogenic Sequelae. *Angle Orthod* 1996, 66, 95–109; discussion 109–110, [https://doi.org/10.1043/0003-3219\(1996\)066<0095:TAAll>2.3.CO;2](https://doi.org/10.1043/0003-3219(1996)066<0095:TAAll>2.3.CO;2).
 212. Jung YH, Liang H, Benson BW, Flint DJ, Cho BH. The Assessment of impacted maxillary canine position with panoramic radiography and cone beam CT. *Dentomaxillofac Radiol.* 2012;41:356–60. <https://doi.org/10.1259/dmfr.14055036>.
 213. Corriero A, Gadaleta RM, Puntillo F, Inchingolo F, Moschetta A, Brienza N. The central role of the gut in intensive care. *Crit Care.* 2022;26:379. <https://doi.org/10.1186/s13054-022-04259-8>.
 214. Bowman SJ, Johnston LE. The esthetic impact of extraction and nonextraction treatments on caucasian patients. *Angle Orthod.* 2000;70:3–10. [https://doi.org/10.1043/0003-3219\(2000\)070%3c0003:TEIOEA%3e2.0.CO;2](https://doi.org/10.1043/0003-3219(2000)070%3c0003:TEIOEA%3e2.0.CO;2).
 215. Lembo T, Hampson K, Kaare MT, Ernest E, Knobel D, Kazwala RR, Haydon DT, Cleaveland S. The feasibility of canine rabies elimination in africa: dispelling doubts with data. *PLoS Negl Trop Dis.* 2010;4:e626. <https://doi.org/10.1371/journal.pntd.0000626>.
 216. Altini M, Cohen M. The follicular primordial cyst - odontogenic keratocyst. *Int J Oral Surg.* 1982;11:175–82. [https://doi.org/10.1016/s0300-9785\(82\)80005-7](https://doi.org/10.1016/s0300-9785(82)80005-7).
 217. Inchingolo AD, Inchingolo AM, Piras F, Malcangi G, Patano A, Di Pede C, Netti A, Ciocia AM, Corriero A, Semjonova A, et al. A systematic review of positional plagiocephaly prevention methods for patients in development. *Appl Sci (Switzerland).* 2022;12:11172. <https://doi.org/10.3390/app122111172>.
 218. Patano A, Cardarelli F, Montenegro V, Ceci S, Inchingolo AD, Semjonova A, Palmieri G, Di Pede C, Mancini A, Maggiore ME, et al. Early functional

- orthodontic treatment of bad oral habits with AMCOP® bio-activators. *J Biol Regul Homeost Agents*. 2022;36:91–110.
219. Dipalma G, Inchingolo AD, Inchingolo AM, Piras F, Carpentiere V, Garofoli G, Azzollini D, Campanelli M, Paduanelli G, Palermo A, et al. Artificial intelligence and its clinical applications in orthodontics: a systematic review. *Diagnostics*. 2023;13:3677. <https://doi.org/10.3390/diagnostics13243677>.
 220. Inchingolo F, Inchingolo AD, Palumbo I, Trilli I, Guglielmo M, Mancini A, Palermo A, Inchingolo AM, Dipalma G. The impact of cesarean section delivery on intestinal microbiota: mechanisms, consequences, and perspectives—a systematic review. *Int J Mol Sci*. 2024;25:1055. <https://doi.org/10.3390/ijms25021055>.
 221. Thilander B, Myrberg N. The prevalence of malocclusion in Swedish schoolchildren. *Scand J Dent Res*. 1973;81:12–21. <https://doi.org/10.1111/j.1600-0722.1973.tb01489.x>.
 222. Kim Y, Park JU, Kook Y-A. Alveolar bone loss around incisors in surgical skeletal class III patients. *Angle Orthod*. 2009;79:676–82. <https://doi.org/10.2319/070308-341.1>.
 223. Inchingolo AM, Patano A, Piras F, Mancini A, Inchingolo AD, Paduanelli G, Inchingolo F, Palermo A, Dipalma G, Malcangi G. The bad relationship between microbiota-gut-brain axis and autism spectrum disorder comparing therapeutic options: a scoping review. *Microorganisms*. 2023;11:1477. <https://doi.org/10.3390/microorganisms11061477>.
 224. Gargiulo Isacco C, Inchingolo AD, Nguyen Cao KD, Malcangi G, Paduanelli G, Pham Hung V, Tran Cong T, Bordea IR, Scarano A, Laforgia A, et al. The bad relationship, osteo-decay and diabetes type 2 searching for a link: a literature review. *J Biol Regul Homeost Agents*. 2021;35:253–69. <https://doi.org/10.2381/21-2suppl-26>.
 225. Inchingolo F, Dipalma G, Paduanelli G, De Oliveira LA, Inchingolo AM, Georgakopoulos PI, Inchingolo AD, Malcangi G, Athanasiou E, Fotopoulou E, et al. Computer-Based quantification of an atraumatic sinus augmentation technique using CBCT. *J Biol Regul Homeost Agents*. 2019;33:31–9 (Dental Supplement).
 226. Strbac GD, Foltin A, Gahleitner A, Bantleon H-P, Watzek G, Bernhart T. The prevalence of root resorption of maxillary incisors caused by impacted maxillary canines. *Clin Oral Investig*. 2013;17:553–64. <https://doi.org/10.1007/s00784-012-0738-9>.
 227. Mason C, Papadakou P, Roberts GJ. The radiographic localization of impacted maxillary canines: a comparison of methods. *Eur J Orthod*. 2001;23:25–34. <https://doi.org/10.1093/ejo/23.1.25>.
 228. Dağsuyu İM, Okşayan R, Kahraman F, Aydın M, Bayraktar İŞ, Uğurlu M. The relationship between dental follicle width and maxillary impacted canines' descriptive and resorptive features using cone-beam computed tomography. *Biomed Res Int*. 2017;2017:2938691. <https://doi.org/10.1155/2017/2938691>.
 229. Preda L, La Fianza A, Di Maggio EM, Dore R, Schifino MR, Campani R, Segù C, Sfondrini MF. The use of spiral computed tomography in the localization of impacted maxillary canines. *Dentomaxillofac Radiol*. 1997;26:236–41. <https://doi.org/10.1038/sj.dmf.4600258>.
 230. Pithon MM. Therapeutic approach in class I malocclusion with impacted maxillary canines. *Dental Press J Orthod*. 2022;27:e22bbo2. <https://doi.org/10.1590/2177-6709.27.2.e22bbo2>.
 231. Laforgia A, Inchingolo AD, Piras F, Colonna V, Giorgio RV, Carone C, Rapone B, Malcangi G, Inchingolo AM, Inchingolo F, et al. Therapeutic strategies and genetic implications for periodontal disease management: a systematic review. *Int J Mol Sci*. 2024;25:7217. <https://doi.org/10.3390/ijms25137217>.
 232. Baart JA, Bakker CJ, Prah-Andersen B. Therapeutic strategies for impacted maxillary canines. *Ned Tijdschr Tandheelkd*. 2000;107:90–3.
 233. Oberoi S, Gill P, Chigurupati R, Hoffman WY, Hatcher DC, Vargervik K. Three-dimensional assessment of the eruption path of the canine in individuals with bone-grafted alveolar clefts using cone beam computed tomography. *Cleft Palate Craniofac J*. 2010;47:507–12. <https://doi.org/10.1597/08/171>.
 234. Almuhtaseb E, Mao J, Mahony D, Bader R, Zhang Z-X. Three-dimensional localization of impacted canines and root resorption assessment using cone beam computed tomography. *J Huazhong Univ Sci Technol Med Sci*. 2014;34:425–30. <https://doi.org/10.1007/s11596-014-1295-z>.
 235. Hampson K, Dushoff J, Cleaveland S, Haydon DT, Kaare M, Packer C, Dobson A. Transmission dynamics and prospects for the elimination of canine rabies. *PLoS Biol*. 2009;7:e53. <https://doi.org/10.1371/journal.pbio.1000053>.
 236. Oxlade O, Murray M. Tuberculosis and poverty: why are the poor at greater risk in India? *PLoS ONE*. 2012;7:e47533. <https://doi.org/10.1371/journal.pone.0047533>.
 237. Inchingolo AM, Inchingolo AD, Viapiano F, Ciocia AM, Ferrara I, Netti A, Dipalma G, Palermo A, Inchingolo F. Treatment approaches to molar incisor hypomineralization: a systematic review. *J Clin Med*. 2023;12:7194. <https://doi.org/10.3390/jcm12227194>.
 238. Botticelli S, Verna C, Cattaneo PM, Heidmann J, Melsen B. Two- versus three-dimensional imaging in subjects with unerupted maxillary canines. *Eur J Orthod*. 2011;33:344–9. <https://doi.org/10.1093/ejo/cjq102>.
 239. Garcovich D, Aiuto R, Serrano E, Re D. Uncovering and autonomous eruption of palatally impacted canines. *Eur J Paediatr Dentist*. 2018;19:300–2. <https://doi.org/10.23804/ejpd.2018.19.04.9>.
 240. Katsnelson A, Flick WG, Susarla S, Tartakovsky JV, Miloro M. Use of panoramic X-Ray to determine position of impacted maxillary canines. *J Oral Maxillofac Surg*. 2010;68:996–1000. <https://doi.org/10.1016/j.joms.2009.09.022>.
 241. Patano A, Malcangi G, Sardano R, Mastrodonato A, Garofoli G, Mancini A, Inchingolo AD, Di Venere D, Inchingolo F, Dipalma G, et al. White spots: prevention in orthodontics—systematic review of the literature. *Int J Environ Res Public Health*. 2023;20:5608. <https://doi.org/10.3390/ijerph20085608>.
 242. Casu C, Mannu C. Atypical afta major healing after photodynamic therapy. *Case Rep Dent*. 2017;2017:8517470. <https://doi.org/10.1155/2017/8517470>.
 243. Casu C, Murgia MS, Orrù G, Scano A. Photodynamic therapy for the successful management of cyclosporine-related gum hypertrophy: a novel therapeutic option. *J Public Health Res*. 2022;11:22799036221116176. <https://doi.org/10.1177/22799036221116177>.
 244. Inchingolo AD, Ferrara I, Viapiano F, Netti A, Campanelli M, Buongiorno S, Latini G, Carpentiere V, Ciocia AM, Ceci S, et al. Rapid maxillary expansion on the adolescent patient: systematic review and case report. *Children (Basel)*. 2022;9:1046. <https://doi.org/10.3390/bambini9071046>.
 245. Laudadio C, Inchingolo AD, Malcangi G, Limongelli L, Marinelli G, Colocchia G, Montenegro V, Patano A, Inchingolo F, Bordea IR, et al. Management of anterior open-bite in the deciduous, mixed and permanent dentition stage: a descriptive review. *J Biol Regul Homeost Agents*. 2021;35:271–81. <https://doi.org/10.23812/21-2suppl1-27>.
 246. Contaldo M, Itro A, Lajolo C, Gioco G, Inchingolo F, Serpico R. Overview on osteoporosis, periodontitis and oral dysbiosis: the emerging role of oral microbiota. *Appl Sci*. 2020;10:6000. <https://doi.org/10.3390/app10176000>.
 247. Contaldo M, Della Vella F, Raimondo E, Minervini G, Buljubasic M, Ogodescu A, Sinescu C, Serpico R. Early Childhood Oral Health Impact Scale (ECHOHIS): literature review and Italian validation. *Int J Dent Hyg*. 2020;18:396–402. <https://doi.org/10.1111/idh.12451>.
 248. Grippaudo MM, Quinzi V, Manai A, Paolantonio EG, Valente F, La Torre G, Marzo G. Orthodontic treatment need and timing: assessment of evolutive malocclusion conditions and associated risk factors. *Eur J Paediatr Dent*. 2020;21:203–8. <https://doi.org/10.23804/ejpd.2020.21.03.09>.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.