



# A general framework for solving differential equations

Luigi Brugnano<sup>1</sup> · Felice Iavernaro<sup>2</sup>

Received: 4 February 2022 / Accepted: 8 June 2022 / Published online: 30 June 2022  
© The Author(s) 2022

## Abstract

Recently, the efficient numerical solution of Hamiltonian problems has been tackled by defining the class of energy-conserving Runge-Kutta methods named *Hamiltonian Boundary Value Methods (HBVMs)*. Their derivation relies on the expansion of the vector field along a suitable orthonormal basis. Interestingly, this approach can be extended to cope with more general differential problems. In this paper we sketch this fact, by considering some relevant examples.

**Keywords** Hamiltonian Boundary Value Methods · HBVMs · Line integral methods · Local Fourier expansion · Spectral methods

**Mathematics Subject Classification** 65P10 · 65L05

## 1 Introduction

The numerical solution of Hamiltonian problems has been the subject of many investigations in the last decades, due to the fact that Hamiltonian problems are not structurally stable against generic perturbations like those induced by a general-purpose numerical method used to approximate their solutions.

The main features of a canonical Hamiltonian system are surely the symplecticness of the map and the conservation of energy. It is well-known that these two properties are mutually exclusive when Runge-Kutta or B-series methods are considered [1]. This is the reason why numerical methods have been devised in order of either defining a

---

Dedicated to the memory of professor Ilio Galligani.

---

✉ Luigi Brugnano  
luigi.brugnano@unifi.it

✉ Felice Iavernaro  
felice.iavernaro@uniba.it

<sup>1</sup> Dipartimento di Matematica e Informatica “U. Dini”, Università di Firenze, Firenze, Italy

<sup>2</sup> Dipartimento di Matematica, Università di Bari, Bari, Italy

symplectic discrete map, giving rise to the class of *symplectic methods* (see, e.g., the monographs [2–5], and references therein, and the review paper [6]), or being able to conserve the energy, resulting in the class of *energy-conserving methods* (see, e.g., [7–14] and the monograph [15]).

The methods we shall deal with belong to the class of energy-conserving methods: they are named *Hamiltonian Boundary Value Methods (HBVMs)* [12, 13, 15, 16]. HBVMs, in turn, can be also regarded as a local projection of the vector field onto a finite-dimensional function space [17]. It is worth mentioning that the same idea has been also named in different ways (e.g., *functionally fitted methods* [18]). Remarkably, this strategy is even more general and can be adapted to other classes of problems, also allowing the use of the methods as spectral methods in time [19–22]. This means that the accuracy of the approximation is attained by increasing the dimension of the functional space, rather than reducing the timestep. Test problems, comparing the computational cost of such methods w.r.t. collocation or other standard methods, can be found in [22], showing the good potentialities of the spectral approach. We mention that spectral methods have been also extensively used in other settings [23–26].

In this paper, we sketch how the approach resulting into HBVMs can be extended to cope with additional differential problems: for some of such instances a thorough analysis is still in progress, but the general framework can be outlined even at this stage.

With this premise, the structure of the paper is as follows: in Sect. 2 we provide a brief introduction to HBVMs; in Sect. 3 the approach is extended to cope with general *initial value problems (ODE-IVPs)*, including *special second-order problems* and *delay differential equations (DDEs)*; Sect. 4 is devoted to *implicit differential algebraic equations (IDEs/DAEs)*, including linearly implicit DAEs; Sect. 5 concerns the case of *fractional differential equations (FDEs)*; a few conclusions are given in Sect. 6.

## 2 Hamiltonian problems

A canonical Hamiltonian problem is in the form

$$\dot{y} = J\nabla H(y), \quad y(0) = y_0 \in \mathbb{R}^{2m}, \quad J = \begin{pmatrix} & I_m \\ -I_m & \end{pmatrix} = -J^\top, \quad (1)$$

with  $H : \mathbb{R}^{2m} \rightarrow \mathbb{R}$  the Hamiltonian function (or *energy*) of the system.<sup>1</sup> As is easily understood  $H$  is a constant of motion, since

$$\frac{d}{dt} H(y) = \nabla H(y)^\top \dot{y} = \nabla H(y)^\top J \nabla H(y) = 0,$$

being  $J$  skew-symmetric. The simple idea on which HBVMs rely on is that of reformulating the previous conservation property in terms of the line integral of  $\nabla H$  along

<sup>1</sup> In fact, for isolated mechanical systems,  $H$  has the physical meaning of the total energy.

the path defined by the solution  $y(t)$ :

$$H(y(t)) = H(y_0) + \int_0^t \nabla H(y(\tau))^\top \dot{y}(\tau) d\tau.$$

Clearly, the integral at the right-hand side of the previous equality vanishes, because the integrand is identically zero, so that the conservation holds true for all  $t > 0$ . The solution of (1) is the unique function satisfying such a conservation property for all  $t > 0$ . However, if we consider a discrete-time dynamics, ruled by a *timestep*  $h$ , there exists infinitely many paths  $\sigma$  such that:

$$\begin{aligned} \sigma(0) &= y_0, & \sigma(h) &=: y_1 \approx y(h), \\ 0 &= h \int_0^1 \nabla H(\sigma(ch))^\top \dot{\sigma}(ch) dc. \end{aligned} \quad (2)$$

The path  $\sigma$  obviously defines a one-step numerical method that conserves the energy, since

$$H(y_1) = H(y_0) + h \int_0^1 \nabla H(\sigma(ch))^\top \dot{\sigma}(ch) dc = H(y_0),$$

even though now the integrand is no more identically zero. The methods derived in this framework have been named *line integral methods*, due to the fact that the path  $\sigma$  is defined so that the line integral (2) vanishes. Such methods have been thoroughly analyzed in the monograph [15] (see also the review paper [27]). Clearly, in the practical implementation of the methods, the integral is replaced by a quadrature of enough high-order, thus providing a fully discrete method. However, for the sake of brevity, we shall hereafter continue to use the integral, and not the quadrature, so that the exposition is greatly simplified and more concise.<sup>2</sup>

After some initial attempts to derive methods in this class [8–10, 12, 30], a systematic way for their derivation was found in [17], which is based on a local Fourier expansion of the vector field in (1). In fact, by setting

$$f(y) = J\nabla H(y), \quad (3)$$

and using hereafter the more convenient notation, depending on the needs, one may rewrite problem (1), on the interval  $[0, h]$ , as:

$$\dot{y}(ch) = \sum_{j \geq 0} P_j(c) \gamma_j(y), \quad c \in [0, 1], \quad y(0) = y_0, \quad (4)$$

<sup>2</sup> This amounts to study HBVMs as *continuous-stage Runge-Kutta methods*, as it has been done in [28, 29].

where  $\{P_j\}_{j \geq 0}$  is the Legendre orthonormal polynomial basis on  $[0, 1]$ ,

$$P_i \in \Pi_i, \quad \int_0^1 P_i(x)P_j(x)dx = \delta_{ij}, \quad i, j \geq 0, \tag{5}$$

$\Pi_i$  is the vector space of polynomials of degree at most  $i$ ,  $\delta_{ij}$  is the Kronecker symbol, and

$$\gamma_j(y) = \int_0^1 P_j(\tau)f(y(\tau h))d\tau, \quad j \geq 0, \tag{6}$$

are the corresponding Fourier coefficients. The solution of the problem is formally obtained, in terms of the unknown Fourier coefficients, by integrating both sides of Eq. (4):

$$y(ch) = y_0 + h \sum_{j \geq 0} \int_0^c P_j(x)dx \gamma_j(y), \quad c \in [0, 1]. \tag{7}$$

A polynomial approximation  $\sigma \in \Pi_s$  can be formally obtained by truncating the previous series to a finite sum,

$$\dot{\sigma}(ch) = \sum_{j=0}^{s-1} P_j(c)\gamma_j(\sigma), \quad c \in [0, 1], \quad \sigma(0) = y_0, \tag{8}$$

and

$$\sigma(ch) = y_0 + h \sum_{j=0}^{s-1} \int_0^c P_j(x)dx \gamma_j(\sigma), \quad c \in [0, 1], \tag{9}$$

respectively, with  $\gamma_j(\sigma)$  defined according to (6), upon replacing  $y$  with  $\sigma$ . Whichever the degree  $s \geq 1$  of the polynomial approximation, the following result holds true, where the same notation used above holds.

**Theorem 1** *Assume that the Hamiltonian  $H$  has continuous partial derivatives. Then,  $H(y_1) = H(y_0)$ .*

**Proof** In fact, one has:

$$\begin{aligned} H(y_1) - H(y_0) &= H(\sigma(h)) - H(\sigma(0)) = h \int_0^1 \nabla H(\sigma(ch))^\top \dot{\sigma}(ch)dc \\ &= h \int_0^1 \nabla H(\sigma(ch))^\top \sum_{j=0}^{s-1} P_j(c)\gamma_j(\sigma)dc \\ &= h \sum_{j=0}^{s-1} \left[ \int_0^1 P_j(c)\nabla H(\sigma(ch))dc \right]^\top \gamma_j(\sigma) \end{aligned}$$

$$\begin{aligned}
 &= h \sum_{j=0}^{s-1} \left[ \int_0^1 P_j(c) \nabla H(\sigma(ch)) dc \right]^T J \left[ \int_0^1 P_j(c) \nabla H(\sigma(ch)) dc \right] \\
 &= 0,
 \end{aligned}$$

due to the fact that  $J$  is skew-symmetric. □

The next result states that the methods has order  $2s$ .

**Theorem 2** *Assume that the function (3) is of class  $C^{(2s+1)}[0, h]$ . Then,  $y_1 - y(h) = O(h^{2s+1})$ .*

**Proof** See [17, Theorem 1]. □

**Remark 1** It is worth noticing that, due to the fact that (see (5))

$$\int_0^1 P_j(x) dx = \delta_{j0}, \tag{10}$$

the new approximation (see (2) and (9)) is given by

$$y_1 := \sigma(h) \equiv y_0 + \int_0^h f(\sigma(t)) dt,$$

which can be regarded as an approximation of the Fundamental Theorem of Calculus.

### 3 Initial value problems

The result in Theorem 2 does not depend on the Hamiltonian structure of the problem. This suggests that the above procedure also makes sense for a generic initial value problem which, on the interval  $[0, h]$  can be cast, without loss of generality, as

$$\dot{y}(ch) = f(y(ch)), \quad c \in [0, 1], \quad y(0) = y_0 \in \mathbb{R}^m. \tag{11}$$

To cope with more general problems, hereafter we consider an alternative but equivalent approach to devise this family of methods. Consider again

$$\dot{\sigma}(ch) = \sum_{j=0}^{s-1} P_j(c) \gamma_j, \quad \sigma(ch) = y_0 + h \sum_{j=0}^{s-1} \int_0^c P_j(x) dx \gamma_j, \quad c \in [0, 1]. \tag{12}$$

We determine the unknown coefficients

$$\gamma_j \equiv \gamma_j(\sigma), \quad j = 0, \dots, s - 1,$$

in order that the residual function

$$r(ch) := \dot{\sigma}(ch) - f(\sigma(ch)), \quad c \in [0, 1], \tag{13}$$

be orthogonal to  $\Pi_{s-1}$ , namely

$$0 = \int_0^1 P_i(c)r(ch)dc = \int_0^1 P_i(c) (\dot{\sigma}(ch) - f(\sigma(ch))) dc, \quad i = 0, \dots, s - 1.$$

Substituting the expansion for  $\dot{\sigma}(ch)$  in (12) and taking into account the orthogonality relations (5) again lead to the equations defining the unknown coefficients  $\gamma_i$  (compare with (6)):

$$\gamma_i = \int_0^1 P_i(c)f \left( y_0 + h \sum_{j=0}^{s-1} \int_0^c P_j(x)dx \gamma_j \right) dc, \quad i = 0, \dots, s - 1. \quad (14)$$

**Remark 2** As was anticipated above, in order to get a fully discrete problem, the integrals appearing in (14) need to be approximated by suitable quadrature formulae: to this purpose, a Gauss-Legendre quadrature of order  $2k$  is employed. The resulting integrator, first devised in the context of Hamiltonian systems, is denoted by HBVM( $k, s$ ) to highlight the dependence on the two discretization parameters. When  $k = s$  the method reduces to the  $s$ -stage Gauss-Legendre collocation method but, if necessary, one can choose  $k$  large enough to ensure that the integrals are approximated to within machine precision. This, in turn, can be done without increasing too much the overall computational cost: as matter of fact, the discretized problem (14) has (block) dimension  $s$ , *independently* of  $k$ . This fact is an important feature, in the case of Hamiltonian problems, as the next Example 1 will show. The resulting set of  $s$  nonlinear equations (14) can be efficiently solved by means of the *blended iteration* described in [31], adapted from [32, 33] (see also [34] for related approaches): in fact, such iteration only requires the factorization of one matrix of the size of the continuous problem, *whichever is the value of  $s$  considered*. We mention that the state-of-art Matlab code `hbvm.m`, implementing HBVMs, is available at the website of the monograph [15].

We observe that, in light of (14) and (12), the new approximation can be written as

$$y_1 = y_0 + h\gamma_0, \quad (15)$$

which is easily evaluated, once the system (14) has been solved.

It is worth mentioning that, in the actual implementation of (12), it is conveniently exploited the known relation between the Legendre polynomials and their integrals:

$$\begin{aligned} \int_0^c P_0(x)dx &= \xi_0 P_0(c) + \xi_1 P_1(c), \\ \int_0^c P_j(x)dx &= \xi_{j+1} P_{j+1}(c) - \xi_j P_{j-1}(c), \quad j \geq 1, \\ \xi_i &= \frac{1}{2\sqrt{|4i^2 - 1|}}, \quad i \geq 0. \end{aligned} \quad (16)$$

**Example 1** Let us consider the Hamiltonian problem defined by [15]:

$$H(q, p) = (q^2 + p^2)^2 - 10(q^2 - p^2), \quad q, p \in \mathbb{R}. \tag{17}$$

When the initial point is chosen as


$$(q_0, p_0) = (0, 10^{-5}), \tag{18}$$

the solution trajectory, in the phase space, is a Cassini oval, as is shown on the left of Fig. 1, with very close branches near the origin (see the right-plot in the same figure). The solution is then periodic, with period  $T \approx 3.131990057003955$ . However, when numerically solving the problem, if the Hamiltonian is not precisely conserved, then the numerical branches may intersect, thus giving a wrong simulation. If we fix  $s = 2$  in (14) and we use a Gauss-Legendre formula of order 8 ( $k = 4$ ), then the HBVM(4,2) method is exactly energy conserving (according to Theorem 1), since the integrals are exact. We use this method and, for comparison, the 2-stage Gauss collocation method (i.e., HBVM(2,2)), with a timestep  $h = 10^{-2}$ . This latter method, though symplectic, is not energy-conserving. As is clearly shown in Fig. 2, the former method (left-plot) provides a correct numerical solution, whereas the latter (right-plot) does not. The two simulations have been done by using the Matlab code `hbvm.m` on an Intel 3GHz Intel Xeon W10 core computer, with 64GB of memory, running Matlab 2020b. Despite the very different outcomes, the computational time for the two simulations is approximately the same (about 10 sec.). This fact confirms that using values of  $k$  larger than  $s$ , for a HBVM( $k, s$ ) method, is not a big issue.


### 3.1 Special second-order problems

A relevant instance of ODE-IVPs is the case of *special second-order problems*, namely problems in the form

$$\ddot{y}(ch) = f(y(ch)), \quad c \in [0, 1], \quad y(0) = y_0, \quad \dot{y}(0) = z_0 \in \mathbb{R}^m. \tag{19}$$



**Fig. 1** Left-plot: solution of problem (17)–(18); right-plot: zoom near the origin



**Fig. 2** Numerical solution of problem (17)–(18) by using HBVM(4,2) (left) and HBVM(2,2) (right) with timestep  $h = 10^{-2}$

As an example, separable Hamiltonian problems fall in this class. The particular structure of the problem allows to gain efficiency, with respect to just recasting the problem as the following first order problem of doubled size:<sup>3</sup>

$$\begin{cases} \dot{y}(ch) = z(ch), \\ \dot{z}(ch) = f(y(ch)), \end{cases} \quad c \in [0, 1], \quad y(0) = y_0, \quad \dot{z}(0) = z_0 \in \mathbb{R}^m. \quad (20)$$

Proceeding as before, we introduce the polynomial approximations

$$\dot{y}(ch) \approx \dot{\sigma}(ch) := \sum_{j=0}^{s-1} P_j(c) dx \hat{\gamma}_j, \quad (21)$$

$$\dot{z}(ch) \approx \dot{\mu}(ch) := \sum_{j=0}^{s-1} P_j(c) dx \gamma_j, \quad c \in [0, 1], \quad (22)$$

satisfying the initial conditions  $\sigma(0) = y_0$ ,  $\mu(0) = z_0$ , and require the residual

$$r(ch) := \begin{pmatrix} \dot{\sigma}(ch) - \mu(ch) \\ \dot{\mu}(ch) - f(\sigma(ch)) \end{pmatrix}, \quad c \in [0, 1],$$

be orthogonal to  $\Pi_{s-1}$ . The  $s$  orthogonality conditions imposed on the first component of  $r(ch)$  allow us to remove the unknowns  $\hat{\gamma}_i$ :

$$\hat{\gamma}_i = \int_0^1 P_i(c) \mu(ch) dc, \quad i = 0, \dots, s - 1,$$

<sup>3</sup> Runge-Kutta-Nyström methods are in fact devised for this purpose.

so that, integrating both (21) and (22) from 0 to  $c$ , we get the two polynomial approximations of degree  $s$  for the solution and its derivative,

$$\sigma(ch) = y_0 + h \sum_{j=0}^{s-1} \int_0^c P_j(x) dx \int_0^1 P_j(\tau) \mu(\tau h) d\tau, \tag{23}$$

$$\mu(ch) = z_0 + h \sum_{j=0}^{s-1} \int_0^c P_j(x) dx \gamma_j, \quad c \in [0, 1], \tag{24}$$

involving only the  $s$  unknown coefficients  $\gamma_j, j = 0, \dots, s - 1$ . Such coefficients are determined from the orthogonality conditions imposed on the second component of  $r(ch)$ :

$$\gamma_i = \int_0^1 P_i(c) f(\sigma(ch)) dc, \quad i = 0, \dots, s - 1. \tag{25}$$

By taking into account (16) and (23)–(24), the expression of the polynomial  $\sigma$ , in terms of the above coefficients, turns out to be:

$$\begin{aligned} \sigma(ch) = & y_0 + chz_0 + c \frac{h^2}{2} \left( \gamma_0 - \frac{\gamma_1}{\sqrt{3}} \right) \\ & + h^2 \sum_{j=1}^{s-2} \int_0^c P_j(x) dx (\xi_j \gamma_{j-1} - \xi_{j+1} \gamma_{j+1}) \\ & + h^2 \int_0^c P_{s-1}(x) dx \xi_{s-1} \gamma_{s-2}. \end{aligned} \tag{26}$$

Once the system (25)–(26) has been solved, the new approximations are obtained, as done before, as

$$\mu(h) =: z_1 \approx \dot{y}(h), \quad \sigma(h) =: y_1 \approx y(h),$$

i.e., recalling (10),

$$z_1 = z_0 + h\gamma_0, \quad y_1 = y_0 + hz_0 + \frac{h^2}{2} \left( \gamma_0 - \frac{\gamma_1}{\sqrt{3}} \right).$$

**Remark 3** Also in this case, there exist efficient iterative procedures for solving the discrete counterpart of (25) (see [31]).<sup>4</sup> In particular, the modified *blended iteration* follows from the analysis in [35] (see also [33]).

We conclude this section by mentioning that the whole approach can be extended, in a straightforward way, to higher-order equations [29].

<sup>4</sup> This discrete counterpart, as usual, is obtained by approximating the involved integrals by using a Gauss-Legendre quadrature of order  $2k$ , thus resulting into a HBVM( $k, s$ ) method for special second-order problems.

### 3.2 Delay differential equations

Delay differential equations have been the subject of many investigations in the past years (we refer to the monographs [36, 37] and references therein for further informations). Here, we extend the approach described in the previous sections to cope with delay differential equations in the form

$$\dot{y}(t) = f(y(t), y(t - \tau)), \quad t \geq 0, \quad y(t) = \phi(t), \quad t \in [-\tau, 0], \quad (27)$$

with  $\tau > 0$  a fixed delay, and  $\phi : \mathbb{R} \rightarrow \mathbb{R}^m$  a known function. In fact, by using a timestep

$$h = \frac{\tau}{K},$$

for a suitable  $K \in \mathbb{N}$ , and setting

$$\sigma_n(ch) = y_{n-1} + h \sum_{j=0}^{s-1} \int_0^c P_j(x) dx \gamma_j^n, \quad c \in [0, 1],$$

the approximation to the solution over the interval<sup>5</sup>

$$[t_{n-1}, t_n], \quad t_n = nh, \quad n \geq 1 - K, \quad y_{n-1} := \sigma_{n-1}(h),$$

then, for  $n \geq 1$ , the unknown coefficients  $\gamma_i^n$ ,  $i = 0, \dots, s - 1$ , solve the system of equations:

$$\begin{aligned} \gamma_i^n &= \int_0^1 P_i(c) f \left( y_{n-1} + h \sum_{j=0}^{s-1} \int_0^c P_j(x) dx \gamma_j^n, \sigma_{n-K}(ch) \right) dc, \\ i &= 0, \dots, s - 1, \end{aligned} \quad (28)$$

with the new approximation given by  $\sigma_n(h) =: y_n \approx y(t_n)$ , i.e.,

$$y_n = y_{n-1} + h\gamma_0^n.$$

We refer to [38] for full details on this approach.

### 4 Implicit differential algebraic equations

The procedure defined above to cope with ODE-IVPs and DDE-IVPs can be quite easily extended to solve implicit differential equations, too. For sake of simplicity we continue to deal with initial value problems, even though different kind of conditions

<sup>5</sup> Actually, the solution is known from the initial condition in (27), for  $n = 1 - K, \dots, 0$ .

could in principle be used, as seen in [39] in the Hamiltonian case. Let us then consider the problem

$$F(\dot{y}(ch), y(ch)) = 0, \quad c \in [0, 1], \quad y(0) = y_0 \in \mathbb{R}^m, \quad (29)$$

with  $F : \mathbb{R}^m \times \mathbb{R}^m \rightarrow \mathbb{R}^m$ . We can then look for a polynomial approximation of degree  $s$  in the form (12), with the new approximation at  $h$  given by  $\sigma(h) =: y_1 \approx y(h)$ , formally still given by (15). As was done before, by requiring the corresponding residual be orthogonal to  $\Pi_{s-1}$ , the unknown coefficients  $\gamma_i, i = 0, \dots, s-1$ , are the solution of the system of equations:

$$0 = \int_0^1 P_i(c) F \left( \sum_{j=0}^{s-1} P_j(c) \gamma_j, y_0 + h \sum_{j=0}^{s-1} \int_0^c P_j(x) dx \gamma_j \right) dc, \\ i = 0, \dots, s-1. \quad (30)$$

This approach, which is still under investigation, will be fully analyzed elsewhere.

#### 4.1 Linearly implicit DAEs

An important simplification is obtained in the case of linearly implicit DAEs, for which the problem is assumed to be in the form

$$M \dot{y}(ch) = f(y(ch)), \quad c \in [0, 1], \quad y(0) = y_0 \in \mathbb{R}^m, \quad (31)$$

with  $M \in \mathbb{R}^{m \times m}$  a (possibly singular) known matrix. In this case, in fact, one may look again for a polynomial approximation of degree  $s$  in the form (12). As a result, we obtain that the unknown coefficients  $\gamma_i, i = 0, \dots, s-1$ , solve the system of equations:

$$M \gamma_i = \int_0^1 P_i(c) f \left( y_0 + h \sum_{j=0}^{s-1} \int_0^c P_j(x) dx \gamma_j \right) dc, \quad i = 0, \dots, s-1. \quad (32)$$

**Remark 4** It is worth mentioning that the iterative solution of (32) can be carried out via a modified *blended iteration*, w.r.t. that defined in [31] for HBVMs, by following similar steps as done in [40].

## 5 Fractional differential equations

Fractional differential equations deserve a separate treatment, due to the fact that the underlying differential operator differs from the usual one (see, e.g., [41]; moreover,

we refer to [42] for a review on numerical methods for such problems). Let us then consider, for a given  $\alpha \in (0, 1]$ , the FDE

$$D^\alpha y(ch) = f(y(ch)), \quad c \in [0, 1], \quad y(0) = y_0 \in \mathbb{R}^m, \quad (33)$$

where  $D^\alpha$  is the Caputo fractional derivative. This latter derivative is defined, for a given differentiable function  $g : \mathbb{R} \rightarrow \mathbb{R}^m$ , as

$$D^\alpha g(t) = \frac{1}{\Gamma(1 - \alpha)} \int_0^t (t - x)^{-\alpha} \dot{g}(x) dx.$$

The corresponding Riemann-Liouville integral of order  $\alpha$  is defined as

$$I^\alpha g(t) = \frac{1}{\Gamma(\alpha)} \int_0^t (t - x)^{\alpha-1} g(x) dx.$$

It is known that:

$$D^\alpha I^\alpha g(t) = g(t), \quad I^\alpha D^\alpha g(t) = g(t) - g(0).$$

With this premise, the polynomial approximation to the solution of (33) is now looked in the form

$$D^\alpha \sigma(ch) = \sum_{j=0}^{s-1} P_j(c) \gamma_j, \quad \sigma(ch) = y_0 + h^\alpha \sum_{j=0}^{s-1} I^\alpha P_j(c) \gamma_j, \quad c \in [0, 1], \quad (34)$$

instead of (12).

**Remark 5** Clearly, when  $\alpha = 1$  (12) and (34) coincide, due to the fact that

$$D^1 g(t) = \lim_{\alpha \rightarrow 1^-} D^\alpha g(t) = \dot{g}(t), \quad \text{and} \quad I^1 g(t) = \int_0^t g(x) dx.$$

As was done before, by formally plugging  $D^\alpha \sigma(ch)$  and  $\sigma(ch)$  into (33), the unknown coefficients  $\gamma_i, i = 0, \dots, s - 1$ , are determined by requiring the residual be orthogonal to  $\Pi_{s-1}$ , thus leading to the system of equations:

$$\gamma_i = \int_0^1 P_i(c) f \left( y_0 + h^\alpha \sum_{j=0}^{s-1} I^\alpha P_j(c) \gamma_j \right) dc, \quad i = 0, \dots, s - 1. \quad (35)$$

In the actual implementation of (35), it is useful knowing how to compute the fractional integrals of the Legendre polynomials.<sup>6</sup> For this purpose, the following known result will be useful:

<sup>6</sup> I.e., we need a result similar to (16) for the problem at hand.

$$I^\alpha x^\eta = \frac{\Gamma(\eta + 1)}{\Gamma(\alpha + \eta + 1)} x^{\alpha+\eta}, \quad \eta \geq 0. \quad (36)$$

In fact, by recalling the explicit expression of the Legendre polynomials,

$$P_j(c) = \sqrt{2j+1} \sum_{i=0}^j (-1)^{j+i} \frac{(j+i)!}{(j-i)!(i!)^2} c^i, \quad j \geq 0,$$

from (36) one derives:

$$I^\alpha P_j(c) = \sqrt{2j+1} c^\alpha \sum_{i=0}^j (-1)^{j+i} \frac{(j+i)!}{(j-i)!i!\Gamma(\alpha+i+1)} c^i.$$

This approach for solving FDEs, at first sketched in [19], will be analyzed in detail elsewhere. We mention that a related approach can be found in [43].

## 6 Conclusions

In this paper we have recalled the basic facts about the numerical solution of differential problems via the expansion of the derivative, either integer or fractional, along the Legendre polynomial basis. Integrating such an approximation then provides a polynomial approximation to the solution.<sup>7</sup> The coefficients of the polynomial approximation, in turn, are determined by imposing the orthogonality of the residual to a suitable polynomial space. In addition, we mention that the obtained methods can be also used as *spectral methods in time*, as was described in [20–22] in the case of Hamiltonian problems (see also [19, 44–46]). Future directions of investigation will concern the extension of this framework to different kinds of differential problems. Also the numerical solution of integral equations could be a further subject of investigation.

**Funding** Open access funding provided by Università degli Studi di Firenze within the CRUI-CARE Agreement.

## Declarations

**Conflict of Interest** The authors declare no conflict of interests.

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

<sup>7</sup> Actually, a polynomial-like approximation, in the case of FDEs.

## References

1. Chartier, P., Faou, E., Murua, A.: An algebraic approach to invariant preserving integrators: the case of quadratic and Hamiltonian invariants. *Numer. Math.* **103**, 575–590 (2006). <https://doi.org/10.1007/s00211-006-0003-8>
2. Sanz-Serna, J.M., Calvo, M.P.: *Numerical Hamiltonian Problems*. Chapman & Hall, London, UK (1994)
3. Leimkuhler, B., Reich, S.: *Simulating Hamiltonian Dynamics*. Cambridge University Press, Cambridge, UK (2004)
4. Hairer, E., Lubich, Ch., Wanner, G.: *Geometric Numerical Integration*, 2nd edn. Springer, Berlin, Germany (2006)
5. Blanes, S., Casas, F.: *A Concise Introduction to Geometric Numerical Integration*. CRC Press, Boca Raton, FL, USA (2016)
6. Sanz-Serna, J.M.: Symplectic Runge-Kutta schemes for adjoint equations, automatic differentiation, optimal control, and more. *SIAM Rev.* **58**, 3–33 (2016). <https://doi.org/10.1137/151002769>
7. McLachlan, R.L., Quispel, G.R.W., Robidoux, N.: Geometric integration using discrete gradients. *Phil. Trans. R. Soc. Lond. A* **357**, 1021–1045 (1999)
8. Iavernaro, F., Pace, B.:  $s$ -Stage trapezoidal methods for the conservation of Hamiltonian functions of polynomial type. *AIP Conf. Proc.* **936**, 603–606 (2007). <https://doi.org/10.1063/1.2790219>
9. Iavernaro, F., Pace, B.: Conservative block-Boundary Value Methods for the solution of polynomial Hamiltonian systems. *AIP Conf. Proc.* **1048**, 888–891 (2008). <https://doi.org/10.1063/1.2991075>
10. Iavernaro, F., Trigiante, D.: High-order Symmetric Schemes for the Energy Conservation of Polynomial Hamiltonian Problems. *JNAIAM J. Numer. Anal. Ind. Appl. Math.* **4**, 87–101 (2009)
11. Celledoni, E., McLachlan, R.I., McLaren, D.I., Owren, B., Quispel, G.R.W., Wright, W.M.: Energy-preserving Runge-Kutta methods. *M2AN Math. Model. Numer. Anal.* **43**, 645–649 (2009). <https://doi.org/10.1051/m2an/2009020>
12. Brugnano, L., Iavernaro, F., Trigiante, D.: Hamiltonian BVMs (HBVMs): A family of “drift-free” methods for integrating polynomial Hamiltonian systems. *AIP Conf. Proc.* **1168**, 715–718 (2009). <https://doi.org/10.1063/1.3241566>
13. Brugnano, L., Iavernaro, F., Trigiante, D.: Hamiltonian Boundary Value Methods (Energy Preserving Discrete Line Integral Methods. *JNAIAM J. Numer. Anal. Ind. Appl. Math.* **5**, 17–37 (2010)
14. Hairer, E.: Energy preserving variant of collocation methods. *JNAIAM J. Numer. Anal. Ind. Appl. Math.* **5**, 73–84 (2010)
15. Brugnano, L., Iavernaro, F.: *Line Integral Methods for Conservative Problems*. Chapman et Hall/CRC, Boca Raton, FL, USA (2016). <http://web.math.unifi.it/users/brugnano/LIMbook/>
16. Brugnano, L., Iavernaro, F., Trigiante, D.: The lack of continuity and the role of infinite and infinitesimal in numerical methods for ODEs: The case of symplecticity. *Appl. Math. Comput.* **218**, 8053–8063 (2012). <https://doi.org/10.1016/j.amc.2011.03.022>
17. Brugnano, L., Iavernaro, F., Trigiante, D.: A simple framework for the derivation and analysis of effective one-step methods for ODEs. *Appl. Math. Comput.* **218**, 8475–8485 (2012). <https://doi.org/10.1016/j.amc.2012.01.074>
18. Li, Y.-W., Wu, X.: Functionally fitted energy-preserving methods for solving oscillatory nonlinear Hamiltonian systems. *SIAM J. Numer. Anal.* **54**, 2036–2059 (2016). <https://doi.org/10.1137/15M1032752>
19. Amodio, P., Brugnano, L., Iavernaro, F.: Spectrally accurate solutions of nonlinear fractional initial value problems. *AIP Conf. Proc.* **2116**, 140005 (2019). <https://doi.org/10.1063/1.5114132>
20. Brugnano, L., Montijano, J.I., Rández, L.: On the effectiveness of spectral methods for the numerical solution of multi-frequency highly-oscillatory Hamiltonian problems. *Numer. Algorithms* **81**, 345–376 (2019). <https://doi.org/10.1007/s11075-018-0552-9>
21. Brugnano, L., Iavernaro, F., Montijano, J.I., Rández, L.: Spectrally accurate space-time solution of Hamiltonian PDEs. *Numer. Algorithms* **81**, 1183–1202 (2019). <https://doi.org/10.1007/s11075-018-0586-z>
22. Amodio, P., Brugnano, L., Iavernaro, F.: Analysis of Spectral Hamiltonian Boundary Value Methods (SHBVMs) for the numerical solution of ODE problems. *Numer. Algorithms* **83**, 1489–1508 (2020). <https://doi.org/10.1007/s11075-019-00733-7>
23. Boyd, J.P.: *Chebyshev and Fourier Spectral Methods*, 2nd edn. Dover Publ. Inc., New York (2000)

24. Canuto, C., Hussaini, M.Y., Quarteroni, A., Zang, T.A.: Spectral Methods. Springer-Verlag, Berlin (2007)
25. Gheorghiu, C.I.: Spectral methods for non-standard eigenvalue problems. Fluid and structural mechanics and beyond. SpringerBriefs in Mathematics. Springer, Cham (2014)
26. Shen, J., Tang, T., Wang, L.-L.: Spectral methods. Algorithms, analysis and applications. Springer Series in Computational Mathematics, vol. 41. Springer, Heidelberg (2011)
27. Brugnano, L., Iavernaro, F.: Line Integral Solution of Differential Problems. *Axioms* **7**(2), 36 (2018). <https://doi.org/10.3390/axioms7020036>
28. Amodio, P., Brugnano, L., Iavernaro, F.: A note on the continuous-stage Runge-Kutta-(Nyström) formulation of Hamiltonian Boundary Value Methods (HBVMs). *Appl. Math. Comput.* **363**, 124634 (2019). <https://doi.org/10.1016/j.amc.2019.124634>
29. Amodio, P., Brugnano, L., Iavernaro, F.: Continuous-Stage Runge-Kutta approximation to Differential Problems. *Axioms* **11**, 192 (2022). <https://doi.org/10.3390/axioms11050192>
30. Brugnano, L., Iavernaro, F., Trigiante, D.: Analysis of Hamiltonian Boundary Value Methods (HBVMs): A class of energy-preserving Runge-Kutta methods for the numerical solution of polynomial Hamiltonian systems. *Commun. Nonlinear Sci. Numer. Simul.* **20**, 650–667 (2015). <https://doi.org/10.1016/j.cnsns.2014.05.030>
31. Brugnano, L., Iavernaro, F., Trigiante, D.: A note on the efficient implementation of Hamiltonian BVMs. *J. Comput. Appl. Math.* **236**, 375–383 (2011). <https://doi.org/10.1016/j.cam.2011.07.022>
32. Brugnano, L., Magherini, C.: Blended Implementation of Block Implicit Methods for ODEs. *Appl. Numer. Math.* **42**, 29–45 (2002). [https://doi.org/10.1016/S0168-9274\(01\)00140-4](https://doi.org/10.1016/S0168-9274(01)00140-4)
33. Brugnano, L., Magherini, C.: Recent advances in linear analysis of convergence for splittings for solving ODE problems. *Appl. Numer. Math.* **59**, 542–557 (2009). <https://doi.org/10.1016/j.apnum.2008.03.008>
34. Brugnano, L., Frasca-Caccia, G., Iavernaro, F.: Efficient implementation of Gauss collocation and Hamiltonian Boundary Value Methods. *Numer. Algorithms* **65**, 633–650 (2014). <https://doi.org/10.1007/s11075-014-9825-0>
35. Brugnano, L., Magherini, C.: Blended Implicit Methods for solving ODE and DAE problems, and their extension for second order problems. *J. Comput. Appl. Math.* **205**, 777–790 (2007). <https://doi.org/10.1016/j.cam.2006.02.057>
36. Bellen, A., Zennaro, M.: Numerical methods for Delay Differential Equations. Clarendon Press, Oxford (2003)
37. Brunner, H.: Collocation methods for Volterra integral and related functional equations. Cambridge University Press, Cambridge (2004)
38. Brugnano, L., Frasca-Caccia, G., Iavernaro, F., Vespi, V.: A new framework for polynomial approximation to differential equations. [arXiv:2106.01926](https://arxiv.org/abs/2106.01926) [math.NA]. doi:<https://doi.org/10.48550/arXiv.2106.01926>
39. Amodio, P., Brugnano, L., Iavernaro, F.: Energy-conserving methods for Hamiltonian Boundary Value Problems and applications in astrodynamics. *Adv. Comput. Math.* **41**, 881–905 (2015). <https://doi.org/10.1007/s10444-014-9390-z>
40. Brugnano, L., Magherini, C., Mugnai, F.: Blended Implicit Methods for the Numerical Solution of DAE Problems. *J. Comput. Appl. Math.* **189**, 34–50 (2006). <https://doi.org/10.1016/j.cam.2005.05.005>
41. Podlubny, I.: Fractional Differential Equations. Academic Press, New York (1999)
42. Garrappa, R.: Numerical solution of fractional differential equations: a survey and a software tutorial. *Mathematics* **6**(2), 16 (2018). <https://doi.org/10.3390/math6020016>
43. Zhang, Z., Zeng, F., Karniadakis, G.E.M.: Optimal error estimates of Petrov-Galerkin and collocation methods for initial value problems of fractional differential equations. *SIAM J. Numer. Anal.* **53**, 2074–2096 (2015). <https://doi.org/10.1137/140988218>
44. Brugnano, L., Montijano, J.I., Rández, L.: High-order energy-conserving Line Integral Methods for charged particle dynamics. *J. Comput. Phys.* **396**, 209–227 (2019). <https://doi.org/10.1016/j.jcp.2019.06.068>
45. Brugnano, L., Gurioli, G., Zhang, C.: Spectrally accurate energy-preserving methods for the numerical solution of the “Good” Boussinesq equation. *Numer. Methods Partial Differential Equations* **35**, 1343–1362 (2019). <https://doi.org/10.1002/num.22353>
46. Barletti, L., Brugnano, L., Tang, Y., Zhu, B.: Spectrally accurate space-time solution of Manakov systems. *J. Comput. Appl. Math.* **377**, 112918 (2020). <https://doi.org/10.1016/j.cam.2020.112918>

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.