



Plant-derived food waste management, valorization, and recycling through sourdough fermentation

Erica Pontonio^{a,*}, Alessandro Stringari^b, Raffaella Di Cagno^b, Pasquale Filannino^a, Carlo Giuseppe Rizzello^c, Andrea Polo^b, Olga Nikoloudaki^b, Marco Gobetti^b

^a Department of Soil, Plant, and Food Sciences, University of Bari Aldo Moro, Bari, Italy

^b Faculty of Science and Technology, Free University of Bolzano, Bolzano, Italy

^c Department of Environmental Biology, "Sapienza" University of Rome, Rome, Italy

ARTICLE INFO

Handling editor: AR Jambrak

Keywords:

Food waste
Sourdough fermentation
Lactic acid bacteria
Nutritional properties
Functional features

ABSTRACT

Background: The food industry generates a vast amount of food waste. Nevertheless, several types of food waste, i. e. those deriving from fruits, vegetables, grains, and other plant-based food production and processing chains, still contain valuable nutritional and bioactive compounds thus having the potential to be converted into value-added products. Several approaches have been investigated as pre-treatment of food waste to improve the nutritional, functional, and technological properties before to re-inclusion in food production. Sourdough fermentation, either spontaneous or through selected microbial strains, appears to be a suitable and sustainable tool for upcycling plant-derived food waste.

Scope and approach: This review reveals the latest insights into the potential of sourdough fermentation to recycle milling by-products, brewers' spent grain, wasted bread, and miscellaneous plant wastes.

Key findings and conclusions: Sourdough biotechnology is suitable for improving the sustainability of several food chains. Nevertheless, due to the significant effect of the presence, growth, and metabolic activity of specific microorganisms on the quality of the final products, an accurate set-up and optimization of tailored fermentation processes is highly suggested.

1. Introduction

Food waste is any edible or inedible loss from its supply chain (O'Connor et al., 2021). In the European Union, over 58 million tons of food waste (131 kg/inhabitant) are generated annually (Food waste and food waste prevention – estimates, 2023). The most abundant part comes from fruits and vegetables, including cereals as the main ingredients for worldwide staple foods, which generate by-products during pre- and post-harvesting, preparation, and processing (FAO, 2019). The overall estimation is that one-third of worldwide food is wasted at various stages along the food supply chain (FAO, 2019). FAO's Food Loss Index estimates that globally, around 14 percent of all food produced is lost from the post-harvest stage up to, but excluding, the retail stage (FAO, 2019). According to the UNEP Food Waste Index 2021, around 931 million tons of food waste were generated in 2019–61% of which came from households, 26% from food service, and 13% from retail – suggesting that 17% of global food production may be wasted at these stages of the food supply chain. Similarly, in the EU, households

generate more than half of the total food waste (54%) with 70% of food waste arising at household, food service and retail (Food waste and food waste prevention – estimates, 2023).

Stockpiling of food wastes poses social, environmental, and economic issues. Global food system causes approximately one-fourth of all greenhouse gas emissions (FAO, 2019). Food wastes require large-scale storage, transportation, and costly disposal applications (FAO, 2019). They have high oxidation potential and water content, and when generated from rendering plants and accidental failure of the cooling system occur, they also suffer from contamination by spoiling and/or pathogen agents (ICMSF, 2000). Given this impact, the United Nations set the Sustainable Development Goals to reduce food waste by 50% per capita by 2030. FAO believes that food waste recycling is the pillar of achieving sustainable development with the concept of a circular economy (FAO, 2019).

Solutions for food waste concern education programs focused on consumers' healthy lifestyles, prevention, improvement of processing efficiency, and valorization. Valorization is the most sustainable

* Corresponding author.

E-mail address: erica.pontonio@uniba.it (E. Pontonio).

<https://doi.org/10.1016/j.tifs.2024.104589>

Received 15 April 2024; Received in revised form 3 June 2024; Accepted 9 June 2024

Available online 10 June 2024

0924-2244/© 2024 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

approach, which should convert wastes into products with added value, thus limiting their disposal in landfills and eliminating greenhouse gas emissions (Al-Obadi et al., 2022). Currently, food wastes are physically, chemically, or biologically transformed to get an array of biofertilizers and soil amendments. Composting and anaerobic digestion are the main large-scale conversion techniques to exploit the potential of food wastes into biofertilizers and soil amendments. Other emerging conversions such as dehydration, biochar production and chemical hydrolysis show promising potentialities in agriculture and soil remediation (O'Connor et al., 2021). Aiming at a “zero waste economy”, food waste can be used as raw material for new products and applications. Nevertheless, it was observed that the food waste needs further processing steps before being used. Overall, the use of microorganisms that guide the production of functional ingredients for novel food formulation has been suggested to be helpful for the development of low-cost bioprocesses of food wasted and by-products leading to a transition toward a bio-economy model (Sabater et al., 2020). Lactic acid bacteria isolated from sourdough can be employed as a biotechnological starter to increase the safety of food industry by-products, to provide added value, to design the synthesis of functional molecules in fermentable substrates, and to moderate the technologies for safer alternative stock (e.g., food waste and by-products) incorporation to the main food (e.g., bread) formulas. Sourdough is a mixture of flour and water, spontaneously fermented by lactic acid bacteria and yeasts, and having acidification and leavening capacities (Arora et al., 2021). Sourdough is one of the oldest examples of natural starters, mostly used for making leavened baked goods as an alternative to baker's yeast and chemical leavening. Almost thirty years of literature accumulated on sourdough show the undoubtedly technological, sensory, and nutritional advantages compared to the other

leavening agents. A recent systematic review (Arora et al., 2021) highlighted the versatility of sourdoughs to ferment an extraordinary variety of cereal, pseudo-cereal, and legume flours, as well as miscellaneous agri-food by-products.

Here, we review the potential of sourdough fermentation to recycle milling by-products, brewers' spent grain, wasted bread, and miscellaneous plant wastes. Sourdough fermentation should represent a natural, low-cost, sustainable, and flexible approach to create added value within the food chain system (Fig. 1).

1.1. Recycling of milling by-products

Milling is the principal conversion procedure in the cereal industry. Dry milling separates by-products such as outer fibrous materials and germs from the grain endosperm. Pearling gradually removes seed coat (testa and pericarp), aleurone and sub-aleurone layers, and germ to get polished grains. Wet milling delivers starch and gluten, leaving steep solids, germ, and bran as by-products. It is estimated that *circa* 13% of all food waste comes from these milling procedures, with 30% of the cereal weight basis lost (FAO, 2019). While milling is inevitable for processing baked goods, most, if not all, milling by-products have high biological and nutritional value to be recycled (Cacace et al., 2022). Chemical and/or physical extraction, and purification procedures are options to separate valuable compounds, but they are quite expensive and contribute *per se* to environmental pollution. On the other hand, the direct reuse of untreated milling by-products as food ingredients causes poor technological and unpleasant sensory attributes. Sourdough fermentation, either spontaneous or driven by an *ad hoc* microbiota (Arora et al., 2021), conjugates the potential to exploit the biological




Fig. 1. Schematic approach of food waste fermentation.

Table 1

Non-exhaustive list of the main advantages related to the application of spontaneous or selected sourdough fermentation to cereal milling by-products and their related products.

Cereal source	Fermentation type/species/strain	Effects	References
Bran			
Wheat	<i>Levilactobacillus brevis</i> E95612 and <i>Kazachstania exigua</i> C81116 with enzymes	Increase of peptides and free amino acids concentration; increase of protein digestibility, soluble fiber concentration. Decrease of pungent flavor and bitter taste of fortified bread.	Coda et al. (2014)
	<i>Lactocaseibacillus rhamnosus</i> (R0011, ATCC 9595, and RW-9595M) and <i>Saccharomyces cerevisiae Lactocaseibacillus rhamnosus</i> 1473	Exopolysaccharides synthesis; increase of total phenolic content, bioaccessibility and antioxidant activity.	Bertsch et al. (2020)
	<i>Enterococcus faecalis</i> M2	Decrease of phytic acid; increase of soluble arabinoxylans; improvement of aroma profile.	Spaggiari, Ricci, et al. (2020)
	Spontaneous fermentation	Increase of soluble dietary fiber, alkylresorcinol, flavonoid and total phenol contents, and antioxidant activity	Mao et al. (2020)
	Commercial baker's yeast, <i>Lactobacillus delbrueckii</i> subsp. <i>bulgaricus</i> and <i>Streptococcus thermophilus</i>	Increase of soluble fiber water-extractable arabinoxylans and free ferulic; total degradation of phytic acid.	Manini et al. (2014)
	<i>Levilactobacillus brevis</i> E-95612 and <i>Candida humilis</i> E-96250 with enzymes	Increase of water extractable arabinoxylans and soluble dietary fiber.	Zhao et al. (2017)
	<i>Kazachstania exigua</i> VTT C-81116 and <i>Levilactobacillus brevis</i> VTT E-95612	Reduction of phytic acid content. Improvement of hydration properties and flavor.	Arte et al. (2015)
	Spontaneous fermentation	Increase of protein solubilization and <i>in-vitro</i> digestibility. Increase of phytase activity and total phenols content.	Savolainen et al. (2014)
	<i>Lactobacillus helveticus</i> FAM22155	Increase of free amino acids and phenolic compounds concentration.	Katina et al. (2012)
	<i>Companilactobacillus paralimentarius</i> (PB 3, PB 4), <i>Lactobacillus helveticus</i> (PB 7), <i>Levilactobacillus brevis</i> (PB 12A)	Reduction of aflatoxin B1	Zhang et al. (2021)
	<i>Liquorilactobacillus uvarum</i> , <i>Lactocaseibacillus casei</i> and <i>Lactocaseibacillus paracasei</i>	Increase of radical scavenging activity of wheat-rye bread	Mikušová et al. (2013)
	<i>Pediococcus acidilactici</i> LUHS29 (DSM 20284)	Lowering content of mycotoxin (below the threshold for human consumption) and biogenic amine.	Bartkiene, Zokaityte, et al., 2021; Zokaityte et al., 2021
	<i>Lactobacillus helveticus</i>	Reduction in DON content in the sourdough by 44–69% and a removal of 15-acetyldeoxyvalenol (15-AcDON), alternariol (AOH), deoxynivalenol-3-glucoside (D3G), toxins H-2 and HT-2.	Zadeike et al. (2021)
Wheat, barley and emmer	<i>Lactobacillus plantarum</i> T6B10 and <i>Weissella confusa</i> BAN8 with commercial xylanase	Decrease of aflatoxin B1 concentration.	Zhang et al. (2021)
	<i>Streptococcus thermophilus</i> , <i>Lactocaseibacillus rhamnosus</i> , <i>Saccharomyces cerevisiae</i> and <i>Candida milleri</i>	Increase of peptides and total free amino acids; Increase of total phenols and antioxidant activity. Decrease of phytic acid. Improvement of the nutritional value of bread.	Pontonio, Dingeo, et al. (2020)
Oat	<i>Streptococcus thermophilus</i> , <i>Lactocaseibacillus rhamnosus</i> , <i>Saccharomyces cerevisiae</i> and <i>Candida milleri</i>	Folic acid fortification.	Korhola et al. (2014)
	<i>Kazachstnia humilis</i>	Increase of fiber solubility.	Degutyte-Fomins et al. (2002)
Rice	<i>Lactocaseibacillus rhamnosus</i> and <i>S. cerevisiae</i> with enzymes	Reduction of the cytotoxicity and inhibition of melanogenesis in B16F1 melanoma.	Chung et al. (2009)
	<i>Lactobacillus acidophilus</i> GIM1.731 and <i>Lactiplantibacillus plantarum</i> subsp. <i>plantarum</i> GIM 1.648 with enzymes	Increase of total phenolic content and antioxidant activity.	Liu, Cao, et al. (2017)
	<i>Weissella koreensis</i> DB1	Increase of ornithine and citrulline content.	Yeong et al. (2020)
	<i>Lactiplantibacillus plantarum</i> EM	Increases in total amino acid content and antioxidant activity of bread.	Moon and Chang (2021)
	<i>Levilactobacillus brevis</i>	High cholesterol removal and strong antimicrobial activity.	Sadeghi et al. (2019)
	<i>Levilactobacillus brevis</i>	Decrease of mycotoxin concentration. Anti-aflatoxigenic effect on aflatoxin B1.	
Rye	Baker's yeast with enzymes	Increase of phenolic compounds in bread.	Koistinen et al. (2016)
	<i>Limosilactobacillus reuteri</i> TMW 1.106	Increase of amount and speed production of exopolysaccharides.	Kaditzky and Vogel (2008)
	<i>Weissella confusa</i>	Exopolysaccharides synthesis.	Kajala et al. (2016)
Germ			
Wheat	Spontaneous fermentation	Improve technological and sensory properties of doughs and breads.	Marti et al. (2014)
	<i>Lactiplantibacillus plantarum</i> subsp. <i>plantarum</i> DSM 32248 and <i>Furfurilactobacillus rossiae</i> DSM 32249	Slow down lipid oxidation by decreasing the activity of lipase and lipoxygenase. Increase of <i>in-vitro</i> protein digestibility, the concentration of total phenols and amino acids, phytase and antioxidant activities (sourdough and bread). Improve the texture and sensory properties of bread.	Rizzello et al., 2010a,b
	<i>Lactiplantibacillus plantarum</i> and <i>Lactobacillus acidophilus</i>	<i>Ex-vivo</i> anti-proliferative effects colon and ovarian carcinoma cell lines.	Rizzello et al. (2013)
	<i>Saccharomyces cerevisiae</i> 5022 and <i>Lactiplantibacillus plantarum</i> 299v	Decrease of the activity of lipase and lipoxygenase and increase in antioxidant activity.	Khosroshahi et al. (2022)
	<i>Lactiplantibacillus plantarum dy-1</i>	Increase of peptide and GABA concentrations and scavenging activity.	Bayat et al. (2022)
	<i>Latilactobacillus sakei</i> TMW1.22 and <i>Fruutilactobacillus sanfranciscensis</i> DSM20451T	Antiproliferative effects and the induction of apoptosis of human HT-29 colon cancer cells.	Zhang et al. (2015)
	<i>Saccharomyces cerevisiae</i>	Degradation of wheat germ agglutinin.	Tovar & Gänzle, 2021
	Spontaneous fermentation	Cancer-fighting characteristics: inhibition metastatic tumor dissemination and proliferation.	Boros et al., 2005; Saiko et al., 2009; Comín-Anduix et al., 2002
Rice	<i>Latilactobacillus sakei</i>	Anti-aging activity.	Zhao et al. (2021)
		Enrich in GABA showing a positive effect on the sleep disturbance of mice.	Mabunga et al. (2015)
Bran and Germ			

(continued on next page)

Table 1 (continued)

Cereal source	Fermentation type/species/strain	Effects	References
Wheat	<i>Lactiplantibacillus plantarum</i> subsp. <i>plantarum</i> DSM 32248 and <i>Furfurilactobacillus rossiae</i> DSM 32249	Increase of fiber content, protein digestibility and nutritional indexes of fortified bread. Decrease of glycemic index.	Pontonio et al. (2017)
Maize	<i>Lactiplantibacillus plantarum</i> subsp. <i>plantarum</i> T6B10 and <i>Weissella confusa</i> BAN8	Increase of the concentrations of free amino acids and peptides, the antioxidant activity. Degradation of phytate. Improve content of dietary fibers and proteins, the protein digestibility, and the starch hydrolysis index in bread	Pontonio et al. (2019)
	<i>Lactobacillus sakei</i> MI401 and <i>Pediococcus acidilactici</i> PA-2	Improve technological properties of albumins and globulins and increase the digestibility and free radical scavenging activity of prolamins. Increase of free amino acids concentration and antioxidant activity.	Zadeike et al. (2022)
Rye/wheat	<i>Kazachstania unispora</i> and <i>Kazachstania servazii</i> and <i>Latilactobacillus curvatus</i>	Improve the complexity of the volatile molecules. Increase in short chain fatty acids, antioxidant activity, total phenol content, bioactive peptides. Decrease in phytic acid content and an increase in prebiotic activity.	Siroli et al. (2022)

value and eliminate the negative attributes of recycled bran and germ in food preparations (Table 1).

1.1.1. Cereal bran

With variations depending on the cereal species, bran is a concentrate of dietary fibers (mainly arabinoxylans), phenolic compounds, proteins, and amino acids with high biological value, as well as minerals and vitamins (e.g., folates) (Chen et al., 2023). Unfortunately, these nutritive features has low bio-accessibility for humans, limiting its recycling potential. The bound forms of dietary fibers and phenolic compounds weaken their bioavailability, digestibility and/or intestinal absorption (Holland et al., 2020). The formation of insoluble complexes with phytate (Saurabh et al., 2021) and constraints imposed by the cell wall matrix limit the digestion and bioavailability of proteins (Alzuwaid et al., 2020) and minerals. Furthermore, the direct recycling of bran in food preparations might be responsible for poor hygiene (mycotoxins and other contaminants), and unpleasant rheology and sensory attributes (Ma et al., 2021). Notwithstanding the potential of other microorganisms (Chen et al., 2021; Li et al., 2022; Wu et al., 2022), sourdough fermentation prior to recycling into food preparations, alone or in combination with enzymes, has the potential to overcome most of these nutritional, technological, and sensory constraints (Chen et al., 2021; Li et al., 2022; Wang, Li, et al., 2022a,b).

The *ad hoc* fermentation with two typical sourdough species, *Levilactobacillus brevis*, and *Kazachstania exigua*, in combination with xylanase, endoglucanase, and β -glucanase enzymes, was successful in breaking the wheat bran cell wall, thus increasing the content of soluble arabinoxylans (Coda et al., 2014). Combining the spontaneous sourdough fermentation and the activity of endogenous or microbial enzymes, the same effect was found for rye arabinoxylans. After fermentation with sourdough, the bran was recycled into the bread formula, and volume and texture improved (Courtin & Delcour, 2002). Usually, the extractability of arabinoxylans increased during spontaneous sourdough backslipping and reached the highest value when the microbiota of lactic acid bacteria and yeasts became mature and stable (Manini et al., 2014). Soluble dietary fibers almost quadrupled, free amino acids increased as the hydrolysis of wheat bran proteins proceeded, and the content of phytate decreased (Mao et al., 2020). Almost the same efficiency was observed under solid-state fermentation of wheat bran steered by *Lactobacillus delbrueckii* subsp. *bulgaricus* and *Streptococcus thermophilus* together with baker's yeast (Zhao et al., 2017). Mixing wheat bran and whey permeate, another agri-food by-product, the fermentation with *Lacticaseibacillus rhamnosus* and *Saccharomyces cerevisiae* allowed the solubilization of fibers and increased the level of free phenolic compounds (Bertsch et al., 2020).

The sourdough fermentation does not affect the total content of phenolic compounds but elevates the ratio between free and bounded forms. The sourdough fermentation of both peeled and native rye bran increased the content of free ferulic acid and other hydroxycinnamic acids liberated from the polymeric structure (Katina et al., 2007). It was estimated that the spontaneous sourdough fermentation increased the

content of bioavailable ferulic acid by 82% (Manini et al., 2014). Lactic acid bacteria, including those populating the sourdough ecosystem, express ferulic acid esterases (Gaur & Gänzle, 2023). Additionally, *Pediococcus acidilactici* was found to significantly increase the contents of caffeic acid and mainly gallic acid during fermentation (Zhang et al., 2023).

When combined with cell wall degrading enzymes, sourdough species such as *Lv. brevis* and *Kazachstania humilis* abundantly released free phenolic compounds (Arte et al., 2015). *Lac. rhamnosus* metabolized conjugated phenolic compounds and broke the linkage with cell wall polysaccharides (Spaggiari, Calani, et al., 2020). The wheat bran solid-state fermentation that started with *Enterococcus faecalis* promoted a remarkable free radical scavenging activity, which depended on the release of free phenolic compounds (Mao et al., 2020). The sourdough fermentation with *Lactobacillus acidophilus*, *Lactiplantibacillus plantarum* and hydrolyzing enzymes modified the profile of phenolic compounds in rice bran, thus increasing the levels of soluble ferulic acid (Liu, Zhang, et al., 2017). This processed rice bran was used as an ingredient for making functional foods, which, due to the enrichment of free phenolic acids and flavonoids, showed remarkable antioxidant activity.

Although the disruption of the cell wall matrix is the main factor for increasing the bio-accessibility of bran proteins (Li et al., 2023), sourdough fermentation initially solubilized wheat bran proteins because of the activation of endogenous proteases at circa pH 4.0 (Arora et al., 2021). Following this primary proteolysis, peptidases by sourdough lactic acid bacteria were responsible for the liberation of small-size peptides and free amino acids (Christensen et al., 2022), which were more readily absorbable at intestinal level comparing to native proteins (Khubber et al., 2022). Processing rye bran with enzymes and sourdough fermentation prior to recycling into bread making led to high *in-vitro* protein digestibility (Nordlund et al., 2013). Sourdough proteolysis also favored the release of biogenic peptides (Pontonio, Verni, et al., 2020; Verni, Verardo, & Rizzello, 2019) and degraded antinutritional factors such as trypsin inhibitors and phytic acid (Patterson et al., 2017; Pontonio, Dingo, et al., 2020). Pre-fermented wheat bran was used to enhance the nutritional profile of bread. Compared to nonfermented bran, the *in-vitro* protein digestibility increased by 40%, and enriched breads showed pleasant sensory and textural attributes (Pontonio, Verni, et al., 2020). The sourdough fermentation with an *ad hoc* microbiota was also efficient for the treatment of rice bran. *Weissella koreensis* enriched rice bran with ornithine, which is a healthcare supplement to activate the immune system and liver function (Yeong et al., 2020). Water-soluble extracts of sourdough fermented rice bran had an anti-photoaging effect on human skin fibroblasts cultures (Seo et al., 2010). The rice bran co-fermented with *Lac. rhamnosus* and *S. cerevisiae* decreased the cytotoxicity and inhibited melanogenesis in B16F1 melanoma through the downregulation of microphthalmia-associated transcription factors (Chung et al., 2009). Rice bran fermented with *Lp. plantarum* attenuated the levels of cholesterol (45–68%) and exerted inhibitory activities towards foodborne pathogenic bacteria and spoiling fungi (Moon & Chang, 2021).

Cereal endogenous or microbial phytases are responsible for the hydrolysis of phytic acid in cereal bran. Wheat endogenous phytases are almost inactive at neutral pH, but sourdough acidification promotes their activation with the consequent decrease of the phytate content and the increase of protein and mineral bioavailability (Ameur et al., 2022). Phytases by several sourdough lactic acid bacteria and yeasts were characterized. *Lac. rhamnosus* had the capability to degrade phytate in wheat bran (Spaggiari, Ricci, et al., 2020), and *Lp. plantarum* and *Weissella confusa* decreased its level by 25%–60% when used as starters for wheat bran fermentation (Moon & Chang, 2021; Pontonio, Dingeo, et al., 2020). Also, the fermentation of rice bran with a functional strain of *Lp. plantarum* EM decreased the content of phytic acid by 50% (Moon & Chang, 2021). The spontaneous sourdough fermentation of bran increased the level of folates by over 100% and the highest detectable levels coincided with the highest cell density of autochthonous lactic acid bacteria and yeasts. Overall, the increase of folate was mainly due to synthesis by yeasts, while the liberation by lactic acid bacteria was strain-dependent. The *ad hoc* substitution of folate-consuming lactic acid bacteria with folate-synthesizing strains significantly increased the vitamin content in sourdough breads enriched with pre-fermented and recycled bran (Mendez-Vilas, 2007). The capability of yeasts to synthesize folate from oat bran was species dependent, with *S. cerevisiae*, *Pseudozyma* sp., *Rhodotorula glutinis* and *Kluyveromyces marxianus* leading to the highest concentrations (Korhola et al., 2014). Intending to increase the folate concentration in bran processing, sourdough fermentation with yeasts was always the most efficient option. Oat bran fermented with *St. thermophilus* or *Lac. rhamnosus*, and *S. cerevisiae* and *K. humilis* led to a folate concentration of 120 ng/g. The intake of 100 g of fermented oat bran supplied 15% of the recommended daily dose of folate (Korhola et al., 2014). The enrichment of rye and wheat brans with exopolysaccharides (EPS) is another option with multiple outcomes. Sourdoughs made with *ad hoc* lactic acid bacteria, for instance *Limosilactobacillus reuteri* (Kaditzky & Vogel, 2008) or *W. confusa*, was the most efficient choice. Usually, the *in-situ* production of EPS in rye bran reached the concentration of 2–3% on dry matter (Kajala et al., 2016) and represented the way to improve the structure-forming capability of bran, making it a recyclable ingredient with prebiotic and *in-vitro* antitumor and immunomodulating activities (Korcz & Varga, 2021).

1.1.2. Wheat germ

The annual world deposit of wheat germ is estimated to be circa 25,000,000 tons, which undoubtedly deserves valorization as it corresponds to the most nutritious part of the grain (Rizzello, Nionelli, Coda, De Angelis, & Gobbetti, 2010). The fast development of rancidity because of the presence of unsaturated fats, lipoxygenases and lipases severely limits the use of germ in baked goods processing (Li et al., 2016). Nowadays, sourdough fermentation is the most efficient solution to stabilize and improve the nutritional attributes of germ (Khosroshahi et al., 2022).

Lowering the value of pH by sourdough fermentation inactivates lipase and lipoxygenase enzymes (Marti et al., 2014; Rizzello, Nionelli, Coda, Di Cagno, & Gobbetti, 2010). Compared to nonfermented wheat germ, the sourdough fermentation with *Lp. plantarum* and *Furfurilactobacillus rossiae* almost inhibited the synthesis of volatile compounds from lipid oxidation during 40 days of storage (Rizzello, Nionelli, Coda, Di Cagno, & Gobbetti, 2010). Other sourdough starters were proven to behave similarly (Khosroshahi et al., 2022). In summary, the sourdough fermentation abolished the technological obstacles to incorporating wheat germ into the bread formula. Improved *in-vitro* protein digestibility, decreased phytase activity, and extended shelf life were the main nutritional benefits derived from the germ supplementation into wheat bread formula (Rizzello, Nionelli, Coda, Di Cagno, & Gobbetti, 2010). The sourdough fermentation also decreased the content of anti-nutritional factors such as phytic acid and raffinose and raised the content of free amino acids to 50%, mainly lysine (Lys) and γ -amino

butyric acid (GABA), a non-protein amino acid with functional features (Bayat et al., 2022; Rizzello, Nionelli, Coda, De Angelis, & Gobbetti, 2010; Rizzello, Nionelli, Coda, Di Cagno, & Gobbetti, 2010). One of the most promising achievements for sourdough fermented wheat germ was represented by the cytotoxic activity towards cancer cell lines. Wheat germ fermented with *Lp. plantarum* dy-1 inhibited the proliferation of HT-29 cells via apoptosis suggesting its use as a potential anticarcinogenic (Zhang et al., 2015). Avemar®, a commercial product made of wheat germ fermented with *S. cerevisiae* (Boros et al., 2005; Saiko et al., 2009), showed *in-vitro* anticancer and autoimmune properties on various human cancer cell lines (Comín-Anduix et al., 2002). Two-methoxy benzoquinone and 2,6-dimethoxybenzoquinone are naturally present in wheat germ but in a glycosylated and non-active form. After activation through β -glucosidase activity of *Lp. plantarum* and *Fu. rossiae*, they exerted *ex-vivo* anti-proliferative effects on colon and ovarian carcinoma cell lines (Rizzello et al., 2013). Based on the metabolism of thiols, sourdough fermentation decreased the content of agglutinins in wheat germ, which is one of the triggering factors for non-celiac wheat sensitivity (Tovar & Gänzle, 2021). A sourdough made combining wheat germ and bran was recycled and used in the bread formula (Pontonio et al., 2017). The bread was enriched in dietary fibers, free amino acids, and phenolic compounds, and as shown by *in-vitro* and *in-vivo* assays, it was classified as a low glycemic index food. After fermentation with *Lactilactobacillus sakei*, the rice germ fermented was enriched in GABA, showing a positive effect on the sleep disturbance of mice (Mabunga et al., 2015). *Lp. plantarum* and *W. confusa* were used to ferment raw maize or heat-treated germ and bran (Pontonio et al., 2019). The concentrations of free amino acids and peptides, the antioxidant activity, and the phytate degradation increased during sourdough fermentation. Incorporation of these fermented maize by-products in bread positively affected the content of dietary fibers and proteins, the protein digestibility, and the starch hydrolysis index (Pontonio et al., 2019).

1.1.3. Mycotoxins degradation

Compared to whole grains, by-products from cereal milling suffer from higher levels of mycotoxin contamination (Hoffmans et al., 2022). Although using implemented good practices, the bio-accumulation of mycotoxins in milling by-products is almost inevitable, especially under suboptimal storage conditions (Khodaei et al., 2021). Decontamination before recycling is a priority. Available options include physical, chemical, and biological methods, with the latter attracting considerable interest (Piotrowska, 2021). Although the effect of sourdough fermentation is somewhat controversial, the degradation of aflatoxins, fumonisins, ochratoxins, deoxynivalenol, and zearalenone was demonstrated with various species of lactic acid bacteria (Bartkiene, Zokaityte, et al., 2021; Muhialdin et al., 2020; Sadeghi et al., 2019; Zokaityte et al., 2021). The mechanism for mycotoxin decontamination occurs through the adhesion of significant amounts of mycotoxins to microbial cells and/or the degradation of mycotoxins into non-toxic metabolites (Muhialdin et al., 2020; Piotrowska, 2021).

Extrusion combined with the fermentation by *Lactocaseibacillus casei* and *Lactocaseibacillus paracasei*, or *Liquorilactobacillus uvarum* reduced the levels of several mycotoxins in wheat bran to levels below the threshold for human consumption (Bartkiene, Zokaityte, et al., 2021; Zokaityte et al., 2021). In addition, the sourdough fermentation with *Lp. plantarum* and *P. acidilactici* alone decreased the mycotoxin contamination of whole wheat flour and rye by 10%–40% (Pontonio et al., 2021). *Lv. brevis* was capable of decreasing the contamination by aflatoxins B1, B2, G1, and G2 (Sadeghi et al., 2019), and *P. acidilactici* decontaminated deoxynivalenol (44–69%), and removed 15-acetyldeoxynivalenol, alternariol, deoxynivalenol-3-glucoside, and toxins H-2 and HT-2 (Zadeike et al., 2021). Usually, the highest level of decontamination was observed during long-time (48 h) sourdough fermentation with mixed sourdough starters (Zadeike et al., 2021). Under solid-state fermentation of wheat bran, *Lactobacillus helveticus* synthesized




Fig. 2. Main advantages of fermentation on brewers' spent grains and wasted bread.

enzymes capable of degrading aflatoxin B1 into four low-toxin derivatives without a lactone ring structure (Zhang et al., 2021).

1.2. Brewers' spent grain

Malting for making beer leaves brewers' spent grains (BSG), whose elevated organic load is critical for environmental pollution (Teixeira et al., 2020). The recycling of BSG has gained attention due to biogenic components such as alkaloids, plant growth factors, food-grade pigments, and phenolic compounds (Verni, Pontonio, et al., 2020). BSG is also rich in cellulose and arabinoxylans and contains proteins up to 20%, particularly rich in lysine (Verni, Pontonio, et al., 2020). The high moisture content (circa 80%), along with the richness in polysaccharides and proteins, make BSG highly prone to microbial contamination with a short lifespan (7–10 days) (Kavalopoulos et al., 2021). Various conservative treatments of BSG have been assessed (Lynch et al., 2016), and among these, sourdough fermentation was one of the most promising (Fig. 2).

The sourdough fermentation with lactic acid bacteria enriched BSG with dextran and oligosaccharides (Koirala et al., 2021). The supplementation with sucrose (4% w/w) and fermentation with *Leuconostoc pseudomesenteroides* and *W. confusa* yielded 11 g/kg of dextran, which improved the BSG viscosity and taste. This fermented BSG was used as

an ingredient for wheat bread formula. Compared to control bread without addition, the bread enriched with BSG had a higher content of proteins and fibers, and positively influenced the gut microbiota functionality. Along with an increase of several essential amino acids, a consistent increase of GABA was also observed during the simulated digestion (Koirala et al., 2022). BSG was also used for making a beverage fermented by *Lp. plantarum*. The beverage had antioxidant potential because of the high content of phenols and flavonoids, which were released through acidification and microbial metabolism (Gupta et al., 2013). The combination of xylanase and sourdough fermentation with *Lp. plantarum* released free phenolic compounds and liberated encrypted biogenic peptides with antioxidant activity and protective effect toward oxidative stress on human keratinocytes (Verni, Pontonio, et al., 2020). Given these attributes, the bioprocessed BSG was used for making pasta. Compared to pasta containing nonfermented BSG, the enriched pasta had higher protective effects against oxidative stress caused by human colon carcinoma cells under simulated gastro-intestinal digestion (Schettino et al., 2021). Positive effects of fermented BSG were also observed when used as an ingredient in bread formula. The rheology and sensory attributes of the enriched bread improved, and the nutritional profile was optimal because of the supplementation with proteins, fibers, and essential amino acids (Waters et al., 2012). The enrichment of bread with spontaneously sourdough-fermented BSG also decreased the

phytate content (Ktenioudaki et al., 2015).

BSG was also used as a solid-state substrate to cultivate and preserve (up to 10 weeks at circa 6 Log cfu/g) sourdough cultures. The fermented BSG was used (10% p/p) in an experimental baking trial leading to sourdough breads characterized by optimal organic acids ratio and acceptability according to a consumer-based study (Vriesekoop et al., 2021).

1.3. Wasted bread

On a global scale, bread is one of the major wasted foods. Industrial waste is generated because of substandard products, crust removal for sandwich production, and unsold bread at retail (Verni, Rizzello, & Coda, 2019). In most cases, the huge amount of wasted bread is still under safe conditions for human consumption. Indeed, innovative, and sustainable recycling solutions must focus on keeping wasted bread - within the food chain. Recycling wasted bread through sourdough fermentation has shown an undoubted potential.

Recycling of wasted bread to prepare a culture medium (wasted bread medium, WBM) for the cultivation of lactic acid bacteria was considered (Verni, Minisci, et al., 2020). The protocol included bread (20%) homogenization with water and hydrolysis with amylases. Its suitability as a valid alternative to common synthetic media was demonstrated for almost all sourdough lactic acid bacteria reaching the most common cell densities after cultivation. The sourdough was also proposed to ferment wasted bread and recycle it as an ingredient for bread making. Sourdough was made from wasted bread crumbs. Several lactic acid bacteria demonstrated the capability of using the bread crumb as a fermentation substrate. *Lp. plantarum* and *P. acidilactici* were capable of a long-time (48–96 h) fermentation without the genesis of off-flavors (Gélinas et al., 1999). A sourdough made with wasted bread was used as an aroma and flavor enhancer for making bread. The formula included 50% whole wheat bread crumb and fermentation with *Lp. plantarum*. The synthesis of lactic and acetic acids resembled that of the traditional sourdough fermentation, and a positive effect on glycemic index and insulinemic responses was hypothesized (Poutanen et al., 2009). Bread is mainly subjected to fungal spoilage during its shelf life. Salts of propionic and sorbic acids and ethanol are the most common chemical preservatives to extend the shelf life, although their negative consumer perception and the healthy recommendations to markedly decrease their use. A protocol to synthesize bioactive peptides from wasted bread was standardized (Nionelli et al., 2020). After hydrolysis by microbial proteases and fermentation with *Lv. brevis*, the wasted bread showed a broad inhibitory spectrum against some of the most diffuse fungal-contaminating species. Nine antifungal peptides, encrypted in wheat protein sequences, were released during sourdough fermentation and identified as the main responsible for fungal inhibition. The shelf-life of the bread was extended up to 10 days (Nionelli et al., 2020). The direct recycling of wasted bread negatively affects the specific volume and softness of the newly manufactured bread. Gelatinized starch and denatured proteins are not reassembled into a new gluten network (Immonen et al., 2020). The hydrolysis of gelatinized starch improved the viscoelastic properties of wasted bread (Immonen et al., 2021), but the most suitable option is the use of sourdough lactic acid bacteria capable of synthesizing EPS (dextran and β -glucan) *in-situ* (Immonen et al., 2020). The dextran synthesis occurring during sourdough fermentation of wasted bread compensated for the adverse effect of the recycled bread increasing the specific volume and decreasing the crumb hardness. EPS synthesized by *Weissella* species also exerted antioxidant and prebiotic activities (Kavitate et al., 2020). A blend of wasted bread and wheat bran was used as the substrate to synthesize GABA (Verni et al., 2022). *Lp. plantarum* yielded the highest content of GABA (circa 800 mg/kg). This fermented wasted bread/bran slurry was used as an ingredient for the manufacture of GABA-enriched bread. In conclusion, tailored sourdough fermentations with selected lactic acid bacteria are key options for recycling wasted bread, gaining suitable

technological features, and creating added functional value along the food chain.

When wasted bread is not edible and recycling into the food chain implies microbiological risks (e.g., presence of mycotoxins), alternative biotechnological solutions are exploited. It is known that lactic acid bacteria act as plant growth-promoting microorganisms (PGPM), indirectly favoring nutrient acquisition in the alkaline soils characterizing the Mediterranean area. Besides, lactic acid bacteria act as biocontrol agents, improving the ability of the host plant to withstand biotic and abiotic stress or producing compounds that directly stimulate plant growth (Lamont et al., 2017). A bioprocessed wasted bread, which was obtained by enzyme treatments coupled with sourdough fermentation was used as an amendment in pot trials. The aim was to evaluate the modification of the soil physicochemical properties and the plant growth-promoting activity using escarole (*Cichorium endivia* var. Cuartana) as the indicator crop (Cacace et al., 2022). Compared to non-amended soils, the supplementation with sourdough fermented wasted bread raised the content of soil organic carbon up to 37% and total nitrogen up to 40%. In addition, the lower pH and the higher organic acid content favored a higher availability of Mn, Fe, and Cu. Escaroles showed improved growth with a higher number of leaves.

Recently, a biotechnological protocol for recycling bread waste at the industrial level has been set up and patented (Ampollini et al., 2023). The approach includes the preliminary drying and fine milling of bread resulting from sandwich production (bread crust). A slurry was obtained with waste bread powder and water, and food-grade proteases were added to the semiliquid formulation to promote fermentation. Lactic acid bacteria (*Lp. plantarum*) and yeast (*S. cerevisiae*) strains were selected for sourdough fermentation in automatic fermenters. It was demonstrated that waste bread sourdough can be added up to 50% of the final weight of bread dough without negatively affecting the textural properties of the bread. Sensory properties were moreover those typical of a sourdough bread.

1.4. Miscellaneous plant wastes

Miscellaneous plant by-products and wastes comprise a very heterogeneous set of matrices (e.g., fruit and vegetables hulling and peeling wastes, pomaces, food surplus) sharing an unquestionable nutritional value. Nevertheless, they are characterized by sources heterogeneity, variation of their chemical composition, and microbial instability. Being rich in carbohydrates, organic acids, mineral salts, and vitamins, these compounds can be utilized as substrates for contaminant (either pathogenic or not) microorganisms. Furthermore, the high fiber and phenolic contents can negatively affect the sensory profile and technological properties of fortified foods (Kosseva, 2020). The exploitation of their potential needs to overcome these weaknesses, either through technological treatments or exploiting fermentation processes, with the latter proven to be an effective, sustainable, and mild approach.

Apples are among the most widely processed fruits, which inevitably generate large amounts of residues. Such availability together with high contents of fibers and phytochemicals make these by-products and wastes ideal ingredients to enrich the wheat bread formula. Fermentation of apple by-products with a binary culture of *Weissella cibaria* and *S. cerevisiae* was proposed for recycling in breadmaking at a rate of 5–10% (w/w of flour) (Cantatore et al., 2019). The fruit and vegetable by-products addition in baked goods causes low acceptability because of the poor sensory and rheological attributes. The dilution of gluten with fruit or vegetable by-products implies a decreased capability of the dough to retain gas, which leads to low volume and high hardness. Fortification with fruit or vegetable by-products does not exceed 10% (w/w of flour) (Gómez & Martínez, 2018). The formula enriched with 5% apple by-products fermented with sourdough improved the wheat dough stability and water absorption and did not interfere with the bread specific volume and color (Cantatore et al., 2019). The sourdough fermentation maximized the hydration properties of apple by-products




Fig. 3. Schematic reconstruction of putative pathways characterizing the fermentation (30 °C for 24 h) of a brewer spent grain -based medium by *Leuconostoc pseudomesenteroides* and *Lactiplantibacillus plantarum* strains. Red colored pathways were unique to *Leuc. pseudomesenteroides*, while blue colored pathways were unique to *Lp. plantarum*. Black colored pathways were present in both species. Genes that were over-expressed compared to growth in standard reference medium (MRS broth) are shown in green (Acin-Albiac et al., 2022). (For interpretation of the references to color in this figure legend, the reader is referred to the Web version of this article.)

because of the synthesis of EPS, and the partial breakdown of insoluble fibers, which resulted in an increased porosity and surface area. Enrichment with sourdough-fermented apple by-products also provided bio-accessible polyphenols (Tlais et al., 2021). In addition, apple pomace was used to fortify sourdough started with a selected culture of *Fructilactobacillus florum* DSM 22689 and baker's yeast (single and co-culture). The sourdough nutritional value was positively affected by the incorporation mainly thanks to the presence of valuable compounds like organic acids and fibers. Additionally, the abundance of glucose and fructose in apple pomace led to higher microbial viability and growth (Martău et al., 2021). Research efforts also focused on the potential of plant by-products as sources of antimicrobial compounds to extend the bread shelf life. By-products having relevant protein contents represent ideal substrates for the biosynthesis of antimicrobial peptides. The sourdough fermentation of palm kernel cake with *Lc. casei* favored the accumulation of antifungal peptides, which, once added to the bread formula, were effective in counteracting the spoilage up for to 10 days (Asri et al., 2020). Another constraint to recycle non-conventional ingredients is their eventual association with increased levels of acrylamide during baking (Bartkiene, Bartkevics, et al., 2021). Oat has a higher potential for acrylamide formation than wheat because of the higher levels of soluble carbohydrates and free asparagine. Sourdough fermentation of oat by-products prevented acrylamide accumulation in baked goods (Bartkiene, Bartkevics, et al., 2021). Solid-state fermentation and the subsequent recycling as ingredients for breadmaking demonstrated how agri-food by-products also served to deliver

microbial enzymes (e.g., feruloyl esterase, xylanase) with appreciable properties (dos Santos Costa et al., 2021). Date seed flour was proposed as an innovative ingredient for sourdough breadmaking (Ameur et al., 2022). Autochthonous lactic acid bacteria and yeasts were isolated from date seeds and selected based on various technological criteria. A mixed starter, comprising *Leuconostoc mesenteroides*, *Lp. plantarum* and *S. cerevisiae*, was used to prepare type I sourdough after consecutive refreshments. An aliquot of durum wheat flour was replaced by date seed flour. The sourdough fermented bread showed an increased radical scavenging activity because of the consistent release of free phenolic compounds, while the perceived bitterness and astringency diminished due to microbial degradation of tannins. Being an economic and sustainable alternative to animal-based proteins, rapeseed proteins isolated from defatted rapeseed press cake have the potential for food applications. Rapeseed protein concentrations higher than 5% in wheat breadmaking led to detrimental effects on textural and sensory attributes. Rapeseed proteins became suitable ingredients for making sourdough bread when fermented with *W. confusa*, which synthesized dextran *in-situ*. Compared to control bread, the sourdough fermentation improved the *in-vitro* protein digestibility, enriched the free amino acid profile mainly with lysine, methionine, and isoleucine, and guaranteed suitable specific volume and low crumb hardness (Wang, Li, et al., 2022a,b).

1.4.1. Lactic acid bacteria glycosyl hydrolases

One of the main secrets beyond the sourdough fermentation as

valuable biotechnology to recycle plant by-products and wastes concerns the hydrolysis of complex polymers (e.g. arabinoxylans, cellulose, and hemicellulose fraction) and the bio-accessibility of phenolic compounds. Sourdough lactic acid bacteria have the aptitude to metabolize diversified carbon sources. This aptitude relies on key metabolic pathways mediated by an extensive set of glycosyl hydrolases. These enzymes undertake the ambivalent role of releasing a wide range of phenolic compounds from glycosylated precursors and degrading the resulting sugar moiety. Glycosyl hydrolases also have the potential to remove undesired bitterness or release aroma compounds. Encoding genes for glycosyl hydrolases are widespread among lactic acid bacteria. Phospho- β -glucosidase genes show a high degree of redundancy. Phospho- β -glucosidases catalyze the degradation of phosphorylated glucosides and fibers-related disaccharides (e.g., cellobiose and gentiobiose), which are activated through the phosphoenolpyruvate (PEP)-dependent carbohydrate phosphotransferase systems (PEP-PTS) (Acin-Albiac et al., 2021). Recently, the complete framework describing the metabolism drift of *Lp. plantarum* and *Leuc. pseudomesenteroides* caused by the lignocellulosic BSG was provided by implementing molecular and phenomics approaches (Fig. 3) (Acin-Albiac et al., 2021, 2022). As confirmed by gene overexpression, *Lp. plantarum* preferred arabinose among pentosans, while *Leuc. pseudomesenteroides* mainly uses xylose. The phenotype switching towards galactose metabolism suffered the greatest fluctuation in *Lp. plantarum*. All lactic acid bacteria strains utilized sucrose more intensively and its plant-derived isomers. Sucrose-6-phosphate hydrolases activity in *Leuc. pseudomesenteroides* likely mediated the increased consumption of raffinose. The increased levels of some phenolic compounds suggested the involvement of 6-phospho- β -glucosidases in β -glucosides degradation. Expression of genes encoding β -glucoside/cellobiose-specific EII complexes and phenotyping highlighted an increased metabolism for cellobiose (Acin-Albiac et al., 2022). The glycoside hydrolase group includes fructan- β -fructosidases and β -fructofuranosidases, which undertake the hydrolysis of inulin and fructooligosaccharides, and β -D-Xylosidases and α -L-arabinofuranosidases, which are responsible for xylooligosaccharides and arabino-xylooligosaccharides utilization by several lactobacilli (Petrova & Petrov, 2017).

2. Conclusion

Valorizing food waste, mainly using sustainable and biological approaches, i.e., either spontaneous or selected sourdough fermentation, is a tool for reducing its environmental and economic burden and transitioning to a circular economy. Fermentation with lactic acid bacteria and/or yeasts has been demonstrated to be an optimal approach to improve the nutritional and functional features (e.g., improved *in-vitro* protein digestibility, antioxidant activity) of several plant-derived food waste and by-products, and to overcome the drawbacks (e.g., decreased concentration of anti-nutritional compounds, improved textural properties) which might impede their re-use in food production. Aiming at facing the advancement of large-scale fermentations and the need for the implementation and diversification of available products on the market, the use of well-adapted and *ad hoc* selected microorganisms for industrial processing is of main importance. Also given the differences among the food waste composition, several studies have involved the selection of strains with various and specific functional properties to be used in tailored fermentation inspired to the sourdough biotechnology. The effects of this approach, suitable for significantly improving the sustainability of several food chains and the quality of the products, depend on the presence, growth, and metabolic activity of specific microorganisms, therefore requiring the accurate set-up and optimization of tailored fermentation processes.

Credit author statement

Erica Pontonio: Writing- Original draft preparation, Investigation,

Writing- Reviewing and Editing, Supervision. **Alessandro Stringari:** Data curation, Investigation. **Raffaella Di Cagno:** Writing- Original draft preparation, Investigation. **Pasquale Filannino:** Writing- Original draft preparation, Investigation. **Carlo Giuseppe Rizzello:** Data curation, Investigation, Writing- Original draft preparation. **Andrea Polo:** Original draft preparation. **Olga Nikoloudaki:** Data curation, Investigation. **Marco Gobetti:** Conceptualization, Methodology, Writing- Original draft preparation, Supervision.

Declaration of competing interest

The authors report there are no competing interests to declare.

Data availability

No data was used for the research described in the article.

Acknowledgements

The authors thank Lena Brigitte Marie Granehäll for language proofreading.

References

- Acin-Albiac, M., Filannino, P., Arora, K., Da Ros, A., Gobetti, M., & Di Cagno, R. (2021). Role of lactic acid bacteria phospho- β -glucosidases during the fermentation of cereal by-products. *Foods*, 10, 97. <https://doi.org/10.3390/foods10010097>
- Acin-Albiac, M., Filannino, P., Coda, R., Rizzello, C. G., Gobetti, M., & Di Cagno, R. (2022). How water-soluble saccharides drive the metabolism of lactic acid bacteria during fermentation of brewers' spent grain. *Microbial Biotechnology*, 15, 915–930. <https://doi.org/10.1111/1751-7915.13846>
- Al-Obadi, M., Ayad, H., Pokharel, S., & Arselene & Ayari, M. (2022). Perspectives on food waste management: Prevention and social innovations. *Sustainable Production and Consumption*, 31, 190–208. <https://doi.org/10.1016/j.spc.2022.02.012>
- Alzuwaid, N. T., Sissons, M., Laddomada, B., & Fellows, C. M. (2020). Nutritional and functional properties of durum wheat bran protein concentrate. *Cereal Chemistry*, 97 (2), 304–315. <https://doi.org/10.1002/cche.10246>
- Ameur, H., Cantatore, V., Filannino, P., Cavoski, I., Nikoloudaki, O., Gobetti, M., & Di Cagno, R. (2022). Date seeds flour used as value-added ingredient for wheat sourdough bread: An example of sustainable bio-recycling. *Frontiers in Microbiology*, 13, Article 873432. <https://doi.org/10.3389/fmicb.2022.873432>
- Ampollini, M., Di Cagno, R., Gobetti, M., Polom, A., Racinelli, F., Rizzello, C., & Stringari, A. (2023). *Bread waste sourdough*. Patent 2023/6042, filed . (Accessed 21 December 2023).
- Arora, K., Ameur, H., Polo, A., Di Cagno, R., Rizzello, C. G., & Gobetti, M. (2021). Thirty years of knowledge on sourdough fermentation: A systematic review. *Trends in Food Science & Technology*, 108, 71–83. <https://doi.org/10.1016/j.tifs.2020.12.008>
- Arte, E., Rizzello, C. G., Verni, M., Nordlund, E., Katina, K., & Coda, R. (2015). Impact of enzymatic and microbial bioprocessing on protein modification and nutritional properties of wheat bran. *Journal of Agricultural and Food Chemistry*, 63(39), 8685–8693. <https://doi.org/10.1021/acs.jafc.5b03495>
- Asri, N. M., Muhiyaldin, B. J., Zarei, M., & Saari, N. (2020). Low molecular weight peptides generated from palm kernel cake via solid state lacto-fermentation extend the shelf life of bread. *Lebensmittel-Wissenschaft & Technologie -LWT*, 134, Article 110206. <https://doi.org/10.1016/j.lwt.2020.110206>
- Bartkiene, E., Bartkevics, V., Pugajeva, I., Borisova, A., Zokaityte, E., Lele, V., Starkute, V., Zavistanaviciute, P., Klupsaite, D., Zadeike, D., & Juodeikiene, G. (2021). The quality of wheat bread with ultrasonicated and fermented by-products from plant drinks production. *Frontiers in Microbiology*, 12, Article 652548. <https://doi.org/10.3389/fmicb.2021.652548>
- Bartkiene, E., Zokaityte, E., Lele, V., Starkute, V., Zavistanaviciute, P., Klupsaite, D., Cernauskas, D., Ruzauskas, M., Bartkevics, V., Pugajeva, I., Berzina, Z., Gruzauskas, R., Sidlauskienė, S., Santini, A., & Juodeikiene, G. (2021). Combination of extrusion and fermentation with *Lactobacillus plantarum* and *L. uvarum* strains for improving the safety characteristics of wheat bran. *Toxins*, 13, 163. <https://doi.org/10.3390/toxins13020163>
- Bayat, E., Moosavi-Nasab, M., Fazaeli, M., Majdinasab, M., Mirzapour-Kouhdasht, A., & Garcia-Vaquero, M. (2022). Wheat germ fermentation with *Saccharomyces cerevisiae* and *Lactobacillus plantarum*: Process optimization for enhanced composition and antioxidant properties in vitro. *Foods*, 11, 1125. <https://doi.org/10.3390/foods11081125>
- Bertsch, A., Roy, D., & La Pointe, G. (2020). Fermentation of wheat bran and whey permeate by mono-cultures of *Lactocaseibacillus rhamnosus* strains and co-culture with yeast enhances bioactive properties. *Frontiers in Bioengineering and Biotechnology*, 8, 956. <https://doi.org/10.3389/fbioe.2020.00956>
- Boros, L. G., Nichelatti, M., & Shoenfeld, Y. (2005). Fermented wheat germ extract (Aveamar) in the treatment of cancer and autoimmune diseases. *Annals of the New York Academy of Sciences*, 1051, 529–542. <https://doi.org/10.1196/annals.1361.097>

- Cacace, C., Rizzello, C. G., Brunetti, G., Verni, M., & Coccozza, C. (2022). Reuse of wasted bread as soil amendment: Bioprocessing, effects on alkaline soil and escarole (*Cichorium endivia*) production. *Foods*, *11*, 189. <https://doi.org/10.3390/foods11020189>
- Cantatore, V., Filannino, P., Gambacorta, G., De Pasquale, I., Pan, S., Gobetti, M., & Di Cagno, R. (2019). Lactic Acid Fermentation to re-cycle apple by-products for wheat bread fortification. *Frontiers in Microbiology*, *10*, 2574. <https://doi.org/10.3389/fmicb.2019.02574>
- Chen, Z., Mense, A. L., Brewer, L. R., & Shi, Y. C. (2023). Wheat bran layers: Composition, structure, fractionation, and potential uses in foods. *Critical Reviews in Food Science and Nutrition*, *2*, 1–24. <https://doi.org/10.1080/10408398.2023.2171962>
- Chen, Q., Wang, R., Wang, Y., An, X., Liu, N., Song, M., Yang, Y., Yin, N., & Qi, J. (2021). Characterization and antioxidant activity of wheat bran polysaccharides modified by *Saccharomyces cerevisiae* and *Bacillus subtilis* fermentation. *Journal of Cereal Science*, *97*, Article 103157. <https://doi.org/10.1016/j.jcs.2020.103157>
- Christensen, L. F., García-Béjar, B., Bang-Berthelsen, C. H., & Hansen, E. B. (2022). Extracellular microbial proteases with specificity for plant proteins in food fermentation. *International Journal of Food Microbiology*, *381*, Article 109889. <https://doi.org/10.1016/j.ijfoodmicro.2022.109889>
- Chung, S. Y., Seo, Y. K., Park, J. M., Seo, M. J., Park, J. K., Kim, J. W., & Park, C. S. (2009). Fermented rice bran downregulates MITF expression and leads to inhibition of α -MSH-induced melanogenesis in B16F1 melanoma. *Bioscience, Biotechnology, and Biochemistry*, *73*, 1704–1710. <https://doi.org/10.1271/bbb.80766>
- Coda, R., Kärki, I., Nordlund, E., Heiniö, R. L., Poutanen, K., & Katina, K. (2014). Influence of particle size on bioprocess induced changes on technological functionality of wheat bran. *Food Microbiology*, *37*, 69–77. <https://doi.org/10.1016/j.fm.2013.05.011>
- Comín-Anduix, B., Boros, L. G., Marín, S., Boren, J., Callol-Massot, C., Centelles, J. J., Torres, J. L., Agell, N., Bassilian, S., & Cascante, M. (2002). Fermented wheat germ extract inhibits glycolysis/pentose cycle enzymes and induces apoptosis through poly (ADP-ribose) polymerase activation in Jurkat T-cell leukemia tumor cells. *Journal of Biological Chemistry*, *277*, 46408–46414. <https://doi.org/10.1074/jbc.M206150200>
- Courtin, C., & Delcour, J. (2002). Arabinoxylans and endoxylanases in wheat flour bread-making. *Journal of Cereal Science*, *35*, 225–243. <https://doi.org/10.1006/j.jcs.2001.0433>
- Deguyte-Fomins, L., Sontag-Strohm, T., & Salovaara, H. (2002). Oat bran fermentation by rye sourdough. *Cereal Chemistry*, *79*, 345–348. <https://doi.org/10.1094/CHEM.2002.79.3.345>
- dos Santos Costa, R., de Almeida, S. S., Cavalcanti, E. A. C., Guimarães Freire, D. M., Moura-Nunes, N., Monteiro, M., & Perrone, D. (2021). Enzymes produced by solid state fermentation of agro-industrial by-products release ferulic acid in bioprocessed whole-wheat breads. *Food Research International*, *140*, Article 109843. <https://doi.org/10.1016/j.foodres.2020.109843>
- FAO. (2019). *The state of food and agriculture 2019*. Food and Agriculture Organization of the United Nations Rome, 2019.
- Food waste and food waste prevention – estimates. Last Modified September 29, 2023. Accessed March 19, 2024. https://ec.europa.eu/eurostat/statistics-explained/index.php?title=Food_waste_and_food_waste_prevention_-_estimates.
- Gaur, G., & Gänzle, M. G. (2023). Conversion of (poly)phenolic compounds in food fermentations by lactic acid bacteria: Novel insights into metabolic pathways and functional metabolites. *Current Research in Food Science*, *6*, Article 100448. <https://doi.org/10.1016/j.crf.2023.100448>
- Gélinas, P., McKinnon, C. M., & Pelletier, M. (1999). Sourdough-type bread from waste bread crumb. *Food Microbiology*, *16*, 37–43. <https://doi.org/10.1006/fmic.1998.0209>
- Gómez, M., & Martínez, M. M. (2018). Fruit and vegetable by-products as novel ingredients to improve the nutritional quality of baked goods. *Critical Reviews in Food Science and Nutrition*, *58*(13), 2119–2135. <https://doi.org/10.1080/10408398.2017.1305946>
- Gupta, S., Jaiswal, A. K., & Abu-Ghannam, N. (2013). Optimization of fermentation conditions for the utilization of brewing waste to develop a nutraceutical rich liquid product. *Industrial Crops and Products*, *44*, 272–282. <https://doi.org/10.1016/j.indcrop.2012.11.015>
- Hoffmans, Y., Schaarschmidt, S., Fahl-Hassek, C., & van der Fels-Klerx, H. J. (2022). Factors during production of cereal-derived feed that influence mycotoxin contents. *Toxins*, *14*, 301. <https://doi.org/10.3390/toxins14050301>
- Holland, C., Ryden, P., Edwards, C. H., & Grundy, M. M. (2020). Plant cell walls: Impact on nutrient bioaccessibility and digestibility. *Foods*, *9*(2), 201. <https://doi.org/10.3390/foods9020201>
- Immonen, M., Maina, N. H., Coda, R., & Katina, K. (2021). The molecular state of gelatinized starch in surplus bread affects bread recycling potential. *Lebensmittel-Wissenschaft & Technologie – LWT*, *150*, Article 112071. <https://doi.org/10.1016/j.lwt.2021.112071>
- Immonen, M., Maina, N. H., Wang, Y., Coda, R., & Katina, K. (2020). Waste bread recycling as a baking ingredient by tailored lactic acid fermentation. *International Journal of Food Microbiology*, *327*, Article 108652. <https://doi.org/10.1016/j.ijfoodmicro.2020.108652>
- Kaditzky, S., & Vogel, R. F. (2008). Optimization of exopolysaccharide yields in sourdoughs fermented by lactobacilli. *European Food Research and Technology*, *228*, 291–299. <https://doi.org/10.1007/s00217-008-0934-7>
- Kajala, I., Mäkelä, J., Coda, R., Shukla, S., Shi, Q., Maina, N. H., Juvonen, R., Ekholm, P., Goyal, A., Tenkanen, M., & Katina, K. (2016). Rye bran as fermentation matrix boosts in situ dextran production by *Weissella confusa* compared to wheat bran. *Applied Microbiology and Biotechnology*, *100*, 3499–3510. <https://doi.org/10.1007/s00253-015-7189-6>
- Katina, K., Juvonen, R., Laitila, A., Flander, L., Nordlund, E., Kariluoto, S., Piironen, V., & Poutanen, K. (2012). Fermented wheat bran as a functional ingredient in baking. *Cereal Chemistry*, *89*, 126–134. <https://doi.org/10.1094/CHEM-08-11-0106>
- Katina, K., Laitila, A., Juvonen, R., Liukkonen, K.-H., Kariluoto, S., Piironen, V., Landberg, R., Åman, P., & Poutanen, K. (2007). Bran fermentation as a means to enhance technological properties and bioactivity of rye. *Food Microbiology*, *24*(2), 175–186. <https://doi.org/10.1016/j.fm.2006.07.012>
- Kavalopoulos, M., Stoumpou, V., Christofi, A., Mai, S., Barampouti, E. M., Moustakas, K., Malamis, D., & Loizidou, M. (2021). Sustainable valorisation pathways mitigating environmental pollution from brewers' spent grains. *Environmental Pollution*, *270*, Article 116069. <https://doi.org/10.1016/j.envpol.2020.116069>
- Kavitake, D., Devi, P. B., & Shetty, P. H. (2020). Overview of an exopolysaccharides produced by *Weissella* genus—A review. *International Journal of Biological Macromolecules*, *164*, 2964–2973. <https://doi.org/10.1016/j.ijbiomac.2020.08.185>
- Khodaei, D., Javanmardi, F., & Khaneghah, A. M. (2021). The global overview of the occurrence of mycotoxins in cereals: A three-year survey. *Current Opinion in Food Science*, *39*, 36–42. <https://doi.org/10.1016/j.cofs.2020.12.012>
- Khosroshahi, E. D., Razavi, S. H., Kaini, H., & Aghakhani, A. (2022). Improvement of stability and antioxidant activity of wheat germ by mixed fermentation versus single fermentation. *Journal of Food Science and Technology*, *59*, 2905–2912. <https://doi.org/10.1007/s13197-021-05316-w>
- Khubber, S., J Marti-Quijal, F. J., Tomasevic, I., Remize, F., & Barba, F. J. (2022). Lactic acid fermentation as a useful strategy to recover antimicrobial and antioxidant compounds from food and by-products. *Current Opinion in Food Science*, *43*, 189–198. <https://doi.org/10.1016/j.cofs.2021.11.013>
- Koirala, P., Costantini, A., Maina, N. H., Rizzello, C. G., Verni, M., Beni, V. D., Polo, A., Katina, K., Cagno, R. D., & Coda, R. (2022). Fermented brewers' spent grain containing dextran and oligosaccharides as ingredient for composite wheat bread and its impact on gut metabolome in vitro. *Fermentation*, *8*, 487. <https://doi.org/10.3390/fermentation8100487>
- Koirala, P., Maina, N. H., Nihtilä, H., Katina, K., & Coda, R. (2021). Brewers' spent grain as substrate for dextran biosynthesis by *Leuconostoc pseudomesenteroides* DSM20193 and *Weissella confusa* A16. *Microbial Cell Factories*, *20*(1), 1–13. <https://doi.org/10.1186/s12934-021-01515-4>
- Koistinen, V. M., Katina, K., Nordlund, E., Poutanen, K., & Hanhineva, K. (2016). Changes in the phytochemical profile of rye bran induced by enzymatic bioprocessing and sourdough fermentation. *Food Research International*, *89*, 1106–1115. <https://doi.org/10.1016/j.foodres.2016.06.027>
- Korcz, E., & Varga, L. (2021). Exopolysaccharides from lactic acid bacteria: Techno-functional application in the food industry. *Trends in Food Science & Technology*, *110*, 375–384. <https://doi.org/10.1016/j.tifs.2021.02.014>
- Korhola, M., Hakonen, R., Juuti, K., Edelmann, M., Kariluoto, S., Nyström, L., Sontag-Strohm, T., & Piironen, V. (2014). Production of folate in oat bran fermentation by yeasts isolated from barley and diverse foods. *Journal of Applied Microbiology*, *117*, 679–689. <https://doi.org/10.1111/jam.12564>
- Kosveva, M. R. (2020). Sources, characteristics and treatment of plant-based food waste. In M. R. Kosveva, & C. Webb (Eds.), *Food industry wastes* (pp. 37–66). Academic Press.
- Ktenioudaki, A., Alvarez-Jubete, L., Smyth, T. J., Kilcawley, K., Rai, D. K., & Gallagher, E. (2015). Application of bioprocessing techniques (sourdough fermentation and technological aids) for brewer's spent grain breads. *Food Research International*, *73*, 107–116. <https://doi.org/10.1016/j.foodres.2015.03.008>
- Lamont, J. R., Wilkins, O., Bywater-Ekegård, M., & Smith, D. L. (2017). From yogurt to yield: Potential applications of lactic acid bacteria in plant production. *Soil Biology and Biochemistry*, *111*, 1–9. <https://doi.org/10.1016/j.soilbio.2017.03.015>
- Li, N., Wang, S., Wang, T., Liu, R., Zhi, Z., Wu, T., Sui, W., & Zhang, M. (2022). Valorization of wheat bran by three fungi solid-state fermentation: Physicochemical properties, antioxidant activity and flavor characteristics. *Foods*, *11*(12), 1722. <https://doi.org/10.3390/foods11121722>
- Li, Y., Wang, H., Wang, L., Qiu, J., Li, Z., & Wang, L. (2023). Milling of wheat bran: Influence on digestibility, hydrolysis and nutritional properties of bran protein during in vitro digestion. *Food Chemistry*, *404*, Article 134559. <https://doi.org/10.1016/j.foodchem.2022.134559>
- Li, B., Zhao, L., Chen, H., Sun, D., Deng, B., Li, J., Liu, Y., & Wang, F. (2016). Inactivation of lipase and lipoxygenase of wheat germ with temperature-controlled short wave infrared radiation and its effect on storage stability and quality of wheat germ oil. *PLoS One*, *11*(12), Article e0167330. <https://doi.org/10.1371/journal.pone.0167330>
- Liu, Q., Cao, X., Zhuang, X., Han, W., Guo, W., Xiong, J., & Zhang, X. (2017). Rice bran polysaccharides and oligosaccharides modified by *Grifola frondosa* fermentation: Antioxidant activities and effects on the production of NO. *Food Chemistry*, *223*, 49–53. <https://doi.org/10.1016/j.foodchem.2016.12.018>
- Liu, L., Zhang, R., Deng, Y., Zhang, Y., Xiao, J., Huang, F., Wen, W., & Zhang, M. (2017). Fermentation and complex enzyme hydrolysis enhance total phenolics and antioxidant activity of aqueous solution from rice bran pretreated by steaming with α -amylase. *Food Chemistry*, *221*, 636–643. <https://doi.org/10.1016/j.foodchem.2016.11.126>
- Lynch, K. M., Steffen, E. J., & Arendt, E. K. (2016). Brewers' spent grain: A review with an emphasis on food and health. *Journal of the Institute of Brewing*, *122*, 553–568. <https://doi.org/10.1002/jib.363>
- Ma, S., Wang, Z., Guo, X., Wang, F., Huang, J., Sun, B., & Wang, X. (2021). Sourdough improves the quality of whole-wheat flour products: Mechanisms and challenges—a review. *Food Chemistry*, *360*, Article 130038. <https://doi.org/10.1016/j.foodchem.2021.130038>

- Mabunga, D. F. N., Gonzales, E. L. T., Kim, H. J., & Choung, S. Y. (2015). Treatment of GABA from fermented rice germ ameliorates caffeine-induced sleep disturbance in mice. *Biomolecules & Therapeutics*, 23, 268–274. <https://doi.org/10.4062/biomolther.2015.022>
- Manini, F., Brasca, M., Plumed-Ferrer, C., Morandi, S., Erba, D., & Casiraghi, M. C. (2014). Study of the chemical changes and evolution of microbiota during sourdough like fermentation of wheat bran. *Cereal Chemistry*, 91, 342–349. <https://doi.org/10.1094/CCHEM-09-13-0190-CESI>
- Mao, M., Wang, P., Shi, K., Lu, Z., Bie, X., Zhao, H., Zhang, C., & Lv, F. (2020). Effect of solid state fermentation by *Enterococcus faecalis* M2 on antioxidant and nutritional properties of wheat bran. *Journal of Cereal Science*, 94, Article 102997. <https://doi.org/10.1016/j.jcs.2020.102997>
- Martău, G. A., Teleky, B.-E., Ranga, F., Pop, I. D., & Vodnar, D. C. (2021). Apple pomace as a sustainable substrate in sourdough fermentation. *Frontiers in Microbiology*, 12, Article 742020. <https://doi.org/10.3389/fmicb.2021.742020>
- Marti, A., Torri, L., Casiraghi, M. C., Franzetti, L., Limbo, S., Morandin, F., Quaglia, L., & Paganì, M. A. (2014). Wheat germ stabilization by heat-treatment or sourdough fermentation: Effects on dough rheology and bread properties. *Lebensmittel-Wissenschaft und -Technologie- Food Science and Technology*, 59(2P1), 1100–1106. <https://doi.org/10.1016/j.lwt.2014.06.039>
- Mendez-Vilas, A. (2007). *Communicating current research and educational topics and trends in applied microbiology*.
- Mikušová, L., Gereková, P., Kocková, M., Šturdík, E., Valachovičová, M., Holubková, A., Vajdák, M., & Mikuš, L. (2013). Nutritional, antioxidant, and glycaemic characteristics of new functional bread. *Chemical Papers*, 67, 284–291. <https://doi.org/10.2478/s11696-012-0258-8>
- Moon, S. H., & Chang, H.-C. (2021). Rice bran fermentation using *Lactiplantibacillus plantarum* EM as a starter and the potential of the fermented rice bran as a functional food. *Foods*, 10, 978. <https://doi.org/10.3390/foods10050978>
- Muhialdin, B. J., Saari, N., & Meor Hussin, A. S. (2020). Review on the biological detoxification of mycotoxins using lactic acid bacteria to enhance the sustainability of foods supply. *Molecules*, 25, 2655. <https://doi.org/10.3390/molecules25112655>
- Nionelli, L., Wang, Y., Pontonio, E., Immonen, M., Rizzello, C. G., Maina, H. N., Katina, K., & Coda, R. (2020). Antifungal effect of bioprocessed surplus bread as ingredient for bread-making: Identification of active compounds and impact on shelf-life. *Food Control*, 118, Article 107437. <https://doi.org/10.1016/j.foodcont.2020.107437>
- Nordlund, E., Katina, K., Aura, A. M., & Poutanen, K. (2013). Changes in bran structure by bioprocessing with enzymes and yeast modifies the in vitro digestibility and fermentability of bran protein and dietary fibre complex. *Journal of Cereal Science*, 58, 200–208. <https://doi.org/10.1016/j.jcs.2013.05.006>
- O'Connor, J., Hoang, S. A., Bradney, L., Dutta, S., Xiong, X., Tsang, D. C. W., Ramadass, K., Vinu, A., Kirkham, M. B., & Bolan, N. S. (2021). A review on the valorisation of food waste as a nutrient source and soil amendment. *Environmental Pollution*, 272, Article 115985. <https://doi.org/10.1016/j.envpol.2020.115985>
- Patterson, C. A., Curran, J., & Der, T. (2017). Effect of processing on antinutrient compounds in pulses. *Cereal Chemistry*, 94(1), 2–10. <https://doi.org/10.1094/CCHEM-05-16-0144-FI>
- Petrova, P., & Petrov, K. (2017). Prebiotic–probiotic relationship: The genetic fundamentals of polysaccharides conversion by *Bifidobacterium* and *Lactobacillus* genera. In A. M. Grumezescu, & A. M. Holban (Eds.), *Handbook of food bioengineering, food bioconversion* (pp. 237–278). Academic Press.
- Piotrowska, M. (2021). Microbiological decontamination of mycotoxins: Opportunities and limitations. *Toxins*, 13, 819. <https://doi.org/10.3390/toxins13110819>
- Pontonio, E., Arora, K., Dingeo, C., Carafa, I., Celano, G., Scarpino, V., Genot, B., Gobbetti, M., & Di Cagno, R. (2021). Commercial organic versus conventional whole rye and wheat flours for making sourdough bread: Safety, nutritional, and sensory implications. *Frontiers in Microbiology*, 12, Article 674413. <https://doi.org/10.3389/fmicb.2021.674413>
- Pontonio, E., Dingeo, C., Di Cagno, R., Blandino, M., Gobbetti, M., & Rizzello, C. G. (2020). Brans from hull-less barley, emmer and pigmented wheat varieties: From by-products to bread nutritional improvers using selected lactic acid bacteria and xylanase. *International Journal of Food Microbiology*, 313, Article 108384. <https://doi.org/10.1016/j.ijfoodmicro.2019.108384>
- Pontonio, E., Dingeo, C., Gobbetti, M., & Rizzello, C. G. (2019). Maize milling by-products: From food wastes to functional ingredients through lactic acid bacteria fermentation. *Frontiers in Microbiology*, 10, 561. <https://doi.org/10.3389/fmicb.2019.00561>
- Pontonio, E., Lorusso, A., Gobbetti, M., & Rizzello, C. G. (2017). Use of fermented milling by-products as functional ingredient to develop a low-glycaemic index bread. *Journal of Cereal Science*, 77, 235–242. <https://doi.org/10.1016/j.jcs.2017.08.022>
- Pontonio, E., Verni, M., Dingeo, C., Diaz-de-Cerio, E., Pinto, D., & Rizzello, C. G. (2020). Impact of enzymatic and microbial bioprocessing on antioxidant properties of hemp (*Cannabis sativa* L.). *Antioxidants*, 9(12), 1258. <https://doi.org/10.3390/antiox9121258>
- Poutanen, K., Flander, L., & Katina, K. (2009). Sourdough and cereal fermentation in a nutritional perspective. *Food Microbiology*, 26, 693–699. <https://doi.org/10.1016/j.fm.2009.07.011>
- Rizzello, C. G., Mueller, T., Coda, R., Reipsch, F., Nionelli, L., Curiel, J. A., & Gobbetti, M. (2013). Synthesis of 2-methoxybenzoquinone and 2, 6-dimethoxybenzoquinone by selected lactic acid bacteria during sourdough fermentation of wheat germ. *Microbial Cell Factories*, 12, 105. <https://doi.org/10.1186/1475-2859-12-105>
- Rizzello, C. G., Nionelli, L., Coda, R., De Angelis, M., & Gobbetti, M. (2010a). Effect of sourdough fermentation on stabilisation, and chemical and nutritional characteristics of wheat germ. *Food Chemistry*, 119(3), 1079–1089. <https://doi.org/10.1016/j.foodchem.2009.08.016>
- Rizzello, C. G., Nionelli, L., Coda, R., Di Cagno, R., & Gobbetti, M. (2010b). Use of sourdough fermented wheat germ for enhancing the nutritional, texture and sensory characteristics of the white bread. *European Food Research and Technology*, 230(4), 645–654. <https://doi.org/10.1007/s00217-009-1204-z>
- ICMSF, International Commission on Microbiological Specifications for Foods Meat and Meat Products. (2000). In A. Roberts, J. I. Pitt, J.-L. Cordier, L. G. M. Gorris, L. Gram, K. M. J. Swanson, & R. B. Tompkin (Eds.), *Microbial ecology of food commodities*, 6. New York: Kluwer Academic/Plenum Publishers.
- Sabater, C., Ruiz, L., Delgado, S., Ruas-Madiedo, P., & Margolles, A. (2020). Valorization of vegetable food waste and by-products through fermentation processes. *Frontiers in Microbiology*, 11, Article 581997. <https://doi.org/10.3389/fmicb.2020.581997>
- Sadeghi, A., Ebrahimi, M., Raeisi, M., & Nematollahi, Z. (2019). Biological control of foodborne pathogens and aflatoxins by selected probiotic LAB isolated from rice bran sourdough. *Biological Control*, 130, 70–79. <https://doi.org/10.1016/j.biocontrol.2018.09.017>
- Saiko, P., Oszvar-Kozma, M., Graser, G., Lackner, A., Grusch, M., Madlener, S., Krupitza, G., Jaeger, W., Hidvegi, M., Agarwal, R. P., & Fritzer-Szekeres, M. (2009). Avemar, a nontoxic fermented wheat germ extract, attenuates the growth of sensitive and 5-FdUrd/Ara-C cross-resistant H9 human lymphoma cells through induction of apoptosis. *Oncology Reports*, 21, 787–791. <https://doi.org/10.3892/or.00000285>
- Saurabh, V., Urhe, S. B., Upadhyay, A., & Shankar, S. (2021). Antinutritional factors in cereals. In S. P. Bangar, A. Siroha, & M. Kumar (Eds.), *Handbook of cereals, pulses, roots, and tubers*. Imprint CRC Press.
- Savolainen, O. I., Coda, R., Suomi, K., Katina, K., Juvonen, R., Hanhineva, K., & Poutanen, K. (2014). The role of oxygen in the liquid fermentation of wheat bran. *Food Chemistry*, 153, 424–431. <https://doi.org/10.1016/j.foodchem.2013.12.062>
- Schettino, R., Verni, M., Acin-Albiac, M., Vincentini, O., Krona, A., Knaapila, A., Cagno, R. D., Gobbetti, M., Rizzello, C. G., & Coda, R. (2021). Bioprocessed brewers' spent grain improves nutritional and antioxidant properties of pasta. *Antioxidants*, 10, 742. <https://doi.org/10.3390/antiox10050742>
- Seo, Y. K., Jung, S. H., Song, K. Y., Park, J. K., & Park, C. S. (2010). Anti-photoaging effect of fermented rice bran extract on UV-induced normal skin fibroblasts. *European Food Research and Technology*, 231, 163–169. <https://doi.org/10.1007/s00217-010-1261-3>
- Siroli, L., Giordani, B., Rossi, S., Gottardi, D., McMahon, H., Augustyniak, A., Menon, A., Vannini, L., Vitali, B., Patrignani, F., & Lanciotti, R. (2022). Antioxidant and functional features of pre-fermented ingredients obtained by the fermentation of milling by-products. *Fermentation*, 8, 722. <https://doi.org/10.3390/fermentation8120722>
- Spaggiari, M., Calani, L., Folloni, S., Ranieri, R., Dall'Asta, C., & Galaverna, G. (2020). The impact of processing on the phenolic acids, free betaine and choline in Triticum spp. L. whole grains and milling by-products. *Food Chemistry*, 311, Article 125940. <https://doi.org/10.1016/j.foodchem.2019.125940>
- Spaggiari, M., Ricci, A., Calani, L., Bresciani, L., Neviani, E., Dall'Asta, C., Lazzi, C., & Galaverna, G. (2020). Solid state lactic acid fermentation: A strategy to improve wheat bran functionality. *Lebensmittel-Wissenschaft & Technologie*, 118, Article 108668. <https://doi.org/10.1016/j.lwt.2019.108668>
- Teixeira, M. R., Guarda, E. C., Freitas, E. B., Galinha, C. F., Duque, A. F., & Reis, M. A. M. (2020). Valorization of raw brewers' spent grain through the production of volatile fatty acids. *New Biotechnology*, 57, 4–10. <https://doi.org/10.1016/j.nbt.2020.01.007>
- Tlais, A. Z. A., Da Ros, A., Filannino, P., Vincentini, O., Gobbetti, M., & Di Cagno, R. (2021). Biotechnological re-cycling of apple by-products: A reservoir model to produce a dietary supplement fortified with biogenic phenolic compounds. *Food Chemistry*, 336, Article 127616. <https://doi.org/10.1016/j.foodchem.2020.127616>
- Tovar, L. E. R., & Gänzle, M. G. (2021). Degradation of wheat germ agglutinin during sourdough fermentation. *Foods*, 10, 340. <https://doi.org/10.3390/foods10020340>
- Verni, M., Minisci, A., Convertino, S., Nionelli, L., & Rizzello, C. G. (2020). Wasted bread as substrate for the cultivation of starters for the food industry. *Frontiers in Microbiology*, 11, 293. <https://doi.org/10.3389/fmicb.2020.00293>
- Verni, M., Pontonio, E., Krona, A., Jacob, S., Pinto, D., Rinaldi, F., Verardo, V., Diaz de Cerio, E., Coda, R., & Rizzello, C. G. (2020). Bioprocessing of brewers' spent grain enhances its antioxidant activity: Characterization of phenolic compounds and bioactive peptides. *Frontiers in Microbiology*, 11, 1831. <https://doi.org/10.3389/fmicb.2020.01831>
- Verni, M., Rizzello, C. G., & Coda, R. (2019). Fermentation biotechnology applied to cereal industry by-products: Nutritional and functional insights. *Frontiers in Nutrition*, 6, 42. <https://doi.org/10.3389/fnut.2019.00042>
- Verni, M., Vekka, A., Immonen, M., Rizzello, C. G., Katina, K., & Coda, R. (2022). Biosynthesis of γ -aminobutyric acid by lactic acid bacteria in surplus bread and its use in bread-making. *Journal of Applied Microbiology*, 133, 76–90. <https://doi.org/10.1111/jam.15332>
- Verni, M., Verardo, V., & Rizzello, C. G. (2019). How fermentation affects the antioxidant properties of cereals and legumes. *Foods*, 8(9), 362. <https://doi.org/10.3390/foods8090362>
- Vriesekoop, F., Haynes, A., van der Heijden, N., Liang, H., Paximada, P., & Zuidberg, A. (2021). Incorporation of fermented brewers spent grain in the production of sourdough bread. *Fermentation*, 7(2), 96. <https://doi.org/10.3390/fermentation7020096>
- Wang, B., Li, G., Li, L., Zhang, M., Yang, T., Xu, Z., & Qin, T. (2022a). Novel processing strategies to enhance the bioaccessibility and bioavailability of functional components in wheat bran. *Critical Reviews in Food Science and Nutrition*, 3, 1–15. <https://doi.org/10.1080/10408398.2022.2129582>
- Wang, Y., Rosa-Sibakov, N., Edelmann, M., Sozer, N., Katina, K., & Coda, R. (2022b). Enhancing the utilization of rapeseed protein ingredients in bread making by

- tailored lactic acid fermentation. *Food Bioscience*, 50, Article 102028. <https://doi.org/10.1016/j.fbio.2022.102028>
- Waters, D. M., Jacob, F., Titze, J., Arendt, E. K., & Zannini, E. (2012). Fibre, protein and mineral fortification of wheat bread through milled and fermented brewer's spent grain enrichment. *European Food Research and Technology*, 235, 767–778. <https://doi.org/10.1007/s00217-012-1805-9>
- Wu, J., Ren, L., Zhao, N., Wu, T., Liu, R., Sui, W., & Zhang, M. (2022). Solid-state fermentation by *Rhizopus oryzae* improves flavor of wheat bran for application in food. *Journal of Cereal Science*, 107, Article 103536. <https://doi.org/10.1016/j.jcs.2022.103536>
- Yeong, M. S., Hee, M. S., & Choon, C. H. (2020). Characterization of high-ornithine-producing *Weissella koreensis* DB1 isolated from kimchi and its application in rice bran fermentation as a starter culture. *Foods*, 9(11), 1545. <https://doi.org/10.3390/foods9111545>
- Zadeike, D., Gaizauskaite, Z., Svazas, M., Gruzauskas, R., Gruzauskas, V., Damasius, J., & Juodeikiene, G. (2022). Application of solid-state fermentation for the improving of extruded corn dry-milling by-products and their protein functional properties. *Life*, 12, 1909. <https://doi.org/10.3390/life12111909>
- Zadeike, D., Vaitkeviciene, R., Bartkevics, V., Bogdanova, E., Bartkiene, E., Lele, V., Juodeikiene, G., Cernauskas, D., & Valatkeviciene, Z. (2021). The expedient application of microbial fermentation after whole-wheat milling and fractionation to mitigate mycotoxins in wheat-based products. *Lebensmittel-Wissenschaft & Technologie – LWT*, 137, Article 110440. <https://doi.org/10.1016/j.lwt.2020.110440>
- Zhang, D., Liu, H., Wang, S., Liu, Y., & Ji, H. (2023). Wheat bran fermented by *Lactobacillus* regulated the bacteria–fungi composition and reduced fecal heavy metals concentrations in growing pigs. *Science of The Total Environment*, 858(3), Article 159828. <https://doi.org/10.1016/j.scitotenv.2022.159828>
- Zhang, Y., Wang, P., Kong, Q., & Cotty, P. J. (2021). Biotransformation of aflatoxin B1 by *Lactobacillus helveticus* FAM22155 in wheat bran by solid-state fermentation. *Food Chemistry*, 341, Article 128180. <https://doi.org/10.1016/j.foodchem.2020.128180>
- Zhang, J., Xiao, X., Dong, Y., Wu, J., Yao, F., & ZhouZhang, X. (2015). Effect of fermented wheat germ extract with *Lactobacillus plantarum* dy-1 on HT-29 cell proliferation and apoptosis. *Journal of Agricultural and Food Chemistry*, 63(9), 2449–2457. <https://doi.org/10.1021/acs.jafc.5b00041>
- Zhao, H. M., Guo, X. N., & Zhu, K. X. (2017). Impact of solid state fermentation on nutritional, physical and flavor properties of wheat bran. *Food Chemistry*, 217, 28–36. <https://doi.org/10.1016/j.foodchem.2016.08.062>
- Zhao, I., Liao, A. M., Liu, N., Huang, J. H., Lv, X., Yang, C. R., Chen, W. J., Hou, Y. C., Ma, L. J., & Hui, M. (2021). Potential anti-aging effects of fermented wheat germ in aging mice. *Food Bioscience*, 42, Article 101182. <https://doi.org/10.1016/j.fbio.2021.101182>
- Zokaityte, E., Lele, V., Starkute, V., Zavistanaviciute, P., Klupsaite, D., Bartkevics, V., Pugajeva, I., Bērziņa, Z., Gruzauskas, R., Sidlauskienė, S., Juodeikiene, G., Santini, A., & Bartkiene, E. (2021). The influence of combined extrusion and fermentation processes on the chemical and biosafety parameters of wheat bran. *Lebensmittel-Wissenschaft & Technologie – LWT*, 146, Article 111498. <https://doi.org/10.1016/j.lwt.2021.111498>