



THE ITALIAN QUATERNARY VOLCANISM

Stefano Branca¹, Alessandra Cinquegrani², Raffaello Cioni³, Aida Maria Conte⁴,
 Sandro Conticelli^{3,4}, Gianfilippo De Astis⁵, Sandro de Vita⁶, Rosanna De Rosa⁷,
 Mauro Antonio Di Vito⁶, Paola Donato⁷, Francesca Forni⁸, Lorella Francalanci³, Mario Gaeta⁹,
 Biagio Giacco⁴, Guido Giordano^{10,4}, Marisa Giuffrida¹¹, Roberto Isaia⁶, Federico Lucchi¹²,
 Fabrizio Marra⁵, Silvia Massaro¹³, Eugenio Nicotra⁷, Danilo M. Palladino⁸, Cristina Perinelli^{3,4},
 Paola Petrosino¹⁴, Marco Pistoletti¹⁵, José Pablo Sepulveda-Birke³, Gianluca Sottilli⁹,
 Claudia Romagnoli¹², Silvio Rotolo^{2,16}, Roberto Sulpizio¹³, Claudio Antonio Tranne¹²,
 Marco Viccaro^{11,1}

¹ Istituto Nazionale di Geofisica e Vulcanologia - Sezione di Catania, Osservatorio Etna, Italy.

² Dipartimento di Scienze della Terra e del Mare, Università degli Studi di Palermo, Palermo, Italy.

³ Dipartimento di Scienze della Terra, Università degli Studi di Firenze, Firenze, Italy.

⁴ CNR - Istituto di Geologia Ambientale e Geoingegneria, Montelibretti e Roma, Italy.

⁵ Istituto Nazionale di Geofisica e Vulcanologia - Sezione di Sismologia e Tectonofisica, Roma, Italy.

⁶ Istituto Nazionale di Geofisica e Vulcanologia - Osservatorio Vesuviano, Napoli, Italy.

⁷ Dipartimento di Biologia, Ecologia e Scienze della Terra, Università della Calabria, Arcavacata di Rende (CS), Italy.

⁸ Dipartimento di Scienze della Terra, Università di Milano La Statale, Milano (MI), Italy.

⁹ Dipartimento di Scienze della Terra, Sapienza, Università di Roma, Roma, Italy.

¹⁰ Dipartimento di Scienze dell'Università degli Studi Roma III, Roma, Italy.

¹¹ Dipartimento di Scienze Biologiche, Geologiche e Ambientali, Università di Catania, Catania, Italy.

¹² Dipartimento di Scienze Biologiche, Geologiche e Ambientali, Università di Bologna, Bologna, Italy.

¹³ Dipartimento di Scienze della Terra e Geoambientali, Università di Bari, Italy.

¹⁴ Dipartimento di Scienze della Terra, dell'Ambiente e delle Risorse, Università degli Studi di Napoli Federico II, Napoli, Italy.

¹⁵ Dipartimento di Scienze della Terra dell'Università degli Studi di Pisa, Pisa, Italy.

¹⁶ Istituto Nazionale di Geofisica e Vulcanologia - Sezione di Palermo, Palermo, Italy.

Corresponding authors: Biagio Giacco <biagio.giacco@cnr.it> and Paola Petrosino <paola.petrosino@unipa.it>

ABSTRACT: The peninsular and insular Italy are punctuated by Quaternary volcanoes and their rocks constitute an important aliquot of the Italian Quaternary sedimentary successions. Also away from volcanoes themselves, volcanic ash layers are a common and frequent feature of the Quaternary records, which provide us with potential relevant stratigraphic and chronological markers at service of a wide array of the Quaternary science issues. In this paper, a broad representation of the Italian volcanological community has joined to provide an updated comprehensive state of art of the Italian Quaternary volcanism. The eruptive history, style and dynamics and, in some cases, the hazard assessment of about thirty Quaternary volcanoes, from the north-earliest Mt. Amiata, in Tuscany, to the southernmost Pantelleria and Linosa, in Sicily Channel, are here reviewed in the light of the substantial improving of the methodological approaches and the overall knowledge achieved in the last decades in the volcanological field study. We hope that the present review can represent a useful and agile document summarising the knowledge on the Italian volcanism at the service of the Quaternary community operating in central Mediterranean area.

Keywords: Italian volcanism, stratigraphy, geochronology, eruptive dynamics, distal tephra.

1. INTRODUCTION (S.C., B.G., P.P.)

Volcanic products undoubtedly represent an important part of the Quaternary deposits in Italy. For this reason, we have decided to take the opportunity offered by the INQUA conference to be held in Rome in 2023 to present a review of the state of the art and knowledge of the Italian Quaternary volcanism. A few remarks should be made to explain why we felt the need to do this.

The activity of the Italian Quaternary volcanoes during the last century has been the subject of intense investigations, although at first the main focus was on mineralogical-petrographical-magmatic aspects other than on the strictly volcanological ones. It was probably in the decade 1980–1990, with the publication of the results of the "Progetto Finalizzato Geodinamica"

of the Italian National Research Council (CNR), that monographic works were published on some volcanic areas, such as the Neapolitan volcanoes or Etna, which, in addition to the petrographic features, proposed a systematic study of the stratigraphic aspects of the products, mapping their distribution and thus providing a reconstruction of their eruptive history (e.g., Santacroce, 1987 for Somma-Vesuvio, Rosi & Sbrana, 1987 for Campi Flegrei, Vezzoli, 1988 for Ischia, Romano et al., 1979 for Etna, etc.).

Since then, and especially in the last two decades, this reconstruction has been progressively refined due to the improved knowledge of volcanic history and stratigraphy, in particular thanks to the Cartografia geologica e geomatica (CARG) project (e.g., Groppelli & Viereck-Goette, 2010), dedicated to updating the geological




Fig. 1 - Location of the Italian Quaternary volcanoes with indication of the time interval of activity (Ma, unless differently indicated).

cal mapping of the Italian territory. A multidisciplinary approach (stratigraphic, geomorphological, geochemical) allowed the reconstruction of the time-space history of most of the Italian Quaternary volcanic areas, which has been furthered by the increased ability to obtain reliable $^{40}\text{Ar}/^{39}\text{Ar}$ ages on both pyroclastic deposits and lavas. In fact, the significant technological developments of noble gas mass spectrometers over the last decade (Mixon et al., 2022), have improved the effectiveness of the method and the possibility of getting accurate and high-precision $^{40}\text{Ar}/^{39}\text{Ar}$ dating allowing a detailed and robust reconstruction of the eruptive history. This has been made possible and facilitated by the peculiar potassium nature of the magmas feeding the peri-Tyrrhenian potassium Quaternary volcanoes (e.g., Peccerillo, 2017), which generated products bearing K-rich minerals (e.g., sanidine and leucite), ideal for direct $^{40}\text{Ar}/^{39}\text{Ar}$ dating. Furthermore, with regard to the explosive activity, the great deal of tephrostratigraphic research on distal stratigraphic successions, i.e., located tens to thousands of km from the volcanic sources, substantially contributed to increase the knowledge on the recurrence, temporal extent and dynamics of explosive activity, providing integrative information that are difficult to detect in the proximal areas (e.g., Monaco et al., 2021, 2022a; Leicher et al., 2023). Finally, the last twenty years have seen an increase in research aimed at evaluating volcanic risk in areas of active volcanism, with particular reference to hazard assessment based on knowledge of past eruptive history (e.g., Cioni et al., 2003; Lirer et al., 2010; Del Negro et al., 2013; Bevilacqua et al., 2022 and references therein).

Given these premises, it seemed important to gather the contributions of various members of the Italian volcanological community in order to present an updated review on the Italian Quaternary volcanoes. To emphasize the volcanological aspects over the petrographic-magmatological ones as companion papers we deliberately decided to not adopt the division into magmatic provinces (see Conticelli et al., 2010, 2015a; Peccerillo, 2017; and references in these papers and book) but to list the Italian Quaternary volcanoes in a simple and neutral geographical way, starting from Tuscany and then moving southward on to those located in Lazio, Campania, Tyrrhenian Sea and Sicily (Fig. 1).

For each volcanic area or volcano dealt with, a summary of the eruptive history is proposed, focusing on the dynamics in relation to time, with emphasis on the most recent researches, and shortly mentioning the chemistry of the products. For the active volcanic areas, particularly those exposed to higher risk, information is also given on the state of hazard and risk assessment research.

We hope that this paper will be a valid support for the studies of the Quaternary community, which very often has to deal with the eruptive history of the Italian volcanoes and for which it is extremely necessary to have an agile, exhaustive and up-to-date general framework.

2. GEODYNAMIC FRAMEWORK OF QUATERNARY VOLCANIC ACTIVITY IN ITALY (S.C., B.G., P.P.)

The Tyrrhenian Sea is one of the most geologically complex areas in the world (Faccenna et al., 2001b;

Cavazza et al., 2004; Beltrando et al., 2010). The reasons for this complexity are mainly linked to its role in the convergence of the African and European plates, which starting from the Cretaceous to the Present time, has led to the formation of two orogenic belts (Alps and Apennine Chain) and several back-arc basins of progressively younger age moving from W to E in the western Mediterranean (Sartori, 2003). Magmatic processes are integral part of this geodynamically complex region and punctuated all stages of its evolution. In the initial stages this activity was concentrated in the Alps (e.g., Conticelli et al., 2009b, 2010; Alagna et al., 2010) along the margins of the colliding plates and, to a lesser extent, within the Africa-Adria plate. The last phase of this complex geodynamic magmatic pattern began in the Upper Eocene-Oligocene and is still on-going. This is coeval with the formation of the Apennine-Maghrebian chain, and the opening of the western Mediterranean basin (Lustrino et al., 2011; Di Capua et al., 2016), and has therefore been referred to as "the Apennine magmatism" (Peccerillo, 2017). Opening of slab tears and incipient slab detachment also occurred in the late evolution of the Apennine orogen, favouring the channelling of intra-plate volcanism that mixed with the orogenic one during the final stage of Vulture volcanism (e.g., Faccenna et al., 2001a, 2001b, 2004; Avanzinelli et al., 2008, 2009; Conticelli et al., 2009b). The Apennine orogen has been subject to back-arc extension since the Late Miocene due to the rapid slab roll-back of the Calabrian block (Malinverno & Ryan, 1988), which has produced two major extensional basins in the abyssal plain of the Tyrrhenian Sea, the Vavilov and the Marsili basins, located at north-west and south-east portions of the Southern Tyrrhenian basin, respectively (Marani and Gamberi, 2004). The opening of the Marsili back-arc basin also coincides with the formation of the Aeolian volcanic arc, whose spatial evolution is linked to the decoupling and roll-back of the slab beneath the Calabrian Arc (Loreto et al., 2020). At the back of the eastward migrating Apennine orogen intense extension occurred since the Upper Miocene (Jolivet et al., 1998, and references therein). In the central Tyrrhenian margin, the onset of extensional tectonics can be dated at Late Miocene, the age of syn-rift sedimentary sequences filling in the extensional basins located between the Tolla-Ceriti and Roccamonfina volcanoes (Mattie et al., 2010). The Tyrrhenian margin of southern Italy, on the counterpart, has been mainly characterized by Plio-Quaternary back-arc extension, with the development of NW-SE normal faults (Acocella and Funiciello, 2006); coeval NE-SW transverse systems are also present. Despite the overall NW-SE alignment of the Quaternary volcanoes along the Tyrrhenian margin, most of them formed where NW-SE normal faults intersect transverse NE-SW tectonic lineaments which, thanks to the higher vertical depth with respect to the NW trending features, represent a preferential structure for magma uprising and fluids upwelling (Acocella and Funiciello, 2002). The volcanic edifices consist of poly-phase calderas and large ignimbrite plateaus, which identify vast areas with negligible topographic relief (Vulsini, Sabatini, the Vulcano Laziale - the early edifice of the Colli Albani volcano, Campi Flegrei) and stratovolcanoes with summit calderas (e.g., Vico, Faeto - the intracaldera edifice of Colli Albani, Roccamonfina and Vesuvius).




Fig. 2 - a) Distribution of ultrapotassic rocks in the Central Italian region revealing the location the Torre Alfina, Radicofani and Monte Amiata volcanoes (spots within the large rectangular shapes); and the distribution of the major magmatic potassiac and ultrapotassiac events. Red spots and areas (kamafugites) and dark green spots (lamproites) mark the location site of the most extreme endmembers of the orogenic magmatic activity. Modified after Conticelli et al. (2010, 2013, 2015a) and Günther et al. (2023); b) Geological sketch map of the Torre Alfina volcano. Legend: 1) Cretaceous carbonaceous and argillaceous sedimentary rocks; 2) blocky olivine-litic lava flows, type B; 3) low viscosity olivine-litic lava flows, type B; 4) low-viscosity olivine-litic lava flows, type A; 5) lapilli and scoriae; 6) leucite-tephrite to leucite-phonolite Vulcini lavas; 7) leucite-phonolite to leucite-trachyte Vulcini pyroclastic rocks. Redrawn after Conticelli (1998) and Avanzinelli et al. (2017).

3. THE CENTRAL ITALY QUATERNARY VOLCANOES

3.1. Tuscany (S.C., J.P.S.-B.)

3.1.1. General background

The oldest Potassiac to Ultrapotassiac volcanic rocks of the Central Mediterranean region are found on the western margin of the Tyrrhenian basin, distributed N-S (Fig. 2), in Corsica at Sisco (14.1 Ma; Civetta et al., 1978), at Capraia Island (7.5-4.6 Ma; Gasparon et al., 2009) and at Cornacya Seamount (12.6 Ma; Mascle et al., 2001). These are leucite-free, lamproite-like, ultrapotassiac rocks associated with shoshonitic and high-K calc-alkaline volcanic rocks (Conticelli et al., 2009a). Following the rollback of the slab and the consequent opening of the Tyrrhenian basin, the volcanism shifted diachronically eastward, from Miocene to Pliocene, from Corsica to Tuscany (Fig. 2) (e.g., Avanzinelli et al., 2009; Conticelli et al., 2010, 2015a).

In Tuscany, lamproite-like, leucite-free ultrapotassiac rocks and associate shoshonitic and calc-alkaline volcanic rocks of ultimate mantle origin are distributed on both sides of the western shoreline of the Italian peninsula (Fig. 2a; e.g., Peccerillo et al., 1987; Conticelli & Peccerillo, 1992; Conticelli et al., 2009a, 2010, 2015; Peccerillo, 2005, 2017), and in some cases these volcanic rocks are intimately associated, in space, but not in time, with crustal-derived rhyolitic to granitic rocks (Peccerillo et al., 1987; Conticelli et al., 2002, 2004, 2010; Conte & Dolfi, 2002).

The oldest ultrapotassiac to high-K calc-alkaline igneous rocks of the so called "Tuscan Magmatic Province" (Conticelli & Peccerillo, 1992) are found at Elba Island, in the archipelago between Italian Peninsula and Corsica with a $^{40}\text{Ar}/^{39}\text{Ar}$ age of 5.89 Ma (Conticelli et al., 2001). These were followed by the emplacement, in the mainland, of the Val d'Ena lamproites made of the hypabissal minette of Montecatini Val di Cecina, emplaced at

4.1 Ma (K/Ar; Borsi et al., 1987), and lamproite-like "orendite" of Orciatico (Conticelli et al., 1992), of the latic dykes at Campiglia (Conticelli et al., 2004), and in the Latium region of the volcanic products of the Tolfa-Manziana-Cerifì dome complexes (Fig. 2a) between 4.3 and 1.9 Ma (e.g., Evernden & Curtis, 1965; Fazzini et al., 1972; Bigazzi et al., 1979; Lombardi et al., 1974; Villa et al., 1989).

The Quaternary volcanic rocks are instead found at Radicofani, Torre Alfina, Monte Cimino, Ponza Island (Monte La Guardia) and in drills in the Campania region. These leucite-free ultrapotassiac, lamproite-like, to shoshonitic and calc-alkaline rocks were emplaced between 1.3 and 0.85 Ma (e.g., Evernden & Curtis, 1965; Niccolatti, 1969; Pasquarello et al., 1983; Solevanti, 1983; D'Orazio et al., 1991; Laurenzi et al., 2015; Conte et al., 2016; Nappi et al., 2022).

3.1.2. Torre Alfina monogenetic volcano

The Torre Alfina Volcano (Fig. 2b) is located a few kilometers north of the Vulcini district (Fig. 2; 42° 45' 19"N-11° 56' 41"E) and consists of a few tiny lava flows erupted from a small volcanic centre, and a small neck cropping out North of Torre Alfina castle. Torre Alfina is a monogenetic volcano characterised mainly by lava flows and minor Strombolian activity (Conticelli, 1998). The Torre Alfina volcanic rocks rest directly on the sedimentary substratum of the Monte Cetona horst.

The Torre Alfina lava flows were erupted at the top of the hill and flowed partly into the canyon formed by the Paglia river. Field characteristics indicate that they had low viscosities, and some lava tunnels may be recognised in the eastern part of the vent. Two slightly different types of lava can be also recognised. They were emplaced contemporaneously. They have different colours, vesiculation, and porphyritic indexes. The most mafic lava (type-A, Fig. 2b) is dark grey, almost aphyric, and

contains abundant green ultramafic microxenoliths 2-4 cm in diameter (Conticelli & Peccerillo, 1990), as well as a few lower crustal xenoliths of variable size (5-10 cm) and nature (Orlando et al., 1994). Type-B lava is the most widely represented at Torre Alfina volcano. It is dark grey to light grey in color, with a variable porphyritic index and vesiculation, showing some flattened vesicles of a few centimeters in length. Type-B lavas contain abundant crustal xenoliths of variable size (2-30 cm), some large (3-20 cm) magmatic mica-rich inclusions, and rare green ultramafic xenoliths. Torre Alfina lavas are leucite free ultrapotassic rocks of lamproitic nature of ultimate mantle origin generated by low degrees of partial melting of a veined mantle wedge at low X_{CO_2} conditions (Conticelli et al., 2015a).

3.1.3. Radicofani monogenetic volcano

The Radicofani volcano is a monogenetic volcano lying within the central portion of the Radicofani basin, few kilometers North of the Torre Alfina volcano (Fig. 2b). The Radicofani basin represents the southernmost branch of the Siena-Radicofani graben, one of the most important post-orogenic extensional tectonic features of the Northern Apennine (e.g., Pasquarello et al., 1983; Acciella et al., 2002; Borini & Sani, 2002). The Radicofani monogenetic volcano is formed by a 90 m-high well-preserved volcanic neck, with remnants of a cinder cone and of a lava lake at its top, and by several lava flows (Fig. 3).

The lava flows are scattered around the neck and lie on top of the Pliocene marine sediments (Liotta, 1996; Disperati & Liotta, 1998; Ghinassi & Lazzarotto, 2005; Conticelli et al., 2011). The original volcanic edifice was made up by a cinder cone a few hundred meters high. Today only a thin layer of red scoriae is preserved on the edge of the top of the neck, just West of the homonym castle. The top of the neck is made up of red vesicular lava that becomes dense and grey downward (Fig. 3). Columnar jointing is present in the middle and lower portions of the neck in grey and dark grey aphyric to sub-aphyric lavas. Oxidized septa dividing the different portions of the neck can also be observed. Rocks of different portions of the volcano yield radiometric ages in the range between 1.315 Ma and 1.08 Ma (e.g., Barberi et al., 1971; Pasquarello et al., 1983; D'Orazio et al., 1991), but the uncertainty in the age determination is relevant.

The Radicofani volcanic rocks have significant compositional variations from basaltic andesite (high-K calc-alkaline), at the bottom of the neck to ultrapotassic shoshonite for the lava-lake lavas, scoria and lava flows. The overall volcanic rocks show aphyric to sub-aphyric textures with small euhedral phenocrysts of olivine set in a microcrystalline matrix made of clinopyroxene, sanidine, ilmenite, and magnetite. Rare microphenocrysts of orthopyroxene are found in the groundmass of calc-alkaline samples. Chemical and isotopic variations along the necks are thought to be a primary feature due to incremental partial melting of a veined mantle wedge under conditions of low X_{CO_2} , favouring the genesis of silica-saturated, leucite-free, ultrapotassic magmas (Conticelli et al., 2011).




Fig. 3 - (a) Scoria and lapilli of the cinder cone containing the lava lake at the root of the neck; (b) reddish shoshonitic lava forming the lava lake at the roof of the neck; (c) oxidized septum that separates shoshonitic lava of the upper neck from the high potassium basaltic andesitic lava of the lower neck (redrawn after Conticelli et al., 2011; Avanzinelli et al., 2022).

3.1.4. The Monte Amiata: a Quaternary "hybrid" volcano

The Monte Amiata volcano is a small linear Late Pleistocene volcano located some tens of kilometers North of the Vulsini district (Fig. 1) and few kilometers west of Radicofani monogenetic volcano (Figs. 2a and 4). It is found where the Roman and Tuscany magmatic provinces overlap in space but not in time (Conticelli et al., 2004, 2010).

The Monte Amiata volcano produced a sequence of several viscous lava flows and exogenous domes outpoured from NNE-SSW tectonic structures (Fig. 2a), both hosting abundant cognate magmatic enclaves (e.g., Ferrari et al., 1996; Marroni et al., 2015). The volume of magma outpoured is smaller than the amount of magma erupted in Roman volcanoes, but consistently larger than that produced by Tuscan volcanoes (Conticelli et al., 2009a; 2010). According to Laurenzi et al. (2015) the onset of magmatic activity was at 305 ka with final lavas emplaced at 230 ka. Lavas and enclaves are characterized by high-K calc-alkaline to shoshonitic affinities ranging in composition with time from trachydacites to trachyte, latite, and olivine latite, with a progressive increase of the amounts of mafic enclaves. Monte Amiata rocks, lavas and enclaves, are leucite-free, with phenocrysts of olivine, confined to the most mafic terms, biotite, plagioclase, sanidine, clinopyroxene and orthopyroxene, set in a microcrystalline to glassy matrix. Fine-grained magmatic enclaves range from porphyritic to aphyric and invariably display chilled margin texture, ranging in compositions from trachybasaltic to shoshonitic and latitic. Petrographic and compositional characteristics of fine-grained magmatic enclaves suggest that they were molten at the time of inclusion by the host trachytic magma. Their chemical, mineralogical, and isotopic characteristics are strongly suggestive of genesis by mixing/mingling between a resident trachydacitic mag-




Fig. 4 - Geological sketch map of the Amiata region. Legend: 1) sedimentary sequences of the Ligurid units; 2) Pliocene marine sediments; 3-4) Radicofani volcano, neck and diemembered lavas, respectively; 5-14) Monte Amiata Volcano - 5-6) Basal trachydacitic lava flows (6); 7-11) trachytic domes and lava flows; 12) iatic lava flows; 14) olivine iatic lava flows; 15) Quaternary alluvial deposits (redrawn after Conticelli et al., 2015a,b; Laurenzi et al., 2015; Maroni et al., 2015).

ma, derived by crystal fractionation from a high-K calc-alkaline basaltic-andesitic magma (Tuscan type), similar in composition to the Radicofani magma and a younger Roman-type, silica-undersaturated, basanitic magma (Conticelli et al., 2015b, and references therein).

3.1.5. The other Leucite-free volcanoes in Latium and Campania

Other leucite-free ultrapotassic associated to shoshonitic and calc-alkalic volcanic rocks of ultimate mantle origin were emplaced along the Tyrrhenian border of the Italian peninsula in a period of time between 2 and 1 Ma (Fig. 2) at Monte Cimino volcanic complex, in Northern Latium underlying the Vico Volcano (Conticelli et al., 2013), at Tolfa-Maziana-Cerite volcanic complexes, in Central Latium underlying the Sabatini volcanoes (Bertagnini et al., 1995), offshore in the Pontine archipelago, in Southern Latium (Conte & Dolfi, 2002; Cadoux et al., 2005), and in the Neapolitan area, in the Central Campania, beneath the Campi Flegrei volcanoes (Albini et al., 1980; Barbieri et al., 1979; Brocchini, 1999). In most of these localities the older leucite-free igneous rocks preceded by a few hundred thousand years the younger strongly silica-undersaturated, melilite- to leucite-bearing Roman volcanic rocks (Conticelli et al., 2010, 2015a).

3.2. Latium and northern Campania

(D.M.P., S.C., J.P.S.-B., M.G.B.G., G.G., F.M., G.S.)

3.2.1. General background

The Quaternary (Middle Pleistocene to present) ultrapotassic magmatism of Central Italy fits into the geo-dynamical context of the Tyrrhenian back-arc extension related to the NE retreat of the west-directed Adriatic slab (Malinverno & Ryan, 1988; Conticelli & Peccerillo, 1992; Faccenna et al., 2001a; Acocella & Funiciello, 2006). This originated a NW-SE trending volcanic belt extending between the Tyrrhenian Sea coast and the Apennine orogen from southern Tuscany, through Latium and Campania. Defined as the Roman Comagmatic Region since Washington (1906), this volcanic region is known worldwide for its peculiar potassio and strongly silica-undersaturated and melilite- to leucite-bearing ultrapotassio magma: feeder magmas display a wide compositional range and multiple differentiation trends, ranging from K-basalts to K-foidites, phonolites and trachytes, and even K-rhyolites. Based on isotope geochemistry, the Roman, Ernici-Roccamontina and Campanian districts are distinguished (Peccerillo, 2005, 2017; Conticelli et al., 2010).

Here, we focus on the Roman Province volcanoes, i.e., from NW to SE, Vulsini, Vico, Sabatini and Colli Albani, as well as on the Volsci Volcanic Field (formerly erroneously defined Ernici volcanoes) and Roccamontina volcano, which belong to the Ernici-Roccamontina districts (Fig. 5). These volcanoes show a full spectrum of




Fig. 5 - Location and geological sketch of the Quaternary peri-Tyrrhenian potassic volcanoes of central Italy.

eruptive dynamics and related morphologies, ranging from small-scale Strombolian and phreatomagmatic events from monogenetic centers (scoria cones, maar-tuff rings and tuff cones), with associated effusive activity, to highly explosive Plinian and pyroclastic flow-forming events, often related to caldera collapses. Major volcanic districts with a dominant areal character (i.e., Vulsini and Sabatini), characterized by multiple caldera depressions and networks of monogenetic eruptive centers, alternate with essentially central volcanoes, such as stratovolcanoes with summit calderas and subordinate eccentric activity (i.e., Vico and Roccamonfina), or polygenetic caldera volcanoes with ignimbrite plateaux (i.e., Vulcano Laziale in the Colli Albani district). In some cases, central volcanic edifices with summit calderas, such as Latra and Montefiascone (Vulsini), Sacrofano (Sabatini) and Faate (Colli Albani), superimposed in time and space with areal volcanism. Quite peculiarly, the Voleci Volcanic Field consists of small-scale, isolated or locally clustered, eruptive centers.

The onset of potassic volcanism is documented by the oldest exposed products in the Volsci Volcanic Field that date back to at least ca. 760 ka (Boari et al., 2009b; Cardellio et al., 2020; Marra et al., 2021) (Fig. 6). However, evidence of even earlier activity in the Roman area is provided by ca. 750-810 ka tephra layers found in the Tiber River delta (Marra et al., 2014; Fig. 6). Since ca. 600 ka, the Vulsini, Sabatini, Colli Albani, Volsci and Roccamonfina volcanoes were all simultaneously active (Fig. 6) (e.g., Marra et al., 2003; Sottili et al., 2004; Roucoul et al., 2008; Boari et al., 2009b; Conticelli et al., 2010; Palladino et al., 2010; Soligo & Tuccimei, 2010), followed at ca. 415 ka by the onset of volcanic activity at Vico (Pereira et al., 2020).

3.2.2. Vulsini Volcanic District (ca. 590-111 ka)

The northernmost and largest (ca. 2200 km²) volcanic district in the Roman Province developed along the southern termination of the NNW-SSE trending Miocene-Pliocene Siena-Radicofani Graben from the superposi-






Fig. 6 - Temporal distribution of the Middle Pleistocene volcanic activity from the peri-Tyrrhenian potassic volcanic systems, plotted against the LR04 benthic stack record (Lisicki & Raymo, 2006). Data source: Vulsini: Palladino et al. (2010); Marra et al. (2020a); Vico: Penni et al. (2004); Persira et al. (2020); Sabatini: Sottili et al. (2010); Marra et al. (2014, 2020b); Colli Albani: Marra et al. (2009); Volsci: Boari et al. (2009); Centamore et al. (2010); Marra et al. (2021); Roccamonfina: Giannetti (1998a, 1998b); Giannetti & De Casa (2000); Rouchon et al. (2008); Scialletti et al. (2008).

tion of five major volcanic complexes (or lithosomes), partially overlapping in space and time: Paleovulsini, Vulsini Fields (formerly defined as Southern Vulsini), Bolsena-Orvieto, Montefiascone and Latera (Vezzoli et al., 1987; Palladino et al., 2010 and reference therein; Figs. 7 and 8).

The most prominent volcano-tectonic feature in the district is the wide caldera depression hosting Lake Bolsena, which is intersected, respectively to the W and SE, by the Latera and Montefiascone central volcanoes, featuring collapse calderas (Nappi et al., 1991; Acciella et al., 2012). The whole area is dotted with several tens of monogenetic cones and craters, either isolated (e.g., Lagaccione maar, tuff cones of Bisentina and Martana Islands in Lake Bolsena) or clustered (e.g., the Valentano scoria cones on the SE rim of Latera caldera) or located along volcano-tectonic lineaments (e.g., the N-S scoria cone alignment along the eastern rim of Bolsena caldera).

Overall, volcanic activity spanned a broad spectrum of eruptive styles, intensities and magnitudes, ranging from small-scale explosive (Strombolian and hydromagmatic)

and effusive eruptions from monogenetic centers, up to Plinian and pyroclastic flow, caldera-forming, events (Sparks, 1975; Conticelli et al., 1987; Nappi et al., 1994; Palladino & Valentine, 1995; Palladino & Agosta, 1997; Palladino & Simei, 2005; Palladino et al., 2014). Figure 9 shows an example of interbedded volcanic units from different source vents and their unconformity boundaries. In the following, the five lithosomes are briefly described: 1) the Paleovulsini lithosome groups the early eruptive products of Vulsini (ca. 590 to <490 ka; Nicoletti et al., 1981; Vezzoli et al., 1987; Cioni et al., 1989; Nappi et al., 1995), which consist of trachytic welded pyroclastic-flow deposits (basal Ignimbrites or Nenfri, ca. 550-505 ka) and associated Plinian pumice fall horizons exposed at the periphery of the district on top of the sedimentary substrate. An early collapse caldera possibly related to the basal ignimbrites, inferred from the Bolsena lake bathymetry and deep drillings, controlled the location and style of the following eruptive activity, which followed as areal volcanism (Vulsini Fields lithosome); 2) the Vulsini Fields (former Southern Vulsini complex;



Towns and other localities: A=Acquapendente, B=Bolsena, Ba=Bagnoregio, C=Canino, F=Farnese, G=Grotte di Castro, M=Montefiascone, O=Orvieto, P=Pigliano, S=Sovana, Si=Sorano, T=Tuscania, TA=Torre Alfina, Cm=Capodimonte, LR=La Rocchetta, Ps=Piansano.

Fig. 7 - Geological sketch map of the Vulsini Volcanic District (modified after Palladino et al., 2010). The boundaries of the Sheet No. 344-Tuscania of the 1:50,000 Geological Map of Italy (CARG project) are also shown. Towns and other localities: A, Acquapendente; B, Bolsena; C, Canino; F, Farnese; G, Grotte di Castro; M, Montefiascone; O, Orvieto; P, Pigliano; S, Sovana; Si, Sorano; T, Tuscania; TA, Torre Alfina; Cm, Capodimonte; LR, La Rocchetta; Ps, Piansano.




Fig. 8 - Sketch of the legend of sheet 344-Tuscania (after Palladino et al. 2010, modified), where representative lithostratigraphic volcanic units are grouped into lithosome units and organized in the framework of the unconformity-bounded stratigraphic (UBS) units recently defined for the volcanic and Tyrrhenian coastal areas of northern Latium. Note that major UBS unit boundaries correlate to even-numbered oxygen isotope stages, corresponding to low-stands of sea level ($\delta^{18}\text{O}$ curve after Shackleton et al., 1990; Shackleton, 1995).

Vezzoli et al., 1987) is a composite lithosome consisting of a gently sloping volcanic plateau, extending throughout the whole district, which developed by long-lasting (ca. 490–111 ka) effusive and subordinate explosive activity from a network of scattered eruptive centers. Volcano-tectonic lineaments related to the early Paleovulsini caldera might have favoured the eruption of poorly differentiated, small magma batches from multiple source vents around the present Lake Bolsena depression. Although this areal volcanism largely preceded the onset of central explosive activities at Latena and Montefiascone at ca. 290 ka (Nappi et al., 1995; Brocchini et al., 2000), it was also contemporaneous and even followed the Montefiascone and Latena explosive climax until the waning phases of activity in the district. In particular, the Monte Bisenzio spatter cone repre-

sents the most recent volcanic product dated so far at Vulsini (ca. 111 ka; Marra et al., 2020a). In lack of evidence of major caldera-forming eruptions that could explain the size of the Bolsena depression, Walker (1984) and Cole et al. (2005) classified it as a downseep caldera end-member. The relationships among eruptive sources and volcano-tectonic lineaments may suggest that the prolonged magma drainage during the Vulsini Fields activity was accompanied by incremental caldera growth that extended the early Paleovulsini caldera collapse and eventually resulted into the wide depression hosting Lake Bolsena (Acocella et al., 2012);

- 3) the Bolsena-Orvieto lithosome includes lava and pyroclastic successions that are mainly exposed in the northeastern sector of Vulsini, from Lake Bolsena to the Tiber Valley, including a series of Plinian fall deposits




Fig. 9. Field pictures of the Vulsini volcanic successions. (a) La Rocchetta locality (6 km SW of Capodimonte): major pyroclastic units of Latara (i.e., Canino, Sorano, Grotte di Castro formations) on top of a lava flow from the Vulsini Fields and intervening stratigraphic discontinuities, such as paleosols (e.g., D1, p) and high-relief erosional surfaces (e.g., D2, D3, D4), related to UBS boundaries of different rank (see Fig. 8). Note the U-shaped channel (D2) mantled by a thin ash-fall layer and filled by pyroclastic current deposits of the Sorano formation (flow direction toward the page; d-example of flow unit boundary). (b) 5 km SW of Montefiascone: interbedded Vulsini Fields and Montefiascone volcanic successions, showing the slightly undulating erosional surface bounding the Barca di Parma (BP) and Fiume Ficra (FF) synths, and local stratigraphic discontinuities representative of temporal hiatuses in the eruptive activity, as evidenced by steep erosional channels (black arrow) or paleosols (p). (c) Close-up of the upper part of the Canino formation at La Rocchetta, showing the depositional boundary between temporally closely spaced pyroclastic flow and fall units, topped by the Canino-Fameise inter-eruptive paleosol.




Fig. 10 - (a) Geological sketch map of the Cimino Volcanic Complex (Tuscan Magmatic Province, Avanzinelli et al., 2009; Conticelli et al., 2010, 2015a). Legend: 1) sedimentary substrate; 2) domes and lava flows of the I Cimino phase; 3) pyroclastic flows of II Cimino phase; 4) leitic to trachytic lava flows of the III Cimino phase; 5) olivine leitic to shoshonitic lava flows of the IV Cimino phase; 6) Roman-type Leucite-bearing Vico volcanic rocks (Roman Magmatic Province); 7) quaternary fluvial deposits (redrawn after Parini et al., 1997, 2004; Conticelli et al., 2007, 2010; 2013, 2015a). (b) Geological sketch map the Vico volcano. Legend: E) caldera rim; 8) Macigno Flysch; 9) Allochthonous Flysch; 10) post-orogeny Pliocene-Pleistocene marine sediments; 11-13) Cimino dome complex (1300-900 ka); 11) lamproitic lava; 12) rhodacitic domes; 13) rhodacitic Peperino Tipico ignimbrite; 16-14) Lago di Vico synthem (second period; the earlier Rio Farina Synthem is not representable at this map scale); 14) stratocone building leucite-bearing lavas; 15) caldera forming phonotitic ignimbrite; 16) Carbognano phreatomagmatic ignimbrite; 17) Monte Venere lithosome; post-caldera scoria cones (third period); 18) travertine; 19) Holocene alluvial and lacustrine deposits; (c) Geological sketch section through the Monte Cimino (1.3-0.9 Ma) and Vico (0.45-0.05 Ma) nested volcanoes (redrawn after Conticelli et al., 2013; Avanzinelli et al., 2017).

(Nappi et al., 1994, 1995) and the interbedded ca. 332 ka Orvieto-Bagnoregio Ignimbrite related to a major caldera-forming event (Palladino et al., 2014; Marra et al., 2019; Palladino & Pettini, 2020);

4) the Monteriascone lithosome, SE of Lake Bolsena, consists of a central volcanic edifice cut by a nearly circular polygenetic caldera, ca. 3 km across. The onset of activity (<286 ka), broadly contemporaneous with Latara, is represented by the lithic-rich Monteriascone basal ignimbrite (Nappi & Marini, 1986), a hydromagmatic event related to an early stage of caldera collapse;

5) Latara is a gently sloping, central volcanic edifice (at least 30 km across) located in the western part of

Vulsini (Fig. 9). Its climactic activity (ca. 253-177 ka; Monaco et al., 2022a and reference therein) produced widespread Plinian fall (Palladino & Agosta, 1997) and ash-pumice flow deposits (Sparks, 1975; Palladino & Valentine, 1995; Palladino & Giordano, 2019; Valentine et al., 2019), resulting in the Latera-Vepe nested caldera system (9 km by 7 km across) through multiple collapse stages (e.g., Nappi, 1969; Sparks, 1975; Barberi et al., 1984; Vazzoli et al., 1987; Nappi et al., 1991; Palladino & Simeoli, 2005). In particular, the Sovana and Onano eruptions, which emplaced typical lag breccia deposits, are identified as major caldera-forming events (Palladino & Simeoli, 2005; Palladino et al., 2014). The late volcanic activity was characterized by effusive (e.g.,




Fig. 11 -Variation of the silica content upsection within the Vico α and Vico β Plinian deposits and the minor pre-Vico α and post-Vico β units (modified from Pereira et al., 2020).

the ca. 157 ka shoshonitic Selva del Lamone lava plateau), Strombolian and phreatomagmatic eruptions from intra- and peri-caldera monogenetic centers.

The Vulcini volcanoes span a full range of potassic rock types, from the least differentiated (i.e., K-basalts,

trachybasalts, basanites, and tephrites), which mostly characterize lava flows and plateaus, as well as Strombolian and hydromagmatic products, to the most differentiated ones (i.e., phonolites, and trachytes), typical of Plinian pumice-fall and ash-pumice flow deposits from major

explosive eruptions (Conticelli et al., 1987, 1991; Nappi et al., 1994; Palladino & Agosta, 1997; Palladino et al., 2010, 2014 and reference therein). In particular, the Montefiascone lavas (e.g., from the circum-caldera Orto Platano eruptive center; ca. 226 ka; Nappi et al., 1995) display peculiar compositional features (i.e., high CaO and MgO and low SiO₂, Al₂O₃, and alkali contents) and are recognized among the most primitive rock types in the Vulsini district and in the whole Roman Province as well (Conte et al., 2009).

In distal setting, the activity of the Vulsini Volcanic District is particularly well documented for the Latara stage (ca. 255–157 ka) in Fucino Basin, central Italy (Monaco et al., 2021). From the same basin, several tephra layers originated from Vulsini have been also found in the interval 430–300 ka (e.g., Orvieto-Bagnoregio unit of 335 ka; Leicher et al., 2023; Monaco et al., 2021), while activity of the Paleovulsini is possibly documented in Lake Ohrid (Leicher et al., 2018; 2021).

3.2.3. Cimino (ca. 1.1–0.8 Ma) and Vico (ca. 415–95 ka)

On the basis of morphological, geological, and mineralogical data the Cimino and Vico volcanoes (Fig. 10) represent two distinct edifices active in different times, but fed by the same plumbing system (e.g., Perini, 1997; Perini et al., 1997, 2000, 2003, 2004; Peccerillo, 2005; Conticelli et al., 2010, 2013, 2015a;). According to Peccerillo et al. (1987) the Cimino volcanic complex, active during the Early Pleistocene, belongs to the Tuscan Magmatic Province, whilst the nested and younger Vico stratovolcano, active during the Late Pleistocene, belongs to the Roman Magmatic Province.

The Cimino volcanic complex is the oldest edifice of the Cimino-Vico area, and it is a relatively simple volcano with an evolution through four main phases (Conticelli et al., 2013) in a short time span from 1.35±0.08 to 0.94±0.20 Ma (e.g., Everenden & Curtis, 1965; Nicoletti, 1969; Borghetti et al., 1981).

The Cimino volcano is made up of: i) an initial alignment of trachydacitic to trachytic cryptodomes along a NW-SE trend (author, "peperino delle atture"); ii) two welded pyroclastic flow units, outpoured when the activity was concentrated in the central part of the volcano (author, "peperino tipico"); iii) lava flows of the Monte Cimino cone; iv) final mafic lava flows unit (e.g., Puxeddu, 1971; Lardini & Nappi, 1987; Perini et al., 2003; Cimarelli & de Rita, 2006a,b; LaBerge et al., 2006, 2008; Conticelli et al., 2013; Nappi et al., 2022).

The Cimino volcanic rocks are invariably leucite-free ranging with time from trachydacite, to trachyte, latite, olivine-latite, and euhedralites. Details on mineralogical and petrographic features are provided in Perini et al. (2003) and Conticelli et al. (2013).

After a hiatus of some hundreds of thousands of years, renewal of volcanic activity occurred in the area, at 450 ka, few kilometers southwest of the Monte Cimino with eruption of Roman-type, silica-under-saturated, leucite-bearing magmas (Fig. 10). The Vico volcano was built in the form of a conic stratovolcano cut by a summit polygenetic caldera (Perini et al., 1997, 2004). Post-caldera activity, subordinate in volume, both within and on the N and W edges of the caldera do occur (Fig. 10b; Perini & Conticelli, 2002; Perini et al., 1997, 2004). Vico volcanic rocks were strong to mild silica-undersaturated,

leucite- to plagioclase-leucite-bearing generated by partial melting of a lithospheric mantle wedge under high X_{CaO} conditions.

The eruptive history of the Vico volcano is subdivided into three periods and corresponding lithosomes (Perini et al., 2004; Conticelli et al., 2013):

- 1) Rio Ferriera Synthem (sensu Perini et al., 2004), involving an early effusive period, alternating with explosive phases at ca. 415–400 ka that produced the Vico α and Vico β Plinian fall markers (Cloni et al., 1987; Pereira et al., 2020);
- 2) Lago di Vico Synthem, involving the build-up of the main stratovolcano by essentially effusive activity (ca. 305–258 ka), followed by a climactic, caldera-forming, explosive phase at ca. 250–144 ka that produced Ignimbrites A, B, C (also known as Tufo Rosso a Scorie Nere Vicano) and D (Locardi, 1965; Bertagnini & Sbrana, 1996), corresponding to the Farine, Ronciglione, Sutri and Carbognano formations (Perini et al., 2004; Bear et al., 2009a,b);
- 3) Monte Venere Synthem (ca. 138–95 ka), characterized by phreatomagmatic and Strombolian-effusive activities from peri- and intra-caldera monogenetic centers.

Vico α and Vico β are geochemically peculiar within the peri-Tyrrhenian volcanism. In fact, in addition to the usual trachy-phonolitic ones, the juvenile fraction of the deposits of both eruptions contains rhyolitic pumice clasts (Fig. 11), which represent a puzzling exception among the Quaternary potassic volcanoes of central Italy. Isotopic and trace element data indicate an origin of these rhyolitic magmas by fractional crystallization processes (Perini et al., 2004), occurred at unusually high-pressure conditions: the rhyolitic glasses in pumice clasts show, indeed, high volatile contents (H₂O ca. 5–7 wt%) and are in equilibrium with amphibole (Monaco, 2022). The deep level of the Vico α and Vico β pre-eruptive magma systems is consistent with the wide dispersal of the fallout products and the lack of co-eruptive pyroclastic currents associated with caldera-forming processes.

Conversely, the Tufo Rosso a Scorie Nere Vicano (ca. 150 ka; Laurenzi & Villa, 1987) is related to the main caldera collapse at Vico (Bear et al., 2009a,b; Palladino et al., 2014). The eruption succession includes: i) a basal Plinian fall horizon containing whitish sub-aphyratic pumice, related to a central feeder conduit; ii) black spatter-bearing, coarse lithic-rich, co-ignimbrite lag breccias, related to developing ring faults during caldera collapse; iii) red tuff with leucite-bearing black scoria, that is the main pyroclastic flow deposit emplaced at the eruption climax, extending with remarkable continuity as far as 25 km from the vent, over an area of 1200 km². The facies architecture and the variation patterns of juvenile textural features and glass compositions, along with mineral chemistry, reveal a magma chamber characterized by thermal and volatiles gradients (which reflect in crystal concentration gradients) and changing magma withdrawal dynamics in the course of a typical caldera-forming event (Bear et al., 2009a; Palladino et al., 2014).

Owing to their peculiar rhyolitic components and/or wide dispersal, Vico α and Vico β and the Vico Ignimbrites C and B are key stratigraphic markers for regional and ultra-regional correlations (Cloni et al., 1987; Pereira et al., 2020; Monaco et al., 2021; Iurino et al., 2022), as




Fig. 12 - Sketch geological map of the Sabatini Volcanic District (after Scattili et al., 2010a), showing the areal distributions of the main activity periods. Insets show the locations of the Sabatini and the other volcanic districts of the Roman province, central Italy. Legend: 1- Sedimentary terrains; 2-M.ti Certi-Toffetano-Manziale lava domes (Pliocene); 3- Morlupo phase products (0.6-0.5 Ma); 4- Southern Sabatini products (0.5-0.4 Ma); 5- Bracciano Caldera products (0.3-0.2 Ma); 6- Sacrofano Caldera products (0.3-0.2 Ma); 7- Vico Volcano (0.4-0.1 Ma); 8- Sabatini Late activity (< 0.15 Ma); 9- Scoria cones; 10- Meas; 11- Caldera rims; 12- Buried source areas.

far as Lake Ohrid, Albania-North Macedonia (Leicher et al., 2021).

3.2.4. Sabatini Volcanic District (ca. 810-70 ka)

The Sabatini Volcanic District extends over an area of ca. 1,800 km² north of Rome (Fig. 12). Volcanism was dominated by explosive activity ranging through a broad spectrum of intensities and magnitudes, from Strombolian and phreatomagmatic eruptions from monogenetic centers to Plinian and pyroclastic flow-forming events associated with caldera collapses (Scattili et al., 2004, 2010a; Marra et al., 2014, 2016). Today, the main landforms in the area are represented by the Bracciano caldera, hosting a lake ca. 9 km in diameter, and by the 6 x 4 km Sacrofano caldera to the East. Although quite subordinate volumetrically, a number of monogenetic tuff rings and scoria cones that postdate the main caldera-forming events are either clustered or scattered in the northern area of the district and between the Bracciano and Sacrofano calderas (Scattili et al., 2010a; Valentine et al., 2015).

Eruptive activity mostly took place in the 0.6-0.1 Ma time span, although older widespread Plinian fall deposits of uncertain source (yet tentatively attributable to a Sabatini paleo-activity) were also found in drill sites along the Tiber Valley and dated at ca. 810-814 ka (Kärner et al., 2001). Conventionally, the Sabatini activity is divided into three main phases (Cioni et al., 1987, 1993; de Rita

et al., 1993, 1996; Barberi et al., 1994; Conticelli et al., 1997; Kärner et al., 2001; Scattili et al., 2004, 2010, 2012, 2019; Mascotta et al., 2010):

- 1) early phase from the Morlupo (ca. 589-500 ka) and Southern Sabatini (ca. 500-400 ka) source areas, respectively to the East and South of the present Lake Bracciano, characterized by widespread pyroclastic flows and Subplinian to Plinian fallout (VEI 3-5). During this phase of activity, the main eruptions of the ca. 548 ka Tufo Giallo della Via Tiberina and the ca. 449 ka Tufo Rosso a Scorie Nere Sabatino occurred, the latter being the volumetrically most important caldera-forming event in the whole activity history of Sabatini (Scattili et al., 2004; 2010; Mascotta et al., 2010; Marra et al., 2014; Palladino et al., 2014);
- 2) intermediate phase of effusive and explosive activity (ca. 400-200 ka) from Lake Bracciano and Sacrofano source areas, marked by two major caldera-forming eruptions, respectively the ca. 310 ka Tufo di Bracciano and the ca. 286 ka Tufo Giallo di Sacrofano;
- 3) late eruptive phase (ca. 150-70 ka), dominated by phreatomagmatic and subordinate Strombolian and effusive activities from monogenetic and polygenetic tuff rings and scoria cones in the central area of Sabatini (Scattili et al., 2012; Valentine et al., 2015), mostly between the Bracciano and Sacrofano calderas (e.g., Baccano, Martignano and Straciacappa centers).

Compositionally, the Sabatini volcanics cover a

wide spectrum of potassic rock types, ranging from trachybasalts to trachytes and phonolites (Conticelli et al., 1997, 2010; Peccerillo, 2005; Sottili et al., 2019). Marra et al. (2020b) highlighted a noticeable parallelism in time and eruption magnitude between Sabatini and Colli Albani during most of their lifetime, and suggested a common regional tectonic trigger, such to define them twin volcanic systems.

The explosive activity of the Sabatini Volcanic District is well documented in distal setting. The most widespread unit is Fall A (~500 ka) of the Morlupo phase, found in central-southern Italy intermountain basins (Sulmona, Mercure and Acerino; Giaccio et al., 2014; Petrosino et al., 2014; Di Rita & Sottili, 2019) and Lake Ohrid (Leicher et al., 2021). Many other tephra layers spanning the interval 550–120 ka, of either well identified or supposed Sabatini origin, have been found in Fucino and Acemo basins, Lake Monticchio and even Lake Ohrid (Wulf et al., 2012; Giaccio et al., 2012; 2017a, 2019; Petrosino et al., 2014; Monaco et al., 2021; 2022a; Leicher et al., 2023). Tephra layers with Sr isotope composition and age comparable to the putative Sabatini paleo-activity (~615–810 ka) have been found in the Sulmona basin (Giaccio et al., 2013b; 2015; Sottili et al., 2019), although their attribution to the Sabatini Volcanic District is still uncertain.

3.2.5. Colli Albani Volcanic District (ca. 608–36 ka)

Following a poorly documented early activity period, the Colli Albani volcanism took place in three main phases marked by different eruptive mechanisms and magma volumes (de Rita et al., 1988, 1995b; Giordano et al., 2006; Fig. 13):


- 1) the climactic activity period, known as Tuscolano-Artemisio Phase (de Rita et al., 1988) or, referring to the evolution of the edifice, as Vulcano Laziale caldera complex (Giordano & the CARG Team, 2010), produced at least five subsequent eruptive cycles, dominated by large mafic and caldera-forming ignimbrites, in the ca. 561–351 ka time span (Kamer et al., 2001; Marra et al., 2009), alternated with effusive and mild explosive activity. Strikingly, large-volume (up to several tens of km³) pyroclastic flows were fed by K-foiditic magmas with SiO₂ as low as 42 wt%, e.g., during the early Trigoria-Tor de' Cenci and Palatino "pisolithic tuff" eruptions and the main ca. 456 ka Pozzolane Rosse and ca. 365 ka Villa Senni caldera-forming eruptions (Freda et al., 1997, 2011; Palladino et al., 2001; Boari et al., 2009a; Marra et al., 2009; Vinkler et al., 2012; Gaeta et al., 2016);
- 2) the Faete Phase (de Rita et al., 1988), during which the Tuscolano-Artemisio peri-caldera ring fracture system and the intra-caldera Faete stratovolcano were built (Giordano & the CARG Team, 2010), occurred at ca. 308–250 ka (Marra et al., 2003) and was characterized mainly by Strombolian and effusive activities;
- 3) following a ca. 50 kyr-long dormancy, the Late Hydromagmatic Phase (ca. 200–36 ka; Marra et al., 2003, 2016; Freda et al., 2006; Giaccio et al., 2007, 2009), or Via dei Laghi lithosome (Giordano & the CARG Team, 2010), was dominated by maar and tuff ring forming eruptions (e.g., Ariccia, Nemi, Valle Marciana, Laghetto or Giurina, Albano). In particular, the Albano polygenetic maar (ca. 69–36 ka) hosted the most recent and

energetic activity (Giordano et al. 2002; Freda et al., 2006; Giaccio et al., 2007; De Benedetti et al., 2006).

The age of the last eruption at Colli Albani has been a matter of debate (see the above references and Soligo & Tuccimei, 2010 for a review). A wide data-set of accurate ⁴⁰Ar/³⁹Ar dating consistently indicates that the last documented eruptive event occurred at the Albano maar at ca. 36 ka (Marra et al., 2003; Giaccio et al., 2007, 2009), in spite of younger ages reported in the literature (e.g., ca. 26 ka; Villa et al., 1999) that are affected by stratigraphic inconsistencies and/or methodological uncertainties. In addition, a series of lahar deposits (Tavolato Formation) have been attributed to the repeated overflows of the Lake Albano and possibly associated with intrusion-triggered limnic eruptions (Funiciello et al., 2003; De Benedetti et al., 2008), although the occurrence and exact age of these events, possibly extending into the Holocene, has been a matter of controversy (Marra & Kamer, 2005; Giaccio et al., 2007; D'Ambrosio et al., 2010). On these grounds, by considering the long-lasting quiescence periods occurred in the past (ca. 40 kyr; Marra et al. 2004), it cannot be excluded that Colli Albani can be potentially active (de Rita et al., 1995a; Marra et al., 2016), also in light of the several geophysical (Chiara et al., 1997) and geochemical (Carapezza et al., 2010) indicators of volcanic unrest.

Although the Colli Albani rock-types span the tephrite, phonotephrite and K-foidite fields of the TAS classification diagram, they show invariably, regardless of eruption size, a leucitic mineral assemblage. The latter is characterized by the virtual absence of feldspars, with a few exceptions in the groundmass of olivine-bearing lavas, and in glass-bearing juvenile pyroclasts formed at high cooling rates (Triglia et al., 1995; Gaeta, 1998; Gaeta & Freda, 2001; Gaeta et al., 2006, 2016; Giordano et al., 2006; Boari et al., 2009a; Marra et al., 2009; Gozzi et al., 2014). The peculiar magma differentiation trend, leading to K-foiditic magmas with SiO₂ contents <50 wt%, has been explained by the assimilation of sedimentary carbonate. This process allows the establishment of CO₂-rich conditions in the Colli Albani magmas and reduces the stability fields of olivine and phlogopite in favour of clinopyroxene and leucite (Gaeta et al., 2000, 2009, 2021; Freda et al., 2006; Di Rocco et al., 2012). The anomalous highly explosive behaviour of SiO₂-poor magmas at Colli Albani has also been attributed to CO₂ entrainment in the pre-eruptive system due to magma-carbonate wall rock interaction at supra-crustal levels (Freda et al., 1997, 2011; Sottili et al., 2010b; Cross et al., 2014). An alternative hypothesis suggests that the main location for magma-carbonate fluids interaction, controlling both the peculiar line of descent and the explosivity, is within the metasomatized mantle source (Boari et al., 2009a; Jorgenson et al., 2020).

The uncommon lithology and K-foiditic composition and other peculiar features of the Colli Albani tephra (e.g., the ⁸⁷Sr/⁸⁶Sr time-dependent variability; Gaeta et al., 2016) make them among the most distinctive markers in the framework of the central Mediterranean tephrostratigraphy (Giaccio et al., 2013a). The pyroclastic products of the main activity phase of Colli Albani (ca. 561–351 ka), as well as of the Late Hydromagmatic Phase (ca. 200–36 ka), are widely dispersed in central-southern Italy (Giaccio et al., 2009; 2013; 2017a; 2019; Petrosino et al.,



(1) Koenig et al., 2001; (2) Marra et al., 2003; (3) Frusta et al., 2006; (4) Marra et al., 2009; (5) Gaeta et al., 2011; (6) Marra et al., 2016; (7) Gaeta et al., 2016.

Fig. 13 - Stratigraphic-chronological scheme and geological sketch of the Colli Albani (Alban Hills) Volcanic District (Marra et al., 2011; Gaeta et al., 2016).

2014; Monaco et al., 2021). The largest events, such as the ~456 ka Pozzolane Rosse, can be traced in the tephra record as far as Lake Ohrid (Leicher et al., 2019; 2021).

3.2.6. Volsci Volcanic Field (ca. 780-230? ka)

This volcanic field (also known as Monti Emici or Middle Latin Valley volcanoes) groups several monogenetic centers (e.g., scoria cones, maar-diatremes and tuff rings) scattered through the Middle Latin Valley and the Volsci carbonate range, as far as the adjoining Pontina Plain (Pasquarè et al., 1985; Boari et al., 2009b; Centamore et al., 2010; Cardello et al., 2020; Marra et al., 2021 and reference therein; Fig. 14). The Volsci eruptive pattern was characterized by small volume (in the order of 0.01-0.1 km³) eruptions from a network of monogenetic centers, totaling nearly 4 km³, i.e., two orders of magnitude lower than the major potassic volcanic districts of the Roman Province. Three main phases of activity have been distinguished (Marra et al., 2021), as follows:

- 1) the "Early Eruptive Phase" (ca. 761-541 ka) mainly occurred along the carbonate Volsci Range (e.g., Patria eruptive center) and was essentially fed by K-rich, leucite-millitite-bearing (HKS) magmas, showing evi-

dence of high degree of carbonate assimilation, possibly due to multi-stage ascent through thick carbonate successions (hence the diffuse phreatomagmatic character);

- 2) subsequently, during the "Main Eruptive Phase" (ca. 424-349 ka), HKS rock types alternated with poorly differentiated, plagioclase-bearing (KS) rock types from eruptive sources close in time and space (e.g., in the Ceccano and Poli areas). In particular, the eruptive activity along the major faults bounding the Middle Latin Valley Graben testifies for the fast ascent of nearly primary KS magmas (K-basaltic in composition; Centamore et al., 2010);
- 3) during the "Late Eruptive Phase" (ca. 300-231? ka), KS magmas prevailed, although still accompanied by HKS events. The low-intensity and low-volume (i.e., Hawaiian-Strombolian, phreatomagmatic, and subordinate effusive) eruptive events, fed by poorly differentiated potassic magmas, reflect the tectonically controlled, fast ascent of primitive magma batches from the mantle source ("bullet eruptions"; Cardello et al., 2020).

Owing to the lack of glass chemical compositions from the Volsci pyroclastic deposits, identifying potential distal correlatives is hardly feasible. Nevertheless, the




Fig. 14 - Geological sketch map of southern Latium, showing the location of the Volsci Volcanic Field (square area) between the major Colli Albani and Roccamontfina volcanoes (after Marra et al., 2021, modified).

Volsci is currently the sole volcanic source for which an activity as old as 760 ka is documented in near-vent setting, which is geochronologically compatible with a series of Sulmona tephra layers dated between ~805 and ~715 ka (Giaccio et al., 2013b; 2015).

3.2.7. Roccamontfina Volcano (630-557 ka)

Roccamontfina volcano (Fig. 15) is the result of the superposition of different morphostructures. The oldest edifice is a stratovolcano made up mostly of leucite-bearing, highly silica-undersaturated (High-K series; HKS; Conticelli et al., 2009b), phonotephritic lava flows and Strombolian deposits (Roccamontfina Synthem, de Rita & Giordano, 1996). An age of 630 ka was obtained for the pyroclastic deposits at the very base of the volcanic pile sampled from a deep borehole drilled at the center of the present caldera (Ballini et al., 1989a). Some K-Ar ages as old as ~1.2 Ma (Gasparini & Adams, 1967) and ~700 ka (Radicati di Brozolo et al., 1988) reported for the products of this early stage may be not reliable for modern analytical standards. As a result of NE-trending extensional faults cutting the edifice, a major lateral sector collapse interrupted the growth of the stratovolcano (de Rita & Giordano, 1996) at around ~440 ka (end of Rio Rava stage; Chiesa et al., 1995; Giannetti, 2001; Rouchon et al., 2008).

Subsequent activity (related to the Riardo Synthem; de Rita & Giordano, 1996) shifted to dominantly explo-

sive and a series of VEI 3-5 Subplinian to Plinian eruptions shaped a polygenetic summit caldera, superimposed to the earlier sector collapse (Cole et al., 1993; de Rita & Giordano, 1996; Giannetti, 2001). Rouchon et al. (2008) subdivided this caldera-forming stage into several substages. The oldest one, between 410 ka and 380 ka, culminated with the first caldera-forming event of the Brown Leucitic Tuff (BLT; Luhr & Giannetti, 1967; Cole et al., 1993), which is the last eruption involving HKS magmas at Roccamontfina. The following activity was fed by slightly silica-saturated leucite-free magmas with a shoshonite affinity (SHO; Conticelli et al., 2009b). Between ~331 ka and ~230 ka, the White Trachytic Tuffs (WTTs; Ballini et al., 1989b; Cole et al., 1993; Giordano, 1998a,b; Giannetti & De Cesa, 2000) originated from multiple trachytic Plinian eruptions. At least two of them (i.e., WTT Cupa and WTT Galuccio of Giordano, 1998a, or LWTT and UWTT of Giannetti & De Cesa, 2000) resulted into caldera collapses in the central area of the volcano. The pyroclastic deposits of the Yellow Trachytic Tuff (Rouchon et al., 2008) were also emplaced during this period.

The youngest activity at Roccamontfina (Vezzara Synthem; de Rita & Giordano, 1996; or 170-150 ka Santa Croce stage; Rouchon et al., 2008), was controlled by N-trending tectonic lineaments and gave birth to the intra-caldera latitic domes of Lattani and M.te Santa Croce. Eruptive activity ended at ~150 ka according to Rouchon




Fig. 15 - Sketch geological map of the Roccamonfina Volcano showing the main volcanic units grouped into the respective synths (redrawn after Cicali et al., 2010).

et al. (2008), though a younger age of ~55 ka is reported in Radicati di Brozolo et al. (1988).

In distal setting, tephra layers attributed to the Roccamonfina activity have been recognized on regional and extra-regional scales. Rio Rava is documented in Molise region, immediately east of Roccamonfina volcano (Russo Ermoli et al., 2010), while BLT has been recently identified in the Fucino basin (central Italy) and associated to the widespread "Bag tephra" of Danubian Middle Pleistocene loess succession (Leicher et al., 2023). The LWTT is also well documented in Molise alluvial-lacustrine successions (Amato et al., 2014; Galli et al., 2017) and, more distally, in the Fucino basin (Leicher et al., 2023). Of note, distal records, spanning the 600-200 ka interval, indicate the occurrence of a number of tephra layers with Roccamonfina-like geochemical compositions that currently lack specific proximal counterparts (e.g., Giaccio et al., 2014; Petrosino et al., 2014; Leicher et al., 2021), providing evidence for a conspicuous, yet still not well-defined, Middle Pleistocene explosive activity at Roccamonfina volcano.

3.2.8. Final remarks: the state of volcanic activity

Ongoing research aims at refining the chronology of volcanic activity in central Italy, in order to better constrain the eruption frequency and the duration of quiescence periods, which is crucial for long-term hazard assessment. In this regard, tephra correlations from distal archives provide additional, detailed information for a comprehensive reconstruction of tempo and dynamics of the peri-Tyrrhenian Quaternary volcanoes (e.g., Giaccio et al., 2019; Leicher et al., 2019; Monaco et al., 2021, 2022a).

Based on the ages of the last documented eruptive

events in the Roman Province, i.e., ca. 111 ka at Vulsini, ca. 95 ka at Vico, ca. 70 ka at Sabatini and 36 ka at Colli Albani (the latest activity at Vico and Roccamonfina is not yet well constrained), it appears that the waning (or, possibly, cessation) of eruptive activity shifted progressively from NW to SE. Strikingly, these volcanoes reveal a "relay behavior" (Marra et al., 2020a), i.e.: the waning stage of activity at Vulsini (ca. 150-110 ka) was concomitant to an activity climax at Vico (third period, Perini et al., 2004; also including the main caldera-forming event of Tufo Rosso a Scoria Nera Vicano); then, waning activity at Vico (ca. 95 ka) occurred during a peak of phreatomagmatic activity at Sabatini; again, the last documented eruption at Sabatini (ca. 70 ka) was accompanied by the onset of the Albano maar cycle at Colli Albani. Finally, the last eruptive phase at Albano (40-36 ka) was broadly coincident with the climactic super-eruption of the 40 ka Campanian Ignimbrite at Campi Flegrei. The focus of volcanic activity seems therefore to have shifted through time from Northern Latium, through the Roman area, toward the Neapolitan area. However, considering that the time lapse since the last documented eruptions at Sabatini (ca. 70 kyr) and Colli Albani (ca. 36 kyr), is in the same order of the dormancies occurred in the activity history of the two volcanoes (ca. 70 kyr and ca. 40 kyr, respectively), the possible occurrence in the near future of eruptive codas from the dormant "twin" volcanoes at the gates of Rome cannot be ruled out (Marra et al., 2020b).

In addition, a recent work by Giordano & Caricchi (2022) points out that defining the state of activity of volcanoes based on recurrence times alone requires the assumption that the magma plumbing system is steady-state, while even at constant magma fluxes, the thermal

and geometric configurations and chemical reactivity of plumbing systems may change significantly over time. This means that as they thermally mature, plumbing systems may allow the growth and coalescence of progressively larger magma bodies, implying progressively a longer quiescence, which may lead to larger eruptions. In this sense, refining our knowledge of the longevity of a volcano is as important as knowing the age of its last eruption, and this is why recent efforts in extending back the distal tephrochronology of the peri-Tyrrhenian Quaternary volcanoes is so important (e.g., Giacconi et al., 2019; Leicher et al., 2019; Monaco et al., 2021). At the same time Giordano & Cariocchi (2022) pointed out that the heat flux of the Roman volcanoes, especially at Vulcini and Sabatini is as high as, and locally even higher than, at the certainly active Ischia and Campi Flegrei volcanoes and much higher than at Vesuvius, suggesting that in order to define the state of activity of these volcanoes a much better imaging of their present magma plumbing system is required, although again none should be considered extinct.

4. PENINSULAR-INSULAR SOUTHERN ITALY VOLCANOES AND SEAMOUNTS

4.1. The Pontine Archipelago (A.M.C., C.P.)

The Pontine Archipelago consists of five major volcanic islands divided in western islands, Ponza, Palmarola and Zannone and eastern islands, Ventotene and S. Stefano (Fig. 16a). The western islands lie on a NE-SW elongated structural high dividing two major areas of sedimentation i.e., the Palmarola and Ventotene intra-slope basins (Fig. 16a) created by the Plio-Pleistocene extensional deformations, which also gave rise to a very steep NW-SE trending continental slope and an intense magmatic activity developed from late Pliocene to Late Pleistocene (Conte & Dolli, 2002; Cadoux et al., 2005). The eastern islands are the emerged portions of the caldera rim of a large strato-volcano rising ca. 700 m from the sea-floor emplaced at the center of the subsiding Ventotene basin (Fig. 16a; Metrich et al., 1988; Bellucci et al., 1999).

In the western islands the Pliocene volcanic cycle (4.5-2.9 Ma; Cadoux et al., 2005; Fig. 17a) produced a large effusion of High-K, Calc-Alkaline (HKCA) rhyolite lavas from extensional fissures in a submarine environment. This led to the emplacement of differently textured hyaloclastites and lava dykes which formed the typical cryptodome-dyke systems that constitute most part of Ponza Island (Fig. 16b, c).

Indeed, three main coalescing domes of about 1 km radius, centered at Cala M.te Pagliaro and aligned along a NE trending regional fracture, have been identified at Ponza on a morphological basis (de Rita et al., 2001). The same likely occurred in the submerged SW sector of Palmarola (Conte et al., 2016; Fig. 16b). At Zannone, as a consequence of local eustatic movements, rhyolite magmas similar, and probably coeval to those of Ponza (Conte et al., 2016), were emplaced as cryptodomes in a subaqueous/subaerial environment and also intruded into a substrate made up of sedimentary and metamorphic units, which are locally exposed (de Rita et al., 1996). This prevented the fine white hyaloclastite to form and led to the formation of a strongly brecciated lava in which

flow structures are often recognized. Similar brecciated lava facies are observed in the NW sector of Ponza island, including the islet of Gavi, suggesting a similar subaqueous/subaerial environment of emplacement.

The second volcano cycle started in Early Pleistocene age (1.64-1.52 Ma; Fig. 17a) during which Palmarola was entirely built owing to the emplacement of a submarine hyaloclastite unit intruded by dykes ranging in compositions from near- to peralkaline trachytes and rhyolites belonging to the Transitional Rock-series (TR; Conte et al., 2016). The Pleistocene activity progressed with the local emission of comendite lava (1.2 Ma; Savelli, 1987) and the emplacement of pyroclastic trachytic products in the southern part of Ponza (1.2-0.9 Ma; Savelli, 1987; Bellucci et al., 1999; Fig. 17a), where the resumption of volcanism coincides with the transition from a submarine to a subaerial environment. The Pleistocene cycle ended at ca. 1.0 Ma with the emplacement of trachytes of potassic series (K-alkaline, KA; Fig. 17b) in the southeastern part of Ponza (i.e., the Monte la Guardia lava dome, Punta della Guardia lava dyke, Fig. 16c). This episode represents the debut of the potassic alkaline magmatism that successively developed southeastward to the eastern Pontine islands up to the Roman-Campanian Magmatic Provinces. The eastern Pontine Islands, Ventotene and S. Stefano, display a similar chronological sequence of the outcropping units, which were erupted during the 0.9 and 0.1 Ma time span (Fig. 17a). It was initially characterized by effusive activity comprising a number of trachybasaltic lava flows (Fig. 17b) followed by a huge explosive phase, which ended with the caldera formation. The composition of the volcaniclastic products ranges from latitic to trachytic and phonolitic (Fig. 17b; Bellucci et al., 1999).

Further insights on the magmatic spectrum existing in the Pontine Archipelago (Fig. 17b) were provided by submarine rocks of relatively undifferentiated compositions (basalt to andesites), so far missing in the literature, collected in outcrops along the continental slope (Conte et al., 2016), and in submarine volcanic edifices (i.e., the Ventotene Volcanic Ridge; Conte et al., 2020; Fig. 16a, 17a, 17b). Geochemical data from these new findings, and thermodynamic modeling allowed to rule out an anatexic origin for both subalkaline and peralkaline rhyolites recognized in the Archipelago (Paone, 2013) and to infer that the whole HKCA-and TR-series felsic rocks, and the KA as well, derived by the respectively poorly evolved rocks by fractional crystallization processes (Fig. 17b). Moreover, the new geochemical data confirm the orogenic signature of the three suites and indicate highly heterogeneous mantle sources, due to crustal components variously recycled in the mantle via subduction. In addition, the age of these new findings (Conte et al., 2020) led to infer that the Pliocene and Pleistocene volcanic cycles developed in the Pontine Archipelago are likely associated to the Pliocene E-W directed rifting stage of the Tyrrhenian back-arc basin, which produced a more intense activity during the time span ~5.0-2.0 Ma (the emplacement time of HKCA and TR series). After that, this fault activity decreased and the extensional direction changed to NW-SE causing the beginning of the KA activity (<~2.0 Ma).




Fig. 18 - (a) Location of the Pontine Archipelago in the central Tyrrhenian Sea. The inset indicates the morphological high of Ventotene Volcanic Ridge (modified after Conti et al., 2016 and 2020); (b) Selected portion of the 1:50,000 geological map of Foglio 413-Borgo Grappa (reported at reduced scale). In the simplified map legend are reported only the main volcanic units; (c) Pliocene: hyaloclastite cryptodome intruded by dykes at Ponza; (d) Pleistocene: trachytic lava dyke at Punta della Guardia, Ponza, showing cooling structures (vertical and curved columnar jointing).




Fig. 17 - (a) Chronology and serial affinity of Pliocene-Quaternary volcanism of the Pontine Archipelago and Ventotene Volcanic Ridge. Yellow, HKCA; pale green, KA; dark green, TR. See the text for the abbreviations; (b) Total Alkali-Silica (TAS; Le Maitre et al., 2002) classification diagram reporting the whole range of compositions of volcanic rocks from Pontine Archipelago. The arrows indicate the liquid lines of descent simulating fractional crystallization trends leading to the felsic rocks constituting the islands from the relatively undifferentiated parental melts recovered in the islands and surroundings. Specifically, a trachybasalt similar to that of Ventotene led to the KA trachyte and trachyte-phonotephrite of Ponza and Ventotene, respectively; andesite rocks recovered by dredge in the deep water (yellow filled circles) led to the HKCA Pliocene rhyolites, and transitional basalts from both deep water (green small filled circles) and Ventotene Volcanic Ridge led the TR Pleistocene peralkaline rhyolites from Palmarola (Conte et al. 2018, 2020).

* The age of Zannone Island is deduced by the morphological continuity with Ponza and the compositional affinities of the products of the two islands. Indeed, since now no absolute dating exists for the Zannone due to the high alteration degree of the volcanic rocks (Dolfi & Conte, 2018).

4.2. The Neapolitan volcanoes: Somma-Vesuvius, Campi Flegrei and Ischia Island

(R.S., R.C., S.d.V., M.A.D.V., R.I., S.M.)

Somma-Vesuvius, Campi Flegrei and Ischia Island are among the most famous active volcanoes in the world and are part of the Neapolitan landscape. The activity of the volcanoes punctuated the human history since Bronze Age, as testified by archaeological findings recognised in the plain and relieves surrounding the volcanic area. The world-famous eruption of 79 CE consigned the Somma-Vesuvius to the history, because of the burying of some important Roman towns like Pompeii and Herculaneum. The Campi Flegrei volcanic field largely contributed to the birth of myths, and was one of the seven gates of Roman hell. Ischia Island was always a favorite place for thermal baths since Roman times.

This brief chapter summarizes the main characteristics of these three world famous volcanoes, resuming their eruptive history, eruptive behavior, composition of their products and finally, the main hazards they pose to the surroundings and distal regions.

4.2.1. Somma-Vesuvius (39 ka-1944 CE)

The Somma-Vesuvius volcanic complex consists of an older volcano dissected by a summit caldera, Monte Somma, and a recent cone, Vesuvius, built within the caldera after the 79 CE "Pompeii" eruption (Cioni et al., 1999). The original Roman name Vesuvius (or Vesibus) was first applied to the old volcano. Starting from the fifth century, chroniclers make mention of Mt. Somma, as the highest ("summa") peak of the mountain. The new cone grew discontinuously during periods of semi-persistent, low-intensity activity (from Strombolian to violent Strombolian, accompanied by important effusive activity). These periods possibly occurred in the first to third centuries, in the fifth to eighth centuries (after the 472 CE "Pollena" eruption), in the tenth to twelfth centuries, and in 1631-1944 time span (Cioni et al., 2006; Santacroce et

al., 2008).

Until relatively recent times, the formation of Somma-Vesuvius (SV) caldera was ascribed to the Pompeii eruption. Roman paintings from Pompeii and Herculaneum prompted Stothers & Rampino (1983) to conclude that, prior to 79 CE, the top of the volcano was asymmetrically shaped, indicating that a Somma-type caldera was already present. The volcanological interpretation of the Roman fresco from Pompeii "Mars and Venus", now at Naples National Archaeological Museum, made by Nazzaro (1997) leaves few doubts about the presence of a pre-existing caldera (Cioni et al., 1999). The SV caldera has a lobate, quasi-elliptical shape with a 5-km-long, east-west major axis (Fig. 18a). The northern rim of the caldera is a well-defined steep wall, with an average elevation of approximately 1000 m. The drainage pattern of the highest portions of the volcano, unaffected by conspicuous deposition of recent products, is clearly radial, suggesting a symmetrical original cone. In this assumption, the shape and size of the ancient Mt. Somma can be constrained, placing the apex of the old volcano around 1900 m elevation and approximately 500 m North of the present Vesuvius crater. Although the general morphology of the volcano (Fig. 18a) could be suggestive of the occurrence of lateral collapses of the seaward, western sector (Milia et al., 1998; Ventura et al., 1999; Rolandi et al., 2004), field evidence discards such a possibility at least for the last 20,000 years (Sulpizio et al., 2008).

About 22 cal ky BP, the mainly effusive activity of Mt. Somma changed into largely explosive (Fig. 18b). At least four high-magnitude Plinian eruptions occurred, staggered with minor events covering a large range of magnitude and intensity. High-intensity, explosive eruptions sporadically occurred, the two largest being the Subplinian events of 472 CE (also known as Pollena eruption; Rosi & Santacroce, 1983; Sulpizio et al., 2005) and 1631 CE (Rosi et al., 1983; Bertagnini et al., 2006).




Fig. 18 - (a) Geological map of Somma-Vesuvius volcano (Santacroce et al., 2003; Sbrana et al., 2020). For the legend the reader has to refer to the original publication; (b) Chronogram of Somma-Vesuvius activity. Arrows refer to explosive eruptions, length and colour reflect the estimated VEI. Blue boxes show recorded or inferred periods of persistent mild Strombolian and effusive activity, punctuated by VEI 2-3 explosive eruptions; (c) Variation of Somma-Vesuvius rocks and glasses from the major explosive eruptions within Total Alkalies vs. Silica (TAS) diagram (Santacroce et al., 2006). The variations of Nb, Sr and Ba of major explosive eruptions of SV discriminate the three groups of rocks, and are peculiar for most of single eruptions. Note the Pompeii trend in the Sr vs. Ba plot, completely cut of the "mildly silica-undersaturated" field (from Santacroce et al., 2006); (d) PDC inundation frequency for the main eruptions of Somma-Vesuvius during the last 22 ka. The map shows areas that relate to high, medium, and low frequency of PDC inundation during the last 22 ka of activity (from Gurioli et al., 2010); (e) Probability hazard map for fallout load exceeding 300 kg/m² for large explosive eruptions at Somma-Vesuvius (from Sandri et al., 2016).

The most recent period (1631-1944) was characterised by summit or lateral lava effusions and semi-persistent, mild explosive activity (small lava fountains, gases and vapour emission from the crater) interrupted by pauses lasting from months to a maximum of seven years (Santacroce, 1987; Arrighi et al., 2001; Fig. 18b). Lateral activity was minor and, starting from the 79 CE, the few eruptive fissures were confined to the western and southern sectors of the volcano (Cortini & Scandone, 1962; Santacroce, 1987; Principe et al., 2004; Acocella et al., 2006).

SV activity has been characterised in the last 20 ka by the eruption of Potassic and Ultra-Potassic magmas spanning a wide compositional spectrum and showing an increase with time of both K₂O content and silica undersaturation (Fig. 18c). The shift from Potassic magmas (represented by the trachytes and latites erupted in the period from the Pomoci di Base to the Greenish Pumice) to K-rich compositions occurred before 8 ka. The products younger than 8 ka have been characterised by increasing alkalinity, and the products of the last 2 ka of activity (following the 79 CE Pompeii Pumice) show the most alkali-rich compositions and the lowest SiO₂ content of the whole set of the erupted products.

Slightly silica - Undersaturated (to saturated) magmas (blue field; Fig. 18c)

Two large eruptions related to Somma-Vesuvius activity occurred in the period preceding the Mercato Eruption: the Subplinian, 19,000 cal yr BP "Greenish Pumice" and the Plinian, 22,000 cal yr BP "Pomoci di Base". The most relevant common geochemical signature of the rocks of all these eruptions, related to their K-trachytic highly evolved composition, concerns the higher silica and lower alkali contents (with respect to rocks with comparable degree of evolution of the other two groups), as well as their moderate Sr and Ba contents (Fig. 18c). As a whole, the geochemical features of these rocks are coherent with evolutionary trends of K-basaltic liquids initially driven by crystallization of mafic phases and plagioclase and later involving K-feldspar fractionation.

Mildly silica - Undersaturated magmas (green field; Fig. 18c)

This group consists of the pyroclastic products of three Plinian eruptions (79 CE "Pompeii", 3910 cal yr BP "Avellino" and 8,900 cal yr BP "Mercato") and at least six other explosive eruptions occurred between the Avellino and Pompeii events (AP1 to AP6; Andronico & Cioni, 2002). Differently from the Mercato deposits, characterized by a strong compositional homogeneity all along the whole eruptive sequence, the products of the other eruptions of this group present important compositional variations, from the most evolved products at the base to the least evolved toward the top. The Mercato and Avellino eruptions (first-erupted white pumice) are characterized by the emission of the most evolved products of Somma-Vesuvius (CaO<2.0%, Nb>100 ppm; Zr>700 ppm, Th up to 100 ppm).

Strongly silica - Undersaturated magmas (pink field; Fig. 18c)

This group comprises all the products erupted after the 79 CE Pompeii eruption. The most relevant common geochemical signatures of these rocks concern their lower silica and higher alkali contents with respect to

rocks with comparable evolution from the other two groups, as well as their very high Sr and Ba contents (Fig. 18c). These features are consistent with evolutionary trends (mostly fractionation within a periodically supplied magma chamber) of K-tephritic liquids dominated by crystallization of leucite and mafic minerals, and characterized by a minor role of plagioclase and the absence of K-feldspar fractionation.

The products of the largest Somma-Vesuvius eruptions are widely dispersed in central Mediterranean area, many of which being important stratigraphic markers for the regional archaeological and paleoclimatic investigations. Codola is dispersed in Tyrrhenian Sea and Adriatic Sea and on land in San Gregorio Magno Basin (Petrosino et al., 2019), Lake Monticchio (Wulf et al., 2004) and Apulia, where acts as a marker for the Palaeolithic successions (Giaccio et al., 2008). Mercato is an important marker for synchronizing the sapropel S1 in Adriatic Sea series with onland Holocene successions of the central-southern Italy and western area of the Balkan Peninsula (e.g., Zanchetta et al., 2011). The Avellino and Pompeii tephra, with their quite opposite dispersal axis, from east to north and south-east, respectively also act as relevant stratigraphic markers for dating and synchronizing marine and terrestrial successions of the southern-central and southern Italy (Zanchetta et al., 2011; Crocitti et al., 2019; Doronzo et al., 2022).

4.2.2 Campi Flegrei (>80 ka-1538 CE)

Campi Flegrei (CF) is a ca. 14 km wide volcanic caldera located in Southern Italy, in a highly urbanized area including the western part of the city of Naples.

The Campi Flegrei volcanic field (Fig. 19a), includes small volcanic apparatus and several monogenetic tuff rings, tuff cones and rarely cinder cones and lava domes. These volcanoes are exposed outside, on the borders, and within a large polygenetic caldera formed by the eruptions of the Campanian Ignimbrite (CI) and the Neapolitan Yellow Tuff (NYT) (e.g., Armenti et al., 1983; Di Girolamo et al., 1984; Rosi & Sbrana, 1987; Orsi et al., 1992, 1995, 1996, 1999; Cole & Scarpati, 1993; Rosi et al., 1996, 1999; Di Vito et al., 1999; De Vivo et al., 2001; Ort et al., 2003). The northern and western parts of the caldera are above sea level and characterized by the presence of many dispersed cones and craters, whereas the southern part is principally submarine and extends into the Gulf of Pozzuoli. The oldest rocks date back till ca. 80 ka and are represented by pyroclastic deposits cropping out outside and along the borders of the caldera (Pappalardo et al., 1999; Scarpati et al., 2012). Pre-caldera volcanism mainly generated monogenetic tuff cones and subordinate lava domes widespread also within the city of Naples (e.g., Rosi & Sbrana, 1987; Scarpati et al., 2012). Thick pyroclastic sequences generated from at least 80 ka also occur (Orsi et al., 1996; Pappalardo et al. 1999; (Fig. 19a). The caldera forming large volume CI eruption (Fisher et al., 1993; Rosi et al., 1996) erupted ca. 300 km³ of trachytic dense rock equivalent magma (DRE). The CI eruption was the largest magnitude eruption to occur in the Mediterranean region during the late Quaternary, and resulted in the formation of a 14-km wide caldera (Rosi & Sbrana, 1987). The eruption began with a Plinian phase, during which a SE-distributed pumice fallout was emplaced. This was followed by a suc-




Fig. 19 - (a) Geological sketch map of the Campi Flegrei (from Vitale and Isaia 2014); (b) Chronostratigraphy of Campi Flegrei activity (from Vitale and Isaia, 2014); (c) Frequencies of PDC arrival computed using EC simulations (SIM) and PDC deposits (OBS) at Campi Flegrei. Yellow star indicates downtown Naples (from Tierz et al., 2016); (d) Conditional probability maps for a loading threshold of 300 kg/m², for size Small, Medium, Large and all sizes (from Selva et al., 2018).

sion of pyroclastic density currents (PDC) that deposited ash and pumice flows and densely-welded ignimbrites that covered the Campanian Plain and surrounding hills (Barberi et al., 1978). Proximal deposits (e.g., Breccia Museo, Piperno) cropping out at the top of the eruption deposits along the caldera margins are interpreted as proximal facies related to the final caldera-forming phase (Rosi & Sbrana, 1987; Rosi et al., 1996). On these deposits, several age data were produced (ca. 37 ka, Deino et al., 2004; ca. 39 ka, De Vivo et al., 2001; ca. 38 ka, Fedele et al., 2008). A recent and high precision ^{14}C and $^{39}\text{Ar}/^{40}\text{Ar}$ age determination of the CI yielded an age of 39.85 ± 0.14 ka (Giaccio et al., 2017b).

After the CI caldera forming eruption volcanic activity occurred almost exclusively through the formation of pyroclastic deposits from several volcanic centers located within or along the border of the caldera. This period of activity culminated with the second large eruption at CF occurred at 15 ka (Deino et al., 2004) named Neapolitan Yellow Tuff (NYT; Cole & Scarpati, 1993; Wohletz et al., 1985; Fig. 19b), whose volume has been estimated larger than 20 km³ DRE of latitic to trachytic magma (Orsi et al., 1992, 1995; Scarpati et al., 1993). The eruption was mainly phreatomagmatic producing pyroclastic density currents and subordinate thin fallout layers. The PDC deposits close to CF caldera were largely zeolitized and form the substrate on which lies the city of Naples. Ashes were mainly dispersed toward North-Northeast from the vent area.

The eruptions of the post-NYT period were confined within the structural boundaries of the caldera and comprised at least 70 known events, dominated by low- to medium-magnitude phreatomagmatic-magmatic eruptions with volumes of <0.1 km³ (Di Renzo et al., 2011; Smith et al., 2011). The Monte Nuovo tuff cone formed as consequence of the most recent eruption in 1538 CE (e.g., D'Orlando et al., 2005; Di Vito et al., 2016). After the NYT eruption, the caldera suffered significant ground deformation phenomena, especially in its central sector, where the uplift is still ongoing.

The primary magmas produced have a mid-ocean ridge basalt (MORB)-like asthenospheric mantle wedge composition, and these are modified by aqueous fluids, oceanic sediments, and continental crust (Tonarini et al., 2004; D'Antonio et al., 2007). The mafic melts at Campi Flegrei are K-basalts (e.g., Webster et al., 2003). These mafic compositions are preserved as melt inclusions in antecrustal Mg-rich olivines and clinopyroxenes in some eruption deposits (e.g., Cannatelli et al., 2007). The composition of the erupted melts range from shoshonitic to phonolitic and trachytic, with the most differentiated compositions dominating (e.g., Mangiacapra et al., 2008; Smith et al., 2011; Tomlinson et al., 2012; Fig. 19c). The phenocrysts in these Campi Flegrei magmas are predominantly plagioclase + K-feldspar + clinopyroxene ± biotite. Magnetite and apatite occur as accessory phases and eruption products occasionally contain olivine or rare feldspathoids.

The SiO₂ and Na₂O contents of the magmas increase with differentiation, whereas CaO, FeO, MgO, and P₂O₅ contents decrease (e.g., Civetta et al., 1991b). There is a noticeable inflection in K₂O melt compositions, denoting K-feldspar-in, at ca. 80 wt% SiO₂ (Fowler et al., 2007; Smith et al., 2011; Tomlinson et al., 2012). The Sr,

Ba, and Eu contents behave compatibly, and reflect the significant amount of felspar fractionation. Other REE (excluding Eu), Y, Nb, Zr, Rb, Th, and Ta are all incompatible (e.g., Civetta et al., 1997; Bohrson et al., 2008; Arlenzo et al., 2010; Tomlinson et al., 2012). These major and trace element compositions follow an evolutionary trend that could be generated through fractional crystallization of a single parental melt (e.g., Civetta et al., 1991a; D'Antonio et al., 1999; Fourmentraux et al., 2012) but isotopic variations indicate that the melts that erupt are derived from different batches of magma (Pappalardo et al., 1999, 2002; D'Antonio et al., 2007; Di Renzo et al., 2011; Fig. 19c).

Samples from Campi Flegrei eruption deposits suggest that only evolved magmas were erupted in the early history (Pappalardo et al., 1999) and it was only after the last caldera-forming NYT eruption (ca. 15 ka) that more compositionally diverse melts were erupted (e.g., D'Antonio et al., 1999; Smith et al., 2011). The isotope (Nd, Pb and Sr) and occasionally the major and trace element glass compositions of the magmas indicate that the eruptions tap distinct batches of melt, and some have interacted at depth (e.g., Di Renzo et al., 2011). This has been well documented for the large caldera-forming events (e.g., Forni et al., 2018) and for eruptions in the last 15 ka (e.g., Tonarini et al., 2009; Fourmentraux et al., 2012). The general trend between 60 and 10 ka is that Nd and Pb isotopic compositions of the erupted magmas became progressively less radiogenic ($^{143}\text{Nd}/^{144}\text{Nd}$ - 0.51252 to 0.51236 and $^{206}\text{Pb}/^{204}\text{Pb}$ - 19.2 to 18.9) while the Sr-isotope composition became more enriched (0.70700 to 0.70884) over time (Pabst et al., 2008; Di Renzo et al., 2011). These changes in the isotopic compositions are consistent with an increase in crustal contamination. However, the Campi Flegrei liquid line of descent and extent of Sr and Pb isotopic heterogeneity is compatible only with very minor assimilation (D'Antonio et al., 2007; Fowler et al., 2007).

The continuous and intense activity of the Campi Flegrei, makes this volcanic system the main source of the late Pleistocene tephra in central Mediterranean area. Y-5, the tephra of the CI, the largest Campi Flegrei eruption, is to great extent the most widespread, and one of the most relevant stratigraphic markers for synchronizing stratigraphic, paleoclimatic and cultural events occurred in western Eurasia around 40 ka (e.g., Giaccio et al., 2017b). Though still poorly known in near vent sections, a number of other widely dispersed tephra preceding the CI eruption, including, among other the X-6, X-5 and C-22 marine layers, have been attributed to previously unknown 92–109 ka Campi Flegrei activity (Monaco et al., 2022b). Following the CI eruption, the two other tephrostratigraphic markers, which significantly exceed the regional dispersal, are those of the relatively poorly known Masseria del Monte eruption of ca. 29 ka (Y-3, Albert et al., 2019) and of the Neapolitan Yellow Tuff, mainly dispersed toward North and found until central Europe (e.g., Lane et al., 2011; 2015).

4.2.3. Ischia Island (>150 ka-1302 CE)

The island of Ischia is part of the Phlegraean volcanic district, related to extensional tectonic phases that accompanied the anticlockwise rotation of the Italian peninsula (Ippolito et al., 1973; D'Argenio et al., 1973;

Finetti & Morelli, 1974; Bartole, 1984; Piochi et al., 2005). Ischia is the emerged part of an active volcanic field, which rises more than 1,000 m above the sea floor (Fig. 20a; Orsi et al., 1999; Bruno et al., 2002), along the margin of an E-W trending scarp that borders to the south the Phlegraean volcanic district. The island covers an area of 46.4 km² and is morphologically dominated by the Monte Epomeo (787 m a.s.l.) in its central portion, and by the NE-SW Monte Vezzi - Monte Cotto alignment in the SE corner.

Volcanism at Ischia dates back to more than 150 ka and continued, with centuries to millennia of quiescence, until the most recent eruption occurred in 1302 CE (Vezzoli, 1988; Sbrana & Toccaceli, 2011; Sbrana et al., 2018; Fig. 20b). The oldest exposed rocks, aged between 150 and 75 ka, are lava flows and/or lava domes, and a sequence of pyroclastic deposits with intercalated paleosols, mainly exposed in the southern part of the island (Scarrupata di Barano Formation in Vezzoli, 1988; Ancient Ischia Synthem in Sbrana & Toccaceli, 2011; Phase 1 in Sbrana et al., 2018; i.e. volcanics older than 75 ka of Figs. 20b and c). Almost in the same time span, a prevailing effusive activity determined the emplacement of a series of small trachytic and phonolitic lava domes, which are presently exposed at the periphery of the island (Fig. 20a).

A very intense period of explosive activity followed between 74 and 55 ka. During this period, many volcanic vents were active mainly in the southern sector of the island, likely producing the highest magnitude eruptions recorded at Ischia. The deposits of at least 10 explosive eruptions, fed by phonolitic to trachytic magmas, generated pyroclastic density currents, fallout deposits, block and ash flows from collapsing lava domes, explosion breccias and hydromagmatic dilute and turbulent pyroclastic density currents (Brown et al., 2008; Rifugio di San Nicola Synthem in Sbrana & Toccaceli, 2011; Phase 2 in Sbrana et al., 2018). The activity of this period culminated with the caldera forming eruptions of the Monte Epomeo Green Tuff (MEGT) between 60 and 50 ka (Brown et al., 2008; Sbrana & Toccaceli, 2011; Sbrana et al., 2018; Fig. 20b).

This tuff consists mostly of trachytic ignimbrites that partially filled the caldera depression in a submarine environment, and were also emplaced on land outside its margins. The caldera depression was later the site of marine sedimentation, which formed a succession of clays, tuffites, sandstones and siltstones by the reworking of MEGT and sedimentary supply from the mainland (Sbrana & Toccaceli, 2011). After the MEGT eruptions, volcanism continued with a series of hydromagmatic and magmatic explosive eruptions up to 33 ka. Most of the vents were located along the present SW and NW offshore of the island. Volcanism in this period was fed by latitic and shoshonitic magmas (Brown et al., 2014; Phase 3 in Sbrana et al., 2018), whose injection into the system triggered the asymmetric resurgence of the caldera floor, generating the Monte Epomeo block (Orsi et al., 1991). After an about 5 ka long period of quiescence, volcanism resumed at ca. 28 ka. The beginning of this period of activity was marked by the eruption of shoshonitic magmas, isotopically distinct from those characterizing the previous eruptions, suggesting the arrival of a new batch of magma into the feeding system of Ischia,

followed by its differentiation and mixing with the resident magma (Poli et al., 1987; Civetta et al., 1991b; Casalini et al., 2017). Volcanism continued sporadically until 18 ka (or 13 ka, Phase 4 in Sbrana et al., 2018), with the emission of shoshonitic to alkali-trachytic magmas that fed effusive and both hydromagmatic and magmatic explosive eruptions, with the emplacement of lava flows and Strombolian fallout deposits, and the construction of small tuff-cones and scoria-cones, exposed along the SE and SW coasts of the island (Fig. 20c; Campotese Sub-synthem in Sbrana & Toccaceli, 2011).

The radiometric dating of the outcropping volcanic products suggests that between 18 and 10 ka there was a period of almost complete quiescence (Vezzoli, 1988), whereas other interpretations (Sbrana & Toccaceli, 2011; Sbrana et al., 2018) seem to indicate the occurrence of some eruptions in this timespan. In any case, all the authors agree in considering the interval between 18 and 10 ka as a period of quiescence or very reduced volcanic activity (Fig. 20b). A period of more sustained activity at Ischia started at ca. 10 ka, and was fed by mainly trachytic and subordinately latitic magmas. These are characterized by a Sr isotopic ratio lower than those of the previous periods, leading to hypothesize the arrival of a new, geochemically distinct magma into the system. Volcanism was mainly concentrated around 5.5 ka and after 2.9 ka, with almost all the vents located in the eastern part of the island (Fig. 20c). Only a few vents are located outside this area, along regional fault systems or along the margins of the resurgent block: these vents generated a multi-vent lava field in the NW corner of the island, a pyroclastic sequence, exposed to the SW, and a lava dome in the N sector of the island. Since 2.9 ka, about 34 effusive and explosive eruptions took place: the effusive eruptions emplaced lava domes and high-aspect ratio lava flows, while the explosive eruptions, both magmatic and phreatomagmatic, generated tuff cones, tuff rings and variably dispersed pyroclastic-fall-and-current deposits, which had a very variable impact on both the island's environment and human settlements (de Vita et al., 2010 and references therein; Sbrana & Toccaceli, 2011; de Vita et al., 2013; Phases 5 and 6 in Sbrana et al., 2018). As volcanism was not continuous since 10 ka, but characterized by the alternation of centuries of quiescence and periods of very intense activity, it has been hypothesized that repeated episodes of magma intrusions occurred intermittently (Tibaldi & Vezzoli, 2004; de Vita et al., 2006; Vezzoli et al., 2009; de Vita et al., 2010), likely accompanied by resurgence. Even if, since the last eruption, no evidence of renewal of uplift due to fault reactivation has been recorded in concurrence with more recent minor mass movements, the magmatic system of Ischia has to be considered still active, as testified by intense volcanism in historical times and widespread fumaroles and thermal springs.

The Ischia volcanoes belongs to the low-K series of the Roman comagmatic province, ranging in composition from shoshonite to latite, trachyte and phonolite; the most abundant exposed rocks are alkali-trachytes (Angiuli et al., 1985; Poli et al., 1987, 1989; Crisci et al., 1989; Civetta et al., 1991; Fig. 20d).

The activity of Ischia volcano is well documented in distal settings (e.g., Wulf et al., 2004; 2008). The most widespread Ischia tephra is the one related to the Monte




Fig. 20 - (a) Geological and structural sketch map of Ischia volcanic field. Deposits of debris avalanches and volcano veins are highlighted in marine areas (from Sorana et al., 2018). For the legend the reader has to refer to the original publication; (b) simplified chronostratigraphic sequence (not to scale) of Ischia volcanic field (from Sorana et al., 2018); (c) Total alkali vs. silica classification diagram showing the variability of composition for Ischia products (from D'Antonio et al., 2021; Santacroce et al., 2008). K-B = potassio basalt; K-TB = potassio trachybasalt; SH = shoshonite; LT = latite; TR = trachyte (these rock names are relative to a potassio alkaline series); TH-Bas = tephrite-basanite; PH-TH = phonotephrite; TH-PH = tephriphonolite; PH = phonolite; (d) distribution of PDC in the last 10 ka (from Selva et al., 2019); (e) distribution of pyroclastic fall-out deposits for all the eruptions in the last 10 ka (from Selva et al., 2019).




Fig. 21 - Geological map of Procida Island with the chronological classification of the main volcanic events and the location of the main coastal sites (from Aucelli et al., 2022, modified after Fedele et al., 2012). The position of the submerged marine terraces was obtained from Putignano et al. (2014).

Epomeo Green Tuff eruption of ca. 56 ka (Tomlinson et al., 2014), matching the layer TM-19 of Monticchio (Wulf et al., 2004), S16 of San Gregorio Magno (Petrosino et al., 2019) and TF-7 of Fucino (Giaccio et al., 2017a), and a still unknown, slightly older, ca. 80 ka, eruption that matches the widespread marine layer Y-7 (Keller et al., 1978) and the Monticchio tephra TM-20 (D'Antonio et al., 2021). Several tephra layers that are compositionally attributable to Ischia, preceding both MEGT and the unknown Y-7-related eruption, found, e.g., in the Tyrrhenian Sea (Paterne et al., 2008) and Fucino (Monaco et al., 2022a), extend the activity of this volcano at least to 200 ka.

4.2.4. Procida Island

The volcanic island of Procida formed due to several monogenetic explosive eruptions that led to the formation of five volcanic centres characterizing the emerged-submerged morphology of the island: Vivara, Pozzo Vecchio, Terra Murata, Fiumicello, and Solchiaro (Fig. 21; Aucelli et al., 2022). These tuff rings, except the tuff cone of Terra Murata, are SW-NE oriented following a fault line crossing the centre of Procida and the nearby island of Ischia (e.g., Pescatore & Rolandi, 1961).

De Astis et al. (2004) tentatively constrained the beginning of the volcanic activity of Procida Island around 70 ka and several authors (Alessio et al., 1976; Llerer et al., 1991; Fedele et al., 2012) identified the last

volcanic episode of the area with the Solchiaro eruption, occurred 22 ka (Fedele et al., 2012). The different volcanic episodes, which led to several volcano-tectonic collapses, modified the original structure of the island together with the activity of the main fault systems (Putignano & Schiattarella, 2010). The deposits deriving from these activities, mainly made up of stratified to massive ash levels, are covered by a tephra succession, cropping out along the SW flank of Vivara and Pozzo Vecchio and partially at Terra Murata, and interbedded with the volcanic formation of Pignatello (55–74 ka, Fedele et al., 2012).

Procida rocks range in composition from basalt to shoshonite and trachyte. The presence of a compositional gap in the range $\text{SiO}_2=54\text{--}59\text{ wt }%$ is evidence of magma bimodality, suggesting that the feeding magmatic system was formed by at least two different reservoirs located at different depths (De Astis et al., 2004).

4.2.5. Hazard assessment

The Neapolitan area represents one of the highest volcanic risk areas in the world due to the presence of three active volcanoes. In the last decades, many studies assessed tephra fall hazard from them combining field data and numerical simulations. Initially, conditional ash load probability maps for one or few reference volcanic scenarios were provided (e.g., Barberi et al., 1995; Cioni et al., 2003; Macedonio et al., 2008; Costa et al., 2009; Folch & Sulpizio, 2010; Sulpizio et al., 2012). The choice of selecting a single scenario was very useful to support civil protection emergency plans (e.g., Emergency Plan of Vesuvius; DPC 2015) and it is computationally less expensive and more feasible for near-real-time applications (e.g., Selva et al., 2014). However, this strategy was not appropriate to achieve an unbiased Probabilistic Volcanic Hazard Assessment (PVHA; Marzocchi et al., 2010; Selva et al., 2010; Sandri et al., 2016) since the uncertainty only accounted for wind conditions (e.g., Macedonio et al., 2008; 2016).

To overcome this limitation, the analysis of the intra-scenario variability is certainly more complete, allowing a reduction of the epistemic uncertainty (e.g., Selva et al., 2014; Sandri et al., 2016; Titos et al., 2022) in both long- and short-term hazard assessments (e.g., Selva et al., 2010; Sandri et al., 2012; 2014; Selva et al., 2014; Thompson et al., 2015; Constantinescu et al., 2016).

In more recent PVHAs at the Neapolitan volcanoes, new sampling procedures have been set to fully explore the intra-size-class aleatory variability comparing the results with the classical approach based on reference volcanic scenarios, in the case of proximal/medial areas and large tephra loads from Somma-Vesuvius and Campi Flegrei (Sandri et al., 2016) or by using ensemble modeling of submarine eruptive vents and tephra total grain-size distributions considering variable mass fraction and aggregates for ash (Selva et al., 2018). Moreover, Montesinos et al. (2022) have developed a new Bayesian Event Tree based on high performance computing applied to Campi Flegrei, aiming to robust and unbiased short- and long-term PVHA on a large-scale (thousands km) and high-resolution (about 2 km) domain to be used by Civil Protection agencies, aviation companies and other stakeholders.

A novel contribution in the quantification of volcanic

hazard regards the estimation of the present state of the Neapolitan volcanoes based on a simple physics-based statistical model that satisfactorily fits the eruptive history of each volcano (Selva et al., 2022). The model is compatible with existing data (including isolated events and long repose periods) and accounts for two activity regimes (high-low) able to describe the temporal modulations in eruptive activity and to provide a homogeneous quantification of the probability of eruption, considering the state of the volcano and the possible transitions.

More recently, Massaro et al. (2023) proposed a new methodology for a probabilistic long-term tephra fallout hazard assessment in Southern Italy from the active Neapolitan volcanoes (Somma-Vesuvius, Campi Flegrei and Ischia). By means of thousands of numerical simulations they quantify the mean annual frequency with which the tephra load at the ground exceeds critical thresholds in 50 years. The output hazard maps account for changes in eruptive regimes of each volcano and are also comparable with those of other natural disasters in which more sources are integrated.

For pyroclastic density currents, Gurioli et al. (2010) provided the high, medium and low inundation frequency from the main eruptions of Somma-Vesuvius during the last 22 ka of activity (Fig. 18d). More recently, quantitative PVHAs were carried out by Tierz et al. (2016) for Somma-Vesuvius and Campi Flegrei testing the energy cone model (Fig. 19d) and by Selva et al. (2019) for Ischia Island (Fig. 20e-f).

4.3. The Mt. Vulture and Monticchio Lakes nested volcanoes (>730–130 ka) (S.C., P.P., J.P.S.-B.)

The Monte Vulture and Monticchio nested volcanoes with few small eccentric explosive monogenetic centres along the Ofanto valley and at Ripacandida make up the Lucanian Magmatic Province (Fig. 1). The volcanism of this magmatic province is located East of the Apennine system, at the outermost edge of the fold and thrust belt. The volcano lies on a structural high made up of Meso-Cenozoic units of the substratum overlain by Pliocene continental sediments (Schiattarella & Beneduce, 2006) and is situated on a deep transfer fault (Trinitapoli-Paestum Line in Ciaranfi et al., 1983; Vulture Line in Schiattarella et al., 2005).

The volcanic activity at Vulture-Monticchio nested volcanoes (Fig. 22a) developed entirely within the Pleistocene starting at about 0.74 Ma (Villa & Buettner, 2009). A reconstruction of the eruptive history of the volcano, after the pioneering work of Heiko Merlin (1967), was carried out by Principe & La Volpe (1989a), who recognized several markers useful to build a stratigraphic framework. The eruptive history has been systematically reinvestigated in the frame of the CARG project by Giannandrea et al. (2006), who adopted USBU systematics to map the primary volcanic products and the sedimentary deposits of the several fluvial/lacustrine successions that characterize the perivolcanic area (Fig. 22b). They highlight the strong tectonic control on the whole of the Vulture activity and identify erosional phases, epiclastic sediments and/or paleosols which testify to the occurrence of variably long quiescence phases. The very first dated products pertain to the Faro d'Olivo sub-synthem (Giannandrea et al., 2006) and are represented by two main ignimbrite units (Ignimbrite A and B of Crisci et al.,




Fig. 22 - (a) Detailed geological sketch map of the Vulture volcanic products. Legend: 1) Monticchio Lakes; 2) Landslides, alluvial and lacustrine deposits; 3) Laghi di Monticchio Synthem (Upper Pleistocene); 4) Valle dei Grigi - Fosso del Corvo Synthem (Middle Pleistocene); 5) Melfi Synthem (Middle Pleistocene); 6) Barile Synthem (Middle Pleistocene); 7) Foggianello Synthem (Middle Pleistocene); 8) Fiumara di Atella Supersynthem (Upper Pliocene-Lower Pleistocene?); 9) Cretaceous to Miocene Apennines units; 10) thrusts; 11) reverse faults; 12) normal faults; 13) strike-slip faults; 14) high-angle faults; 15) crater rim; 16) secondary volcano eruptive centres. (b) Unconformity bounded stratigraphic units forming the whole volcanic products of the Vulture volcano. Modified from Schiattarella et al. (2005) and Giano et al. (2002); (c) TAS classificative diagram (Le Maître et al., 2002) for the Vulture volcanic rocks (modified from Conticelli et al., 2010); (d) View of the Lago Grande di Monticchio (available at fondoambiente.itluoghi/monticchio?dc).

1983), for the lowermost of which an $^{40}\text{Ar}/^{39}\text{Ar}$ age of 739 ± 12 ka has been determined (Villa & Buettner, 2009). These ignimbrites, to which a caldera collapse has been ascribed (Principe & Giannandrea, 2006), unconformably overlay the deposits of an older explosive activity (Campanile subsynthem, Giannandrea et al., 2006). A subaerial erosional phase, with the associated epiclastic sediments, marks the passage to the Barile synthem, to which the Toppe San Paolo subsynthem, a lava dome, on the NW rim of the Vulture, belongs, together with the Rionero subsynthem, a complex sequence of pyroclastic fall and pyroclastic flow deposits. Among these deposits, the fallout of Masseria Boccaglie (De Fino et al., 1982) has the prominent role of a stratigraphic marker (marker M3 of La Volpe & Principe, 1989) younger than 720 ± 25 ka (Buettner et al., 2006). Starting from the Rionero, but mainly in the Vulture-San Michele subsynthems (Principe & Giannandrea, 2006) the main stratovolcano was built

up between ca. 670 and 600 ka (Broccolini et al., 1994; Villa & Buettner, 2009) by alternating mildly explosive and effusive activity, respectively emplacing block and ash flow with minor pyroclastic fall deposits, and thin lava flows. A phreatomagmatic eruptive phase (Ventaruo subsynthem) marks the beginning of an intense phase of destruction of the stratovolcano, occurred mainly during the Melfi synthem deposition, in which parasitic vents along the north and north-eastern side of the volcano produced lava flows with the emission of the Melfi haüynophire (557 ± 7 ka, Bonadonna et al., 1998; 573 ± 4 ka, Villa & Buettner, 2009) and Piano di Croce lavas. The volcanic and tectonic activity of the Vulture area has been sealed by a thick paleosol (Schiattarella et al., 2005), which La Volpe & Principe (1989) had identified as marker M18. The last deposits, grouped in the Monticchio supersynthem by Giannandrea et al. (2006), were erupted starting from ca. 500 ka (Buettner et al., 2006

and references therein) from a series of monogenetic centers, and mostly show evidence of phreatomagmatic activity. This supersynthem also includes the ash tuff deposits of the two maar-like centres of Lago Grande and Lago Piccolo di Monticchio (Fig. 22d). The recommended age of the Lago Piccolo subsynthem, which gives the youngest $^{40}\text{Ar}/^{39}\text{Ar}$ age determined so far for the Vulture products, is 141±11 ka (Villa & Buettner, 2009).

The bulk of the erupted products mostly belong to a basanite-tephrite-phonolite series (Fig. 22c), with mafic products (basanites and tephrites) dominating in both lavas and pyroclastic rocks (e.g., De Fino et al., 1986; Beccaluva et al., 2002). More silica undersaturated rocks, though minor in volume, are typical of this volcanic complex (Hieke Merlin, 1964, 1967). Melilites, hauyne-bearing leucitites are also found at the end of the Monte Vulture volcanic activity in the form of dykes, plugs and lava flows in the north-western flank of the volcano. On the other hand, trachyphonolites are found as dykes in the neighboring of the volcanic complex, and may be representatives of an earlier, and less silica undersaturated stage of activity (Melluso et al., 1996; Beccaluva et al., 2002).

A small carbonatitic lava flow has been also found at Toppo del Lupo (e.g., Stoppa & Principe, 1997; D'Orazio et al., 2008; Stoppa et al., 2008). Within the hydro-magmatic units of the Monticchio supersynthem, round lapilli tuff units with ejecta of intrusive carbonatites and mantle nodules are visible (e.g., Jones et al., 2000; Rosatelli et al., 2000; Downes et al., 2002). K₂O contents of the volcanic products decrease with time passing from Monte Vulture to Monticchio volcanoes. Sodic compositions are also found among lapilli tufts of the Monticchio lake activity (Avanzinelli et al., 2008, 2009).

Two distal tephra attributed to Vulture activity and tentatively correlated with ignimbrites A and B of Crisci et al. (1983) despite the incomplete age overlap, were found embedded in the Montalbano Jonico marine sequence (Petrosino et al., 2015).

4.4. Aeolian Islands (1.3 Ma-Present)

(F.L., G.D.A., R.D.R., P.D., F.F., L.F., E.N., M.P., R.S., C.R., C.A.T., M.V.)

4.4.1. General background

The Aeolian Islands arc is the most active volcanic structure in the Mediterranean area, and is composed of seven volcanic islands (Alicudi, Filicudi, Salina, Lipari, Vulcano, Panarea, Stromboli) and several seamounts rising ca. 2000-3000 m above the seafloor in the Southern Tyrrhenian Sea (Romagnoli, 2013; Romagnoli et al., 2013) (Fig. 23a). They are arranged in an articulated, tectonically-controlled arc-shaped structure, crossed in its central sector by the NNW-SSE lithospheric Tindari-Letojanni fault system, which has developed in a complex subduction-type scenario during the Quaternary (Ventura, 2013). The oldest products from the submarine sectors are dated to 1.3 Ma (Beccaluva et al., 1995), whereas the age of emergent islands ranges between 270-250 ka (Locardi, 2011) and the Present (Fig. 23b). Different stages of volcanic activity through time are established on the basis of the available radiometric ages and stratigraphic relationships with marine terraces and tephra layers of Campanian and/or Aeolian origin (Lucchi

et al., 2013a and references therein). The Aeolian Islands volcanoes are typically characterized by a large variety of volcanic landforms and rock types which mostly depends upon the wide compositional range observed in the erupted products from basalts to rhyolites with calc-alkaline (CA) to potassic signature (De Astis et al., 2013; Forni et al., 2013; Francalanci et al., 2013; Lucchi et al., 2013a, b, c, d, e; Nicotra et al., 2020) (Fig. 23c), and the corresponding eruption types (from Strombolian/Hawaiian to Subplinian), under the direct control of regional fault systems and recurrent calderas and lateral collapses. The western Aeolian volcanoes (Salina, Filicudi and Alicudi) are considered extinct, and currently active to quiescent volcanoes (Vulcano, Stromboli, Panarea, Lipari) are basically restricted to the eastern and the outer portions of the sector of the Aeolian Islands arc. Major explosive eruptions involving more evolved magmas occurred during the last 75 ka on Stromboli (Petrarza Tufts), Salina (Grey Ponni Tufts, Lower and Upper Pollara Tufts) and Lipari (Monte Guardia, Vallone del Gabellotto, Monte Pilato) producing widespread tephra layers recorded in the Tyrrenian, Ionian and Adriatic deep sea cores and other lake or terrestrial archives in southern Italy (Albert et al., 2017; Meschiali et al., 2020 for references).

Hereafter we describe the most important geological features and chronology of the Alicudi, Filicudi, Salina, Lipari and Panarea volcanic complexes (from West to East), whereas a particular attention will be given to Vulcano and Stromboli in dedicated paragraphs.

The island of Alicudi is the emerged portion of a composite volcano constructed between 106 and 28 ka by lava flows, domes and Strombolian scoriae with a compositional range between CA basalts and high-K CA andesites (Lucchi et al., 2013d). Six Eruptive Epochs were characterized by a mainly central-type volcanism interrupted by periods of dormancy and three summit caldera-type collapses (Lucchi et al., 2013d). The Alicudi rocks are characterized by the most primitive geochemical and isotopic signature of the entire Aeolian archipelago, with a progressive shift toward more acid products due to the evolution of the magma feeding system and polybaric crystal fractionation, and a lower control by extensional regional tectonics (Peccerillo et al., 2013).

The island of Filicudi is the emerged portion of a volcanic complex, result of the superimposition of lava flows, lava domes and coulees and subordinate pyroclastic deposits ranging in composition from CA basalts to (minor) HK CA dacites (Lucchi et al., 2013e). Four Eruptive Epochs developed between 246 and 29 ka, separated by quiescence stages, erosional episodes and partial collapses of portions of the volcanic edifices (Lucchi et al., 2013e). Four partially overlapping stratovolcanoes and scoria cones (Casa Fiocria, Fossa Felci, Chiumento and Monte Guardia) were mostly controlled by the WNW-SEE regional tectonic trend, with a progressive shifting of activity toward South-East. A variation of compositions toward the dacitic terms was recorded in the lava domes and coulees and explosive eruptions in the south-eastern sector of Filicudi at ca. 64-50 ka (Monte Terrone, Capo Graziano, Monte Montagnola and Casa dello Zucco Grande; Lucchi et al., 2013e). Volcanic activity of Filicudi has been regulated by the interaction of different evolutionary processes, such as fractional crystallization, crus-




Fig. 23 - (a) Sketch bathymetry of the Aeolian Islands and seamounts, southern Italy (modified from Beccaluva et al., 1985). The trace of the Tindari-Lenocanni fault system (TL) is shown. Depth contour lines in metres below sea level. (b) Schematic chronological framework of Aeolian Islands volcanoes based on the available radiometric ages and correlation of marine terraces and tephra layers (see Lucchi et al., 2013a for age references). (c) SiO_2 vs. K_2O wt% (Peccerillo & Taylor, 1976) classification diagram of volcanic rocks from the Aeolian Islands volcanoes: CA=calcareous; HKCA=high potassium calc-alkaline; SHO=shoshonitic; Bas=basalt; BasAnd=basaltic andesite; And=andesite; Lat=latite; Dac=dacite; Trachy=trachyte; Rhy=rhyolite. Source of data: Alicudi (Lucchi et al., 2013d); Filicudi (Lucchi et al., 2013e); Selene (Lucchi et al., 2013b); Lipari (Forni et al., 2013); Vulcano (De Astis et al., 2013); Panarea (Lucchi et al., 2013c); Stromboli (Francalanci et al., 2013). (d) Panoramic view from the north of the Rocche Rosse obsidian lava flow (rr) effused from the NE side of the Monte Piano pumice cone (mp), the flanks of which are intensely dissected by quarry activity. (e) Aerial view of the submarine gas bursts (arrows) occurred in 2002-2003 in the area of the minor islets; in the background the main island of Panarea is visible.

tal assimilation and mixing. The youngest activity in this area is recorded by the Canna neck and lava flows (29 ka), representing the few remnants of an independent volcanic center offshore the NW coast of Filicudi.

The island of Salina, in the center of the Aeolian archipelago, is the emerged portion of a large volcanic complex developed through time under the influence of the three regional tectonic trends acting in the archipelago (NNW-SSE, WNW-ESE, NE-SW). Six Eruptive Epochs between 244 and 15 ka produced lava flows, Strombolian scoriae and hydromagmatic and subplinian pyroclastic deposits with a CA basaltic to high-K CA dacitic composition (Lucchi et al., 2013b; Nicotra et al., 2014; Sulpizio et al., 2016), interrupted by major periods of quiescence, volcano-tectonic collapses and marine ingression phases during MIS 7 and MIS 5 stages (Lucchi et al., 2013b). The typical twin-volcanoes shape of the island is the result of the superimposition of the Monte dei Porri stratovolcano with the compound volcano formed by the nested Pizzo Capo, Monte Rivi and Monte Fossa delle Felci stratovolcanoes. Magma compositions and their variation through time are the result of contamination of primary magmas with the Calabro-Peloritano lower crust and subsequent differentiation dominated by polybaric fractional crystallization becoming dominant during the activity of Monte Fossa delle Felci and Monte dei Porri, whereas the older Pizzo Capo and Monte Rivi were fed by deep magma reservoirs located at the crust-mantle boundary (Nicotra et al., 2014). During the latest eruptive activities of the Pollara crater (27-15 ka) magma mixing and mingling processes triggered high-energetic basaltic to rhyolitic explosive eruptions.

The island of Panarea (Fig. 23d), together with the Basiluzzo islet, the small rocks of Le Formiche and the minor islets of Dattilo, Panarelli, Liscia Bianca, Bottaro, Liscia Nera are the remnants of CA to high-K CA andesite to dacite lava dome-fields, with minor pyroclastic products, representing the subaerial culminations of a large, mainly submerged, irregularly truncated cone-shaped volcanic complex. Volcanism on Panarea dates back to 155-149 ka, with different growth phases interrupted by erosional unconformities, whereas the latest explosive eruption occurred in the time range between 27-26 and 8.7 ka from an undefined eruptive vent in the area of minor islets (Calanchi et al., 1999; Lucchi et al., 2013c). Although long considered an extinct volcano going through a weak fumarolic phase of activity, Panarea was recently (2002-2003) characterized by signs of a possible eruptive renewal manifesting through spectacular submarine gas bursts occurred in 2002-2003 in the area of the minor islets (Fig. 23d) and accompanied by seismic swarms and emission of high temperature fluids with magmatic geochemical signature (Cairo et al., 2004; Capaccioni et al., 2007).

The island of Lipari is the above-sea-level culmination of a broad, largely submerged volcanic complex belonging to the Vulcano-Lipari-Salina volcanic belt. This volcanic complex is composed of a large variety of lava flows, domes and coulees and pyroclastic successions produced by a spectrum of hydromagmatic, Strombolian to Subplinian and effusive activities, under the control of major regional tectonic trends and recurrent volcano-tectonic collapses. Volcanism has developed between c. 287 ka and the Medieval age, showing progressive West-

East and South-North shifts of the eruptive vents through time, together with an overall chemical change of erupted products from early CA and high-K basaltic andesites to latest rhyolites with a notable gap in the dacites field and a steep increase in the K₂O content through time (Forni et al., 2013). The youngest volcanic eruptions produced the prominent Monte Pilato pumice cone (VIII century CE) (Fig. 23e) and the obsidian lava flows of Rocche Rosse and Forgia Vecchia (XIII century CE), sited in the north-eastern sector of the island, as well as highly dispersed white-coloured, fine-grained rhyolitic tephra layers (Pistolesi et al., 2021). At present, Lipari is in a quiescent stage with a few low-temperature fumaroles and hot springs.

4.4.2. Vulcano volcanic complex (130 ka-Present)

The Island of Vulcano is the exposed summit of a largely submerged volcanic complex that is separated from the nearby Lipari only by a shallow-water and narrow strait of sea. Its development has occurred from about 130 ka to the well-known eruptive cycle of 1888-90 CE which gave origin to the definition of the "Vulcanian" eruptive style (Mercalli & Silvestri, 1891). Volcanism displayed a wide spectrum of eruptive-explosive activities from a number of volcanic edifices which become gradually younger moving north-westwards under control of tectonic structures associated to the NNW-SSE oriented Tindari-Letojanni fault system. The development of the two Il Piano (ca. 100 ka) and La Fossa multi-stage caldera structures (78-8 ka) have the same NNW-SSE elongation and display a progressive NNW-shifting through time (De Astis et al., 2013) (Fig. 24). Erupted products have variable K-alkaline character and degrees of evolution, ranging from basalts to rhyolites and changing through time parallel to the evolution of the volcanic system. It is noteworthy that successive large-scale hydromagmatic eruptions from the La Fossa Caldera emplaced the widely distributed Brown Tufts (70-8 ka) pyroclastic products (Meschini et al., 2020).

The Holocene activity mostly occurred from the La Fossa and Vulcanello eruptive centres (Fig. 24), fed by magmas ranging from shoshonites to rhyolites and alternating each other (De Astis et al., 2013; Nicotra et al., 2018).

The La Fossa stratovolcano (LFO) (Fig. 24a and b), in the middle of La Fossa caldera, was built during the last 5.5 ka through recurrent hydromagmatic to Vulcanian explosive eruptions, which gave rise to multiple dilute pyroclastic currents and fallout deposits, alternating with a few trachytic to rhyolitic lava flows. Compositionally, the LFO products reach the maximum degrees of magma evolution and alkali contents on Vulcano. The progressive growth of the cone occurred during different eruptive cycles interrupted by short intervals of quiescence. After the early activity (5.5-2.9 ka), when most of the present cone was built up to 250-300 m in elevation, a younger eruptive cycle during Roman to early Medieval times produced pyroclastic successions deposited by dilute to concentrated pyroclastic currents, two fallout layers and four lava flows from two different, intersecting craters (De Astis et al., 2013; Malaguti et al., 2022). The activity of the eccentric Forgia crater (Fig. 24c) in Medieval times is still not chronologically well constrained. Another explosive cycle of activity started from 1727 CE and culminated




Fig. 24 - Simplified geological map of Vulcano merged on a shaded relief DEM of the island, showing the areal distribution of the erupted products and vents during its eruptive history, as well as the collapse rims corresponding to the multi-stage formation of the Il Piano and La Fossa calderas (modified from De Astis et al., 2013). Main landforms and calderas rims in the NE shallow-water submarine portion are also shown (from Casalbore et al., 2018). Ago references from De Astis et al. (2013) and references therein. Magma compositions: basalt (Bas), andesite (And), shoshonite (Sho), latite (Lat), trachite (Tr), myiolite (Rhy), high-K (HK), shoshonitic (SHO). (a) Panoramic view of the central sector of the island of Vulcano with the field evidence for the major volcano-tectonic collapse rims (vc1, vc2, vc4, vc6) leading to formation of the multi-stage Il Piano and La Fossa calderas, and the La Fossa stratocone (LFO) standing out in the middle of La Fossa caldera. (b) View from the south of the summit area of LFO showing the crater (cr1) and vents (ve) of the AD 1888-1890 eruption, and its products characterized by bread-crust bombs and lava blocks; this crater is nested within older crater rims (cr2, cr3). The active fumarolic field (arrow) is visible along the northern rim of the AD 1888-1890 crater. (c) Outcrop of the intensely hydrothermally altered rocks of the Faraglione cone (fa) near to the Spiaggia di Levante beach, where intense gas emission in shallow-water areas (arrow) has occurred during the 2021-2022 unrest. In the background, the eccentric Forgia crater (fo) is visible along the northern flank of LFP. (d) View from Lipari of the Vulcanello lava platform (la) and nested scoria cones (sc), with the LFO in the background. All photographs credit F. Lucchi.

ed with the effusion of the outstanding Pietre Cotte phytoclastic lava flow in 1739 CE, the last effusive event from LFO (De Flore, 1922). The Gran Cratere explosive eruptive cycle developed between 1739 CE and the latest paroxysmal activity of 1888-1890 CE, and was characterized by the distinctive emplacement of bread-crust bombs and lava blocks widely dispersed in the summit area of LFO (Fig. 24b). Since the last eruption, the LFO has shown only fumarolic activity, associated with shallow seismicity, fed by a shallow latitic magma body at a depth of 2.5-3.5 km (Paonita et al., 2013; Mandarano et al., 2016). Gas emissions are concentrated in several high-temperature active fumaroles along the northern rim of LFO crater, and subordinate points of emission and hot springs in the submarine areas near the Porto di Levante harbour (Granieri et al., 2006) (Fig. 24c). Unrest periods of increase in temperature above the usual range of 330-400 °C and in the magmatic component of the total gas flux (usually called 'crises') have occurred periodically (1920s, 1987-93, 1996-98, 2004-2009), with the last one started in September 2021 and still ongoing at the time of writing (Aiuppa et al., 2022).

Vulcanello is located along the northern border of La Fossa caldera and consists of three NE-SW-aligned, nested small Strombolian scoria cones from which a lava plateau, made up by a composite lava flow field of pahoehoe- to aa-lavas, formed (Fig. 24d). The oldest reported submarine to shallow-water eruption occurred during Roman times (II century BC, Stothers & Rampino, 1983), whereas most of the subaerial activity took place up to early Medieval times (De Astis et al., 2013; Malaguti et al., 2022), with latest eruptions around the XVI century and fumarolic activity persisting until 1678 CE (Keller, 1980). In the early stages, Vulcanello developed as an independent islet and was successively linked to the main island of Vulcano through the progressive accumulation of ash erupted from La Fossa forming the isthmus area between 1525 CE and 1550 CE (Barbano et al., 2017).

4.4.3. Stromboli composite volcano (85 ka-Present)

Stromboli is a largely submerged composite volcano well-known for its persistent "Strombolian" eruptive activity from historical times. Controlled by a regional NE-SW structural system, the exposed portion of Stromboli was entirely developed from ca. 85 ka to present times (e.g. Gillot and Keller, 1993; Risica et al., 2019), whereas the Strombolicchio neck is dated to ca. 204 ka. The successive epochs of activity, represented by Paleostromboli I-III (85-34 ka), Vancori (26-13 ka), Neostromboli (13-4 ka) and Recent Stromboli (<2.4 ka) (Fig. 25), mainly erupted mafic products (ca. <56 wt% SiO₂) covering a large compositional affinity from calc-alkaline to shoshonitic and potassic-alkaline series (Hornig-Kjaergaard et al., 1993; Francalanci et al., 2013). These epochs were repeatedly interrupted by small, summit calderas and lateral collapses cutting the NW and SE flanks of the volcano down to the submerged basis of the edifice (Tibaldi, 2001; Romagnoli et al., 2009) (Fig. 25a, b). They largely conditioned the location of active vents and erupted products, with the current Sciara del Fuoco (SDF) scar developed during the last 13 ka. The activity has been mostly supplied from summit vents through alternating effusive and Strombolian eruptions, whilst more in-

tense Subplinian explosions only characterized the Paleostromboli I and Vancori epochs, and subordinate (but notable) flank eruptions along the NE and SW buttressed flanks of the volcano took place during the Neostromboli one. Recurrent NW-dipping lateral collapses occurred during the Holocene in alternation with sequences of eruptions, and caused the unloading of the magmatic-hydrothermal system, likely triggering atypical hydromagnetic eruptions and pyroclastic density currents (PDC) that affected the lower slopes of the volcano (Lucchi et al., 2018). The Recent Stromboli epoch started during Roman times and resulted in the construction of the summit Pizzo scoria cone, whilst the San Bartolo fissure erupted pahoehoe lava flows that largely invaded the NE coastal sector forming a fan-shaped delta, and represented the last effusive activity outside the SDF. The present-day activity of Stromboli, starting from (at least) early Medieval times, has occurred from a crateric terrace located at elevations of c. 750 m a.s.l. near the SDF headwall (Rosi et al., 2013)(Fig. 25c). This activity is typically characterized by continuous active degassing (puffing) and rhythmic, discrete, weak to moderate explosions lasting a few seconds (Fig. 25d) that eject dark highly-porphyritic (hp) scoriaceous lapilli, bombs and ash, accumulating in the area around the craters and along the steep SDF scar. Occasional more violent explosions, covering a large intensity range from major explosions to paroxysms (Bevilacqua et al., 2020), produce convective plumes rising up to 7-8 km (Fig. 25e) and eject mingled scoriaceous hp and "pumiceous", light-coloured and low-porphyritic lapilli, bombs and spatter, together with lithic blocks derived from deeper craterization of the upper conduit and vent system. Ballistic fallout can overcome the SDF walls impacting the flanks of the cone, and occasionally reaching the inhabited areas of Stromboli and Ginostra, whereas short-lived PDCs due to column collapse generally flow within the SDF (Fig. 25f). During historical times, paroxysms occurred repeatedly in the XVI-XVII and the XX centuries, with the more energetic event in 1930 and 1944, whilst recent paroxysms occurred on 5 April 2003, 15 March 2007, and the close events of 3 July and 28 August 2019 (Giordano & De Astis, 2021; Melrich et al., 2021; Ripepe et al., 2021; Viccaro et al., 2021). Effusive eruptions from summit craters or lateral vents/fissures opened at various elevations within SDF, recurrently interrupt the 'normal' Strombolian activity, lasting from ca. 1 day to a few months (e.g. in 2002-2003, 2007, 2014) (Fig. 25g). For (at least) the past three centuries, lava flows have descended the steep slopes of SDF, and eventually reached the sea (e.g. in 2007, 2014, 2022), representing a negligible hazard for people and infrastructures. Up to the Middle Age, lava flows have episodically surrounded the SDF lateral rims and expanded along the upper to lower flanks of the cone.

Recurrent gravitational instabilities and collapses at various scales along the oversteepened slopes of SDF are recorded during the last century (Barberi et al., 1993), eventually resulting in tsunami waves along the Stromboli and surrounding coasts, as observed associated to the December 2002 submarine and subaerial landslide developed along the SDF (Tinti et al., 2005).




Fig. 25 - Simplified geological map of Stromboli (modified from Francalanci et al., 2013) and morpho-structural sketch map (a) of its submerged flanks (modified from Romagnoli et al., 2013). Altitude points are in metres above sea level. The map is based on a DEM-shaded relief image courtesy of DICEA (Dipartimento di Ingegneria Civile Edile e Ambientale), Università di Roma La Sapienza (research project funded by the Italian Department of Civil Protection). Age references from Francalanci et al. (2013) and references therein. Magma compositions: basalt (Bas), basaltic-andesite (Bas-And), andesite (And) shoshonite (Sho), latite (Lat), trachyte (Tr), high-K (HK), shoshonitic (SHO). (b) Aerial view of the Stromboli composite volcano and the Scira del Fuoco collapse scar cutting its NW flank, with the active craters visible near its headwall (Credit Italian Civil Protection). (c) Aerial view of the summit area of the Stromboli composite volcano showing the NE-SW aligned active craters near the headwall of the Scira del Fuoco collapse scar (Credit INGV). (d) Strombolian activity of the summit craters (Credit M. Pistolesi). (e) Eruptive column formed during the 5 April 2003 paroxysm (Credit INGV). (f) Pyroclastic currents formed along the Scira del Fuoco collapse during the 3rd July 2019 paroxysm (unknown author). (g) Pahoehoe lava flows forming a lava delta along the Scira del Fuoco collapse during the 2007 effusive activity (Credit C. Romagnoli).

4.4. Mt. Etna (500 ka to Present) (S.B., M.G., E.N., M.V.)

The eruptive history of Mt. Etna began about 500 ka (De Beni et al., 2011) in the eastern Sicily, an area of geodynamic complexities, where the Apennine-Maghrebian Chain overlaps the undeformed margin of the African continental plate, the Hyblean Foreland, bounding westwards the oceanic Ionian lithosphere (Lentini, 1982; Cristofolini et al., 1985; Branca et al., 2004; 2008; 2011a,b; Fig. 26).

Volcanic activity in the area developed after an early long period of scattered basaltic magmatism in the Hyblean Plateau that migrated northwards during Middle-Late Pleistocene towards the region of the present Mt. Etna (Neri et al., 2018). Volcanism in this area began with fissure-type eruptions with tholeiitic affinity emplacing lava flows in submarine and subaerial environment (Tanguy et al., 1978). Remnants of the submarine activity are pillow-lavas and columnar outcrops on the eastern shore north of Catania (about 500 ka; Gillot et al., 1994; Corsaro & Cristofolini, 1997; 2000; De Beni et al., 2011; Fig. 26). Subaerial tholeiites date back to 330 ka (De Beni et al., 2011) and mostly crop out on the SW slopes of the volcano (Fig. 26). The erupted lavas gradually changed in composition from subalkaline towards Na-alkaline, while the fissure activity mainly concentrated on the Ionian coast along the Timpe fault system starting from about 220 ka (Tanguy et al., 1997; Corsaro & Cristofolini, 1997; Branca et al., 2004; 2008; 2011a,b; Fig. 26).

During the last 130 ka the major eruptive axes migrated westwards, gradually aligning themselves on a central axis. This transition from fissure to central vent activity led to the construction of a series of central-conduit edifices producing Strombolian to Plinian eruptions in the area of the present-day Valle del Bove. As the central activity stabilized at ca. 57 ka (De Beni et al., 2011), a huge stratovolcano, the Etna, grew up. The intense eruptive activity both effusive and highly explosive of the Etna created a roughly 3600 m-high cone, which constitutes the frame of the present edifice of Mt. Etna (Fig. 26). The Etna activity ended 15 ka ago after four Plinian eruptions (Coltell et al., 2000; Del Carlo et al., 2017) that created a 4 km-wide caldera. This became the site of persistent volcanic activity that progressively led to the emplacement of a new stratovolcano, the Mongibello, over the last 15 ka (Fig. 26). The formation of the Valle del Bove, a horseshoe depression about 7 x 4.5 km wide that is located on the eastern flank of Mt. Etna, dates back to the activity of the Mongibello stratovolcano, namely to 7478 - 7134 BCE (Malaguti et al., 2023).

The activity of Mongibello marks the beginning of the recent volcanic history of Mt. Etna, which also displayed a wide range of eruptive styles from purely effusive to violent explosive (Branca & Del Carlo, 2004). Historical records of eruptions date back to the Greek colonization of Sicily (734 BCE), passing through other chronicles during the Roman Age and the XVII century (Branca & Abate, 2019) with reference, respectively, to the 122 BCE Plinian eruption (Coltell et al., 1998) and the impressive 1669 eruption (Corsaro et al., 1996; Nicotra & Viccaro, 2012; Kahl et al., 2017). The last few centuries were dominated by effusive and Strombolian eruptions (Branca & Del Carlo, 2005) with emission of ha-

watic products and subordinate mugearites (Viccaro & Cristofolini, 2008). Over the last five decades, the eruption frequency and explosivity increased, while the composition of volcanic products shifted towards K-trachybasalts (Tanguy et al., 1997; Tonarini et al., 2001; Viccaro et al., 2011; Viccaro & Zuccarello, 2017).

Mt. Etna has been particularly active during the last 30 years, experiencing continuous morphological modifications (Harris et al., 2011). Most of the morphological changes were caused by the opening of eruptive fissures on the flanks of the volcano that produced lava flows, as in 1991-1993, 2004-05 and 2008-09, and large scoria cones, as during the 2001 and 2002-2003 eruptions (Tonarini et al., 1995; Ciochetti et al., 2004; Métrich et al., 2004; Andronico et al., 2005; Allard et al., 2006; Ferlito et al., 2012; Palano et al., 2017). Since nineties, there was also a clear growth in the number of paroxysmal eruptions from the summit craters (Fig. 26), which generated extensive eruptive columns, abundant tephra fallouts and ash-lapilli dispersion up to hundreds of km far from craters (Andronico et al., 2021). Main paroxysmal episodes occurred in short-term sequences of multiple events lasting weeks to years. Remarkable sequences were in 1996-1999 (23 events in 6 months), 2000 (64 events in 6 months) and 2007 (7 events in 7 months).

The last decade has also seen the development of some of the most energetic, yet long-lasting paroxysmal sequences of recent times, which have been fed by the most volatile-rich magmas ever found at Mt. Etna (Zuccarello et al., 2021). These include a three-year-long sequence of 44 lava fountains during 2011-2013 (Behncke et al., 2014; Giuffrida & Viccaro, 2017; Giuffrida et al., 2018; Zuccarello et al., 2022), and the very recent sequence of 62 events that took place between December 2020 and February 2022. These sequences alternate with periods of dominant effusive activity from both the summit and the volcano flanks (Viccaro et al., 2019; Borzi et al., 2020; Giuffrida et al., 2021), with the only brief interlude of two short, but powerful paroxysmal series in December 2015 (4 events) and May 2016 (3 episodes) (Corsaro et al., 2017; Calvari et al., 2018; Cannata et al., 2018; Zuccarello et al., 2022).

The recurrence of highly explosive events at the summit, which produce severe tephra fallout and episodic formation of pyroclastic density currents due to partial collapses of the South East Crater cone (Behncke, 2009a; Ferlito et al., 2010; Scollo et al., 2013; Costa et al., 2023), and effusive flank eruptions from vents located at low altitude (Del Negro et al., 2013) have the potential to cause significant socio-economic damages because of the high probability to impact the densely populated areas around the volcano, as well as hundreds of thousands of tourists who visit Mt. Etna yearly.

4.5. The Hyblean volcanism (M.G., M.V.)

The Hyblean volcanism began in Upper Triassic ca. 200 Ma (Cristofolini, 1986) and continued until Lower Pleistocene, with both submarine and subaerial episodes of mafic volcanism widely scattered across the area of southeastern Sicily (Carbone & Lentini, 1981; Schmincke et al., 1997). No volcanism is known to have occurred from the Upper Cretaceous to the Middle-Upper Miocene. The Messinian volcanism produced lavas of Na-alkaline affinity that outcrop in the southernmost and eastern part




Fig. 26 - Synthetic geological map of Mt. Etna volcano with the main morphotectonic features (upper panel, modified after Branca et al., 2011); a scheme of the stratigraphic relationships of the volcanic succession using the supersyntem, syntem and lithosomes classifications and the radiocarbon ages have been also reported down-left in the figure (modified after Branca et al., 2011b) and aerial view of the Mt. Etna summit area taken on January 2021 with the 5 active craters, namely (lower panel): Voragine (VOR), North East Crater (NEC), Bocca Nuova 1 and Bocca Nuova 2 (BN1 and BN2), South East Crater (SEC).

of the Hyblean Plateau. A hiatus of ca. 50 Ma separates these earlier emissions from the later activity that resumed during Miocene in the central-northern area of the plateau with eruptions of both tholeiitic and alkaline magmas (De Rosa et al., 1991; Tonarini et al., 1996; Beccaluva et al., 1998; Trus et al., 1998). The Neogene-Quaternary volcanism includes two different eruptive cycles: a) the Miocene cycle, prevalently alkaline in composition (mostly basanites and alkali-olivine basalts), which is dominated by explosive eruptions testified by diatreme-related volcanoclastic deposits bearing a significant amount of mantle xenoliths (Scribano, 1987a; b; Scribano et al., 2009; Suttorp & Schmincke, 2010; 2012); b) the Plio-Pleistocene cycle almost exclusively characterized by relatively basic magmas, ranging in composition from extremely silica-undersaturated alkaline magmas to silica-oversaturated tholeites. The Plio-Pleistocene cycle was the most widespread, taking place over an area of ca. 500 km² at the northern margin of the Hyblean Plateau. Large volumes of tholeiitic products were erupted in a shallow marine environment during the Upper Pliocene, followed by at least five episodes of alkali basaltic volcanism between the end of the Upper Pliocene and the Lower Pleistocene. The first of these episodes emplaced in the northernmost sector of the Hyblean Plateau a considerable volume of alkali basalts as pillow breccias, subaerial lavas and scoria deposits (Behncke, 2009b), while the latest eruptions (ca. 1.4 Ma; Schmincke et al., 1997) emitted scarce volumes of basanitic and nephelinitic lavas.

4.6. The Sicily Channel

(A.C., M.G., S.R., M.V.)

4.6.1. General background

The Sicily Channel is set in the thinned continental crust of the Pelagian block (Fig. 27a), a rifted zone in the Sicilian Maghrebian Chain foreland (Catalano et al., 2014). The main morphotectonic features are three deep tectonic troughs, NW-SE oriented, namely the Pantelleria, Linosa and Malta grabens, which exert control on magma generation and ascent (Fig. 27a). Magmatism in the Sicily Channel is widespread and results in the occurrence of several volcanic submarine centres (Civile et al., 2006; Rotolo et al., 2006) and two subaerial volcanic complexes, the Pantelleria (Fig. 27b) and Linosa islands. Submarine volcanism began around 10 Ma, with the latest event of the 'Ferdinandea' ephemeral islet emersion in 1831 CE and the 1891 CE 'Fœstner' entirely submarine eruption (Conte et al., 2014; Coltell et al., 2016; Kelly et al., 2014). Submarine volcanic rocks are mafic and dominantly mildly alkalic regarding the serial affinity (Rotolo et al., 2006; White et al., 2020). Ascent of mafic magmas is driven by tectonic structures, with the exception of Pantelleria, where the route to the surface is facilitated in the northern part of the island (Gluffrida et al., 2020) and inhibited in the central sector, with production of abundant felsic derivative products.

4.6.2. Pantelleria (320-4 ka)

The subaerial volcanic activity at Pantelleria started around 320 ka and is characterized by emission of pantelleritic products emplaced as lava flows and pumice fall-out deposits (Civetta et al., 1984; Mahood & Hildreth, 1986; Rotolo et al., 2013; Jordan et al., 2018). Nine ignimbrite

eruptions covered the island (entirely to partly) in the age interval 181-46 ka and alternated with inter-ignimbrite periods, during which tens of local eruptive centers were active (Jordan et al., 2018; Rotolo et al., 2021; Fig. 27c). Two ignimbrites out of nine, namely the Capra and Green Tuff Formations (age 140 and 46 ka, respectively) are associated with caldera collapses and related scarp remnants (La Vecchia and Cinque Denti caldera, respectively; Fig. 27b; Speranza et al., 2012; Rotolo et al., 2013). Other five potential caldera collapses (now buried) have been hypothesized based on the occurrence of thick pyroclastic breccia horizons within or at the top of some ignimbritic deposits (Jordan et al., 2018). Over the period 181-46 ka, basaltic manifestations were rare and limited only to some lava flow units dating back to 118 ka and 83 ka (Civetta et al., 1984) that were found in the northwestern part of the Island (Fig. 27c). The post-Green Tuff volcanic activity was dominated by low-energy Strombolian eruptions and effusive events, mostly inside and along the rim of the Cinque Denti caldera. This activity produced pumice fall deposits, lava flows and domes of dominant pantelleritic compositions (Civetta et al., 1988; Rotolo et al., 2007). An updated stratigraphy of the post-Green Tuff evolution is given in Rotolo et al. (2021) based on high resolution ⁴⁰K/³⁹Ar ages (Scialletti et al., 2011) and paleomagnetic data (Speranza et al., 2010), with two periods of activity after the Green Tuff eruption, instead of the five proposed by Civetta et al. (1988).

Basaltic volcanism accounts for ca. 5 % of outcropping rocks and is centered in the NW sector of the island with short lava flows and scoria cones (Gluffrida et al., 2020). Mafic volcanism is dated (K/Ar) in the age interval 31±13 ka and 27±10 ka, with possible younger ages close to 10 ka (unpublished data; Civetta et al., 1984; Fig. 27d).

4.6.3. Linosa (1.06-0.5 Ma)

Linosa is a small volcanic island (ca. 6 km²) whose subaerial portions represent only 4% of the entire volcanic edifice, which extends for ca. 20 km along the NW-SE tectonic lineament characterizing the whole Sicily Channel (Fig. 28; Romagnoli et al., 2020). While no age data are available for its submarine activity, the subaerial eruptive history can be constrained between ca. 1.06 and 0.5 Ma, during which volcanism was effusive to explosive, mostly hydromagmatic, and occurred over three major periods (Rossi et al., 1996). Erupted magmas show limited compositional variation, being only mafic (alkali basalts to hawaiites/mugearites), slightly more primitive than the Pantelleria basalts. Rare intrusive lithics in pyroclastic products have more evolved, syenitic compositions never erupted as magma.

4.7. Seamounts of the Tyrrhenian Sea (6 Ma-Present)

(R.D.R., S.d.V., P.D., F.L., E.N., R.S., C.A.T., M.V.)

The Tyrrhenian Sea started its opening 15 Ma during the Apennine orogenesis as a result of back-arc extensional processes. In the southern sector, the roll-back of the Ionian slab caused the SE migration of the Calabrian Arc and accelerated the process of back-arc extension, leading to the progressive thinning and substitution of the continental crust with a new oceanic one. This process is responsible for the opening of the Vavilov (6




Fig. 27 - (a) Schematic structural map of the Sicily Channel (PT: Pantelleria through; MT: Melite through; LT: Linosa through; GB: Graham Bank) and (b) Geological and structural sketch map of the island of Pantelleria (modified after Catalano et al., 2014); (c) volcanic succession in the area of Kazen at Pantelleria: pre-Green Tuff scoriaeous basaltic rocks are visible at the bottom of the succession, which is constituted in the upper portion by other pre-Green Tuff ignimbrites and finally by the Green Tuff ignimbrite; (d) Pantelleria Island (Safo La Vecchia): La Vecchia Caldera scarp. The caldera wall dips 80° NW (left) and separates flat-lying pre-caldera ignimbrites (to the right of the caldera wall), with concordant post-caldera ignimbrites and one fallout deposit. The sequence is topped by the Green Tuff ignimbrite (ago 46 ka), the last of the nine Pantelleria ignimbrites; (e) Linosa Island (Punta Galcarella): hydromagnetic and fallout deposits of three major volcanic phases.

or 4.5 - 2.5 Ma) and Marsili (1.8 Ma-pres.) oceanic basins and the birth and growth of a great number of seamounts (Fig. 28a; Trua et al., 2004), whose ages decrease from NW toward SE.

Marsili is a huge (70x30 km) volcanic submarine complex, elongated along a N-S direction with its summit placed 3 km above the seafloor. Although the oldest products date back to 1 Ma, Marsili is considered an active volcanic system due to: i) seismic activity typical of volcanic and hydrothermal areas (D'Alessandro et al., 2012); ii) volcanic gases detected above the water column (Lupton et al., 2011); iii) the products of the two eruptive episodes occurred 3000 and 5000 years ago, recognized at its top (Iezzi et al., 2014). A temporal progression characterizes the Marsili magmatic activity from

IAB-like basalts with calc-alkaline affinity to OIB-like basalts (Trua et al., 2007). The summit area is marked by volcanic cones perfectly preserved, with some evidence of lava flow fields and volcano-tectonic structures (Ventura et al., 2013). Magnaghi and Vavilov are the other two important seamounts of the Tyrrhenian Sea, with an elongated shape over 20 km. Magnaghi seamount is related to the ancient activity (3-2.7 Ma; Serri et al., 2001), whereas Vavilov seamount formed during the Quaternary (0.73-0.1 Ma) and it could still be considered active (Savelli & Ligi, 2017). Although they lie on a seafloor with MORB-like geochemical signature (Trua et al., 2007), the few rocks collected at Magnaghi and Vavilov seamounts show a mildly alkaline OIB-like geochemical affinity (Trua et al., 2007). Palinuro seamount, consid-




Fig. 2B - (a) Morphometric map of the Tyrrhenian Sea; seamounts are labeled in white, whereas emerged volcanoes in yellow; (b) Simplified geological map of Ustica; (c) Panoramic view of the western coast of the Ustica island and of the Monte Costa del Fallo stratocone (Eruptive Epoch 2); (d) Panoramic view of the northern scarp of the Ustica island and of lava and scoriae belonging to Eruptive Epoch 3.

ered active or quiescent (Milano et al., 2012), presents a 50 km E-W elongated shape, resulting from the superimposition of different volcanic edifices. The composition of emitted products is very similar to the emerged portion of the Aeolian Islands (Trua et al., 2004).

4.6. Ustica (730-ca.130 ka)

(R.D.R., S.d.V., P.D., F.L., E.N., R.S., C.A.T., M.V.)

The island of Ustica is the exposed summit of a largely submerged NE-SW elongated volcanic complex rising about 2000 m above the Tyrrhenian seafloor, including an extensive field of submerged volcanic cones (up to thirty) to the north of the island. It is located over a transitional zone between a northern domain characterized by thinned oceanic crust (ca. 8 km thick) and a variably thick (25-30 km) western-southern domain of continental crust (Savelli, 1986; Sulli, 2000).

Based on literature data and new unpublished studies, the eruptive history of the Island of Ustica, entirely developed during the Quaternary (de Vita et al., 1998; Bonomo & Ricci, 2010), is arranged into five epochs of activity (Fig. 28b) that emplaced submarine to subaerial volcanic rocks with a Na-alkaline signature. The first epoch (730-516 ka) includes submarine basaltic lavas emitted from fissural vents and basaltic to hawaiitic and mugearitic lavas relevant to the large Mt. Guardia dei Turchi stratovolcano, which progressively emerged from the sea. Basaltic to hawaiitic and mugearitic lava flows and pyroclastic products of the second epoch (490-441 ka) are related to the Monte Costa del Fallo stratovolcano and its lateral eruptive vents, partly located offshore the west coast of the island. The third epoch (424-412 ka) is characterized by the emplacement of hawaiitic/mugearitic lava flows (Fig. 28c and d) and trachytic pumice pyroclastic products from the Monte Costa del Fallo stratovolcano, together with the products of a series of lateral vents including the trachytic, amphibole-bearing sub-intrusive body exposed at Punta San Paolo (Aletti et al., 2005). The fourth eruptive epoch was fed by lateral dykes in the NW sector of Monte Costa del Fallo stratovolcano that emplaced hawaiitic to mugearitic lava flows and minor strombolian deposits. Finally, the fifth eruptive epoch includes the alkali-basaltic tuff cone and lava neck of Capo Falconiera, which was active at about 130 ka (de Vita et al., 1998). The whole island of Ustica has been eroded and reworked by different orders of marine terraces, whereas the volcanic products of Capo Falconiera are interlayered within marine terraces of the Last Interglacial (de Vita et al., 1998).

ACKNOWLEDGMENTS

Biagio Giaccio and Paola Petrosino, together with all the co-authors, would like to thank an anonymous reviewer whose valuable suggestions helped to improve the first version of the ms. They are deeply indebted to the associate editor Ilaria Mazzini for inviting them to design and coordinate this review paper on the Italian Quaternary volcanism.

REFERENCES

- Acocella, V., Funiciello, R. (2002) - Transverse structures and volcanic activity along the Tyrrhenian margin of central Italy. *Memorie della Società Geologica Italiana*, 1, 739-747.
- Acocella V., Funiciello R. (2006) - Transverse systems along the extensional Tyrrhenian margin of central Italy and their influence on volcanism. *Tectonics*, 25, TC2003.
- Acocella V., Porreca M., Neri M., Matti M., Funiciello R. (2006) - Fissure eruptions at Mount Vesuvius (Italy): insights on the shallow propagation of dikes at volcanoes. *Geology*, 34, 673-676. Doi: 10.1130/G22552.1
- Acocella V., Palladino D.M., Ciolfi R., Russo P., Simei S. (2012) - Caldera structure, amount of collapse and erupted volumes: the case of Bolsena Caldera, Italy. *Geological Society of America Bulletin*, 124(9-10), 1562-1576.
- Aluppa A., Bitetto M., Calabrese S., Delle Donne D., Lages J., La Monica F.P., Chioldi G., Tamburello G., Cotteri A., Fulignati P., Gioncoada A., Liu E.J., Moretti R., Pistolesi M. (2022) - Magmatic magma feeds degassing unrest at Vulcano Island, Italy. *Communications Earth & Environment*, 3, 255.
- Alagna K., Peccerillo A., Martin S., Donati, C. (2010) - Tertiary to Present Evolution of Orogenic Magmatism in Italy. In: Beltrando M., Peccerillo A., Matti M., Conticelli S., Doglioni C. (Eds.) *The Geology of Italy: tectonics and life along plate margins*, Journal of the Virtual Explorer, 36, paper 18. Doi: 10.3909/jvtx.2010.00233
- Albert P.G., Tomlinson E.L., Smith V.C., Di Traglia F., Pistolesi M., Morris A., Donato P., De Rosa R., Sulipizio R., Keller J., Rosi M., Menzies M. (2017) - Glass geochemistry of pyroclastic deposits from the Aeolian Islands in the last 50 ka: a proximal database for tephrochronology. *Journal of Volcanology and Geothermal Research*, 336, 81-107.
- Albert P.G., Giaccio B., Isaia R., Costa A., Niespolo E.M., Nomade S., Pereira A., Renne P.R., Hinchliffe A., Mark D.F., Brown R.J., Smith V.C. (2019) - Evidence for a large-magnitude eruption from Campi Flegrei caldera (Italy) at 29 ka. *Geology*, 47 (7), 595e599. Doi: 10.1130/G45805.1
- Albini A., Cristofolini R., Di Girolamo P., Stanzione D. (1980) - Rare-earth and other trace-element distributions in the calc-alkaline volcanic rocks from deep boreholes in the Phleorean Fields, Campania (south Italy). *Chemical Geology*, 29, 123-133.
- Alessio M., Bella F., Improta S., Belluomini G., Calderoni G., Cortesi C., Turi B. (1979) - University of Rome Carbon-14 dates XIV. *Radiocarbon*, 18, 321-349.
- Allard P., Behncke B., D'Amico S., Neri M., Gambino S. (2006) - Mt. Etna 1993-2005: anatomy of an evolving eruptive cycle. *Earth Science Reviews*, 78, 85-114.
- Aletti M., Pomilio M., Rotolo S.G. (2005) - Melitic and ultramafic enclaves in Ustica Island lavas: Inferences on composition of lower crust and deep magmatic processes. *Lithos*, 84 (3-4), 151-167.
- Amato V., Auocelli P.P.C., Cesario M., Jicha B., Lebreton V., Orain R., Pappone G., Petrosino P., Russo Ermoli E. (2014) - Quaternary evolution of the largest intermontane basin of the Molise Apennine (central-southern Italy). *Rendiconti Lincei. Scienze Fisiche e Naturali*, 25(2), 197-216.

- Doi: 10.1007/s12210-014-0324-y
 Andronico D., Cioni R. (2002) - Contrasting styles of Mount Vesuvius activity in the period between the Avellino and Pompei Plinian eruptions, and some implications for assessment of future hazards. *Bulletin of Volcanology*, 64, 372-391.
 Doi: 10.1007/s00445-002-0215-4
- Andronico D., Branca S., Calvari S., Burton M., Cefalù T., Corsaro R.A., Del Carlo P., Garfì G., Lodato L., Miraglia L., Murè F., Neri M., Peccora E., Pompilio M., Salerno G., Spampinato L. (2005) - A multi-disciplinary study of the 2002-03 Etna eruption: insights into a complex plumbing system. *Bulletin of Volcanology*, 67, 314-330.
- Andronico D., Carnàtia A., Di Grazia G., Ferrari F. (2021) - The 1986-2021 paroxysmal episodes at the summit craters of Mt. Etna: insights into volcano dynamics and hazard. *Earth Science Reviews*, 220, 103686.
- Angiulli G., De Francesco A.M., Lomonaco L. (1985) - Nuovi dati geochemica sull'isola di Ischia. *Mineralogica et Petrografia Acta*, 29, 41-60.
- Arienzo I., Moretti R., Civetta L., Orsi G., Papeo P. (2010) - The feeding system of Agnano-Monte Spina eruption (Campi Flegrei caldera, Italy): dragging the past into present activity and future scenarios. *Chemical Geology*, 270, 135-147.
 Doi: 10.1016/j.chemgeo.2009.11.012
- Armenti P., Barberi F., Bizouard H., Cicchiatti R., Innocenti F., Metrich N., Rosi M., Sbrana A. (1983) - The Phlegraean Fields: Magma evolution within a shallow chamber. *Journal of Volcanology and Geothermal Research* 17, 289-301.
- Arrighi S., Principe C., Rosi M. (2001) - Violent strombolian and subplinian eruptions at Vesuvius during post-1631 activity. *Bulletin of Volcanology*, 63, 126-150.
 Doi: 10.1007/s004450100130
- Avanzinelli R., Elliott T., Tommasini S., Conticelli S. (2008) - Constraints on the genesis of the potassium-rich Italian volcanics from U/Th disequilibrium. *Journal of Petrology*, 49, 195-223.
- Avanzinelli R., Lustrino M., Mattei M., Melluso L., Conticelli S. (2009) - Potassic and ultrapotassic magmatism in the circum-Tyrrhenian region: the role of carbonated pelitic vs. pelitic sediment recycling at destructive plate margin. *Lithos*, 113, 213-227.
- Avanzinelli R., Cioni R., Conticelli S., Giordano G., Isala R., Mattei M., Melluso L., Sulpizio R. (2017) - The Vesuvius and the other volcanoes of Central Italy. *Geological Field Trips and Maps*, 9: No.1. 1, pp. 158.
 Doi: 10.3301/GFT.2017.01
- Avanzinelli R., Benvenuti M.G., Brogi A., Citelli F., Cioni R., Conticelli S., Laurenzi M.A., Liotta D., Marroni M., Pandolfi L., Papini M., Paternostro S., Rook L., Sepulveda Birke J.P., Tavarelli E., Valeriani L., Vacelli O. (2022) - Itinerario 9 - Baccinello, Monte Amiata, Val d'Orcia e Monte Cetona. In: *Guide Geologiche Regionali - Toscana* (Conti P., Conticelli S., Cormamusini G., Marroni M. (Edts). Società Geologica Italiana, Roma, 15, 243-267. ISBN: 9-788894-484434.
- Aucelli P.P.C., Mattei G., Caporizzo C., Di Luccio D., Tursi M.F., Pappone G. (2022) - Coastal vs volcanic processes: Procida Island as a case of complex morpho-evolutive response. *Marine Geology*, 106814.
 Doi: 10.1016/j.margeo.2022.106814
- Ballini A., Barberi F., Laurenzi M.A., Mezzetti F., Villa I.M. (1989a) - Nuovi dati sulla stratigrafia del vulcano di Roccamontagna. *Bollettino GNV*, 5, 533-555.
- Ballini A., Frullani A., Mezzetti F. (1989b) - La formazione piroclastica del Tufo Trachitico Bianco del vulcano di Roccamontagna. *Bollettino GNV*, 5, 557-574.
- Barbano S., Castelli V., Pirrotta C. (2017) - Materiali per un catalogo di eruzioni di Vulcano e di terremoti delle isole Eolie e della Sicilia nordorientale (secc. XV-XIX). *Quaderni di Geofisica*, 142.
- Barberi F., Ferrara G., Innocenti F., Marinelli G., Mazzuoli R. (1971) - A magmatic province of anatetic origin: The Tuscan-Latian province (Italy). Abstract, XV UCGI Congress, Moscow.
- Barberi F., Innocenti F., Lirer L., Munno R., Pescatore T.S., Santacroce R. (1978) - The Cl: a major pre-Historic eruption in the Neapolitan area (Italy). *Bulletin of Volcanology*, 41, 1-22.
- Barbieri M., Di Girolamo P., Locardi E., Lombardi G., Stanzione D. (1979) - Petrology of the calc-alkaline volcanics of the Parete 2 well (Campania, Italy). *Periodico di Mineralogia*, 48, 53-74.
- Barberi F., Innocenti F., Landi P., Rossi U., Saitta M., Santacroce R., Villa I.M. (1984) - The evolution of Latena caldera (central Italy) in the light of subsurface data. *Bulletin of Volcanology*, 47, 125-141.
- Barberi F., Rosi M., Sodini A. (1993) - Volcanic hazard assessment at Stromboli based on review of historical data. *Acta Vulcanologica*, 3, 173-187.
- Barberi F., Buonasorte G., Cioni R., Fiorelli A., Foresi L., Iaccarino S., Laurenzi M.A., Sbrana A., Vernia L., Villa I.M. (1994) - Plio-Pleistocene geological evolution of the geothermal area of Tuscany and Lazio. *Memorie Descrittive della Carta Geologica d'Italia*, 49, 77-134.
- Barberi F., Contelli M., Frullani A., Rosi M., Almeida E. (1995) - Chronology and dispersal characteristics of recently (last 5000 years) erupted tephra of Cotopaxi (Ecuador): implications for long-term eruptive forecasting. *Journal of Volcanology and Geothermal Research*, 69, 217-239.
- Bartole R. (1984) - Tectonic structure of the Latian-Campanian shelf (Tyrrhenian Sea). *Bollettino di Oceanografia Teorica e Applicata*, II, 3, 197-230.
- Bear A.N., Cas R.A.F., Giordano G. (2009a) - The implications of spatter, pumice and lithic clast rich proximal co-ignimbrite lag breccias on the dynamics of caldera forming eruptions: the 151 ka Sutri eruption, Vico Volcano, Central Italy. *Journal of Volcanology and Geothermal Research*, 181, 225-255.
- Bear A.N., Cas R.A.F., Giordano G. (2009b) - Variations in eruptive style and depositional processes associated with explosive, phonolitic composition, caldera-forming eruptions: the 151 ka Sutri eruption, Vico Caldera, central Italy. *Journal of Volcanology and Geothermal Research*, 184, 1-24.
- Beccaluva L., Gabbianelli G., Lucchini F., Rossi P.L., Savelli C. (1985) - Petrology and K/Ar ages of volcanics dredged from the Aeolian seamounts: impli-

- cations for geodynamic evolution of the Southern Tyrrhenian basin. *Earth Planetary Science Letters*, 74, 187-208.
- Beccaluva L., Siena F., Coltorti M., Di Grande A., Lo Giudice A., Macciotta G., Tassanini R., Vaccaro C. (1998) - Nephelinitic to tholeiitic magma generation in a transcurrent tectonic setting: an integrated model for the Iblean volcanism, Sicily. *Journal of Petrology*, 39, 1547-1576.
- Beccaluva L., Coltorti M., Di Girolamo P., Melluso L., Milani L., Morra V., Siena F. (2002) - Petrogenesis and evolution of Mt. Vulture alkaline volcanism (Southern Italy). *Mineralogy and Petrology*, 74, 277-297.
- Behncke B. (2009a) - Hazards from pyroclastic density currents at Mt. Etna (Italy). *Journal of Volcanology and Geothermal Research*, 180, 148-160.
- Behncke B. (2009b) - Late Pliocene volcanic island growth and flood basalt-like lava emplacement in the Hyblean Mountains (SE Sicily). *Journal of Geophysical Research*, 109, B09201.
- Behncke B., Branca S., Corsaro R.A., De Beni E., Miraglia L., Proietti C. (2014) - The 2011-2012 summit activity of Mount Etna: birth, growth and products of the new SE crater. *Journal of Volcanology and Geothermal Research*, 270, 10-21.
- Bellucci F., Lirer L., Munno R. (1999) - Geology of Penza, Ventotene and Santo Stefano islands (with a 1:15,000 scale geological map). *Acta Vulcanologica*, 11, 197-222.
- Beltrando M., Rubatto D., Manatschal G. (2010) - From passive margins to orogens: The link between ocean-continent transition zones and (ultra)high-pressure metamorphism. *Geology*, 38(6), 559-562.
- Bertagnini A., Sbrana A. (1986) - Il vulcano di Vico: stratigrafia del complesso vulcanico e sequenze eruttive delle formazioni piroclastiche. *Memorie della Società Geologica Italiana*, 35, 699-713.
- Bertagnini A., de Rita D., Landi P. (1995) - Mafic inclusions in the silica-rich rocks of the Tofa-Ceriti-Manziana volcanic district (Tuscan Province, Central Italy): chemistry and mineralogy. *Mineralogy and Petrology*, 54, 261-278.
- Bertagnini A., Cioni R., Guidoboni E., Rosi M., Neri A., Boschi E. (2008) - Eruption early warning at Vesuvius: The A.D. 1631 lesson. *Geophysical Research Letters*.
Doi: 10.1029/2006GL027297
- Bevilacqua A., Bertagnini A., Pompilio M., Landi P., Del Carlo P., Di Roberto A., Aspinall W., Neri A. (2020) - Major explosions and paroxysms at Stromboli (Italy): a new historical catalog and temporal models of occurrence with uncertainty quantification. *Scientific Reports*, 10, 17357.
Doi: 10.1038/s41598-020-74301-6
- Bevilacqua A., Macedonio G., Neri A., Orsi G., Petrosino P. (2022) - Volcanic Hazard Assessment at the Campi Flegrei Caldera, Italy. *Active Volcano World*.
Doi: 10.1007/978-3-642-37060-1_12
- Bigazzi G., Bonadonna F.P., Iaccarino S. (1979) - Geochronological hypothesis on Plio-Pleistocene boundary in Latium region. *Bullettino Società Geologica Italiana*, 92, 391-422.
- Boari E., Avanzinelli R., Melluso L., Giordano G., Mattei M., De Benedetti A., Morra V., Conticelli S. (2009a) - Isotope geochemistry (Sr-Nd-Pb) and petrogenesis of leucite-bearing volcanic rocks from "Colli Albani" volcano, Roman Magmatic Province, Central Italy: inferences on volcano evolution and magma genesis. *Bulletin of Volcanology*, 71, 977-1005.
- Boari E., Tommasini S., Laurenzi M.A., Conticelli S. (2009b) - Transition from ultrapotassic Karmalugitic to Sub-alkaline Magmas: Sr, Nd, and Pb isotope, trace element and $^{40}\text{Ar}/^{39}\text{Ar}$ age data from the middle Latin Valley Volcanic Field, Roman Magmatic Province, Central Italy. *Journal of Petrology*, 50, 1327-1357.
- Bohnenstiehl R.W., Spera F.J., Fowler S.J., Belkin H.E., De Vivo B., Poland G. (2006) - Petrogenesis of the Campanian Ignimbrite: implications for crystal-melt separation and open-system processes from major and trace elements and Th isotopic data. In *Volcanism in the Campania Plain: Vesuvius, Campi Flegrei and Ignimbrites*, 9, 249-268.
- Bonini M., Sani F. (2002) - Extension and compression in the Northern Apennines (Italy) hinterland: Evidence from the late Miocene-Pliocene Siena-Radicofani Basin and relations with basement structures. *Tectonics*, 21, 1-19.
- Bonomo R., Ricci V. (2010) - Application of unconformity-bounded stratigraphic (UBS) units to the geological survey of the volcanic island Ustica (Italy). *Stratigraphy and Geology of Volcanic Areas*, 51.
- Borghetti G., Sbrana A., Sollevanti F. (1981) - Vulcanotettonica dell'area dei Monti Cimini e rapporti cronologici tra vulcanismo cimino e vicino. *Rendiconti della Società Geologica Italiana*, 4, 253-254.
- Borsi S., Ferrara G., Tongiorgi E. (1967) - Determinazione con il metodo K/Ar delle età delle rocce magmatiche della Toscana. *Bullettino della Società Geologica Italiana*, 86, 403-410.
- Borzi A.M., Giuffrida M., Zuccarello F., Palano M., Viccaro M. (2020) - The Christmas 2018 eruption at Mt. Etna: enlightening how the volcano factory works through a multi-parametric inspection. *Geochemistry Geophysics Geosystems*, 21, 10, e2020GC009226.
- Branca S., Abate T. (2019) - Current knowledge of Etna's flank eruptions (Italy) occurring over the past 2500 years. From the iconographies of the XVII century to modern geological cartography. *Journal of Volcanology and Geothermal Research*, 385, 159-178.
- Branca S., Del Carlo P. (2004) - Eruptions of Mt. Etna during the past 3,200 years: a revised compilation integrating the historical and stratigraphic records. *Geophysical Monograph Series* 143, 1-27.
- Branca S., Del Carlo P. (2005) - Types of eruptions of Etna Volcano AD 1670-2003: Implications for short-term eruptive behaviour. *Bulletin of Volcanology*, 67, 732-742.
- Branca S., Coltelli M., Groppelli G. (2004) - Geological evolution of Etna volcano. In: Bonaccorso A., Calvari S., Coltelli M., Del Negro C., Falsaperla S. (Eds.). *Mt. Etna Volcano Laboratory*. AGU (Geophysical monograph series), 143, 49-63.
- Branca S., Coltelli M., De Beni E., Wijbrans J. (2008) - Geological evolution of Mount Etna volcano (Italy) from earliest products until the first central volcan-

- ism (between 500 and 100 ka ago) inferred from geochronological and stratigraphic data. International Journal of Earth Sciences, 97, 135-152.
- Branca S., Cobelli M., Groppelli G. (2011a) - Geological evolution of a complex basaltic volcano: Mount Etna, Italy. Italian Journal of Geosciences, 130 (3), 306-317.
- Branca S., Cobelli M., Groppelli G., Lentini F. (2011b) - Geological map of Etna volcano, 1:50,000 scale. Italian Journal of Geosciences, 130 (3), 265-291.
- Broccolini D. (1999) - Cronologia ed evoluzione del vulcanismo nell'area del Somma-Vesuvio. Ph.D. thesis, Pisa, 1-168.
- Broccolini D., La Volpe L., Laurenzi M.A., Principe C. (1994) - Storia evolutiva del Mt. Vulture. Plinius 12, 22-25.
- Broccolini D., Di Battistini G., Laurenzi M.A., Vernia L., Bargossi G.M. (2000) - New $^{40}\text{Ar}/^{39}\text{Ar}$ datings on the southeastern sector of the Vulcian volcanic district (central Italy). Bollettino della Società Geologica Italiana, 119, 113-120.
- Brown R. J., Orsi G., De Vita S. (2008) - New insights into late Pleistocene explosive volcanic activity and caldera formation on Ischia (southern Italy). Bulletin of Volcanology, 70, 583-603.
- Brown R.J., Civetta L., Arzeno I., D'Antonio M., Moretti R., Orsi G., Tomlinson E.L., Albert P.G., Menzies M.A. (2014) - Geochemical and isotopic insights into the assembly, evolution and disruption of a magmatic plumbing system before and after a cataclysmic caldera collapse at Ischia volcano (Italy). Contribution to Mineralogy and Petrology, 168, 1035.
- Bruno P.P.G., de Alteris G., Florio G. (2002) - The western undersea section of the Ischia volcanic complex (Italy, Tyrrhenian Sea) inferred by marine geophysical data. Geophysical Research Letters, 29(9), 1343.
- Buettner A., Principe C., Villa IM., Broccolini D. (2006) - Geocronologia $^{39}\text{Ar}/^{40}\text{Ar}$ del Monte Vulture. La Geologia del Monte Vulture (a cura di Claudia Principe) In C. Principe ed. La geologia del Monte Vulture, 73-86.
- Calanchi N., Tranne C.A., Lucchini F., Rossi P.L., Villa I.M. (1999) - Explanatory notes to the geological map (1:10 000) of Panarea and Basiluzzo Islands, Aeolian Arc, Italy. Acta Vulcanologica, 11, 223-243.
- Caliro S., Caracausi A., Chiodini G., Ditta M., Italiano F., Longo M., Minopoli C., Nuccio P.M., Peonia A., Rizzo A. (2004) - Evidence of a new magmatic input to the quiescent volcanic edifice of Panarea, Aeolian Islands, Italy. Geophysical Research Letters, 31, L07819.
- Cadeux A., Pinti D. L., Aznar C., Chiesa S., Gillot P. (2005) - New chronological and geochemical constraints on the genesis and geological evolution of Ponza and Palmarola volcanic islands (Tyrrhenian Sea, Italy). Lithos, 81, 121-151.
- Calanchi N., Tranne C.A., Lucchini F., Rossi P.L., Villa I.M. (1999) - Explanatory notes to the geological map (1:10 000) of Panarea and Basiluzzo Islands, Aeolian Arc, Italy. Acta Vulcanologica, 11, 223-243.
- Caliro S., Caracausi A., Chiodini G., Ditta M., Italiano F., Longo M., Minopoli C., Nuccio P.M., Peonia A., Rizzo A. (2004) - Evidence of a new magmatic input to the quiescent volcanic edifice of Panarea, Aeolian Islands, Italy. Geophysical Research Letters, 31, L07819.
- Capaccioni B., Tassi F., Vaselli O., Tedesco D., Poreda R. (2007) - Submarine gas burst at Panarea Island southern Italy, on November 3, 2002: A magmatic versus hydrothermal episode. Journal of Geophysical Research, 112, B05201.
- Caporizzo C., Di Luccio D., Tursi M.F., Pappone G. (2022) - Coastal vs volcanic processes: Procida Island as a case of complex morpho-evolutive response. Marine Geology, 448, 106814.
- Carapezza M.L., Barberi F., Tarchini L., Ranaldi M., Ricci T. (2010) - Volcanic hazards of the Colli Albani. The Colli Albani Volcano, Special Volume Geological Society London, 3, 1-279.
- Carbone S., Lentini F. (1981) - Caratteri deposizionali delle vulcaniti del Miocene superiore negli Iblei (Sicilia sud-orientale). Geologica Romana, 20, 79-101.
- Cardillo G.L., Consorti L., Palladino D.M., Carminati E., Carlini M., Doglioni C. (2020) - Tectonically controlled carbonate-seated maar-diatreme volcanoes: the case of the Volsci Volcanic Field, central Italy. Journal of Geodynamics, 139, 101763.
- Casalbore D., Romagnoli C., Bosman A., De Astis G., Lucchi F., Tranne C.A., Chiocci F.L. (2018) - Multi-stage formation of La Fossa Caldera (Vulcano Island, Italy) from an integrated subaerial and submarine analysis. Marine Geophysical Research, DOI: 10.1007/s11001-018-9358-3
- Casalini M., Avanzinelli R., Heumann A., de Vita S., Saneivero F., Conticelli S., Tommasini S. (2017) - Geochemical and radiogenic isotope probes of Ischia volcano, Southern Italy: Constraints on magma chamber dynamics and residence time. American Mineralogist, 102, 262-274.
- Catalano S., Tortorici L., Viccaro M. (2014) - Regional tectonic control on large size explosive eruptions: insights into the Green Tuff ignimbrite unit of Pantelleria. Journal of Geodynamics, 73, 23-33.
- Cavazza W., Roure F., Ziegler P.A. (2004) - The Mediterranean area and the surrounding regions: active processes, remnants of former Tethyan oceans and related thrust belts. In The Transmed Atlas, The Mediterranean Region from Crust to Mantle, 1-29.

- Springer, Berlin, Heidelberg.
- Centamore E., Dramis F., Di Manna P., Fumanti F., Milli S., Rossi D., Palombo M.R., Palladino D.M., Trigila R., Zanon V., Chiocchini M., Didaskalou P., Potetti M., Niaic S. (2010) - Note illustrativa del Foglio 402 Ceccano. Carta Geologica d'Italia 1:50.000. Servizio Geologico d'Italia, Rome.
- Chiarabba C., Amato A., Delaney P.T. (1997) - Crustal structure, evolution, and volcanic unrest of the Alban Hills, Central Italy. *Bulletin of Volcanology*, 59, 161-170.
- Chiesa S., Floris B., Gillot P.Y., Prosperi L., Vezzoli L. (1995) - Il Vulcano di Roccamonfina. In: ENEA (Ed.), Lazio Meridionale. ENEA, 128-150.
- Ciaranfi N., Guida M., Iaccarino G., Pescatore T., Pieri P., Rapicardi L., Ricchetti G., Sgrocco I., Torre M., Tortorici L., Turco E., Scarpa R., Cuscho M., Guerra I., Iannaccone G., Panza G.F., Scandone P. (1983) - Elementi sismotettonici dell'Appennino meridionale. *Bollettino della Società Geologica Italiana*, 102, 201-222.
- Cimarelli C., De Rita D. (2006a) - Relatively rapid emplacement of dome-forming magma inferred from strain analyses: the case of the acid Latian dome complexes (Central Italy). *Journal of Volcanology and Geothermal Research*, 158, 108-116.
- Cimarelli C., De Rita D. (2006b) - Structural evolution of the Pleistocene Cimini trachytic volcanic complex (Central Italy). *Bulletin of Volcanology*, 68, 538-548.
- Cioni R., Sbrana A., Bertagnini A., Bucnascorta G., Landi P., Rossi U., Salvati L. (1987) - Tephrostratigraphic correlations in the Vulsini, Vico and Sabatini volcanic successions. *Periodico di Mineralogia*, 56, 137-155.
- Cioni R., Laurenzi M.A., Sbrana A., Villa L.M. (1989) - Geochronology and stratigraphy of basal pyroclastites of the Vulsini Volcanic District. *Plinius European Journal of Mineralogy*, 1, 46-47.
- Cioni R., Laurenzi M.A., Sbrana A., Villa L.M. (1993) - $^{40}\text{Ar}/^{39}\text{Ar}$ chronostratigraphy of the initial activity in the Sabatini volcanic complex (Italy). *Bollettino della Società Geologica Italiana*, 112, 251-263.
- Cioni R., Santacroce R., Sbrana A. (1999) - Pyroclastic deposits as a guide for reconstructing the multi-stage evolution of the Somma-Vesuvius caldera. *Bulletin of Volcanology*, 61, 207-222. Doi: 10.1007/s004450050272
- Cioni R., Longo A., Macedonio G., Santacroce R., Sbrana A., Sulpizio R., Andronico D. (2003) - Assessing pyroclastic fall hazard through field data and numerical simulations: example from Vesuvius. *Journal of Geophysical Research*, 108, 2063. Doi: 10.1029/2001JB000642
- Cioni R., Bertagnini A., Santacroce R., Andronico D. (2008) - Explosive activity and eruption scenarios at Somma-Vesuvius (Italy): Towards a new classification scheme. *Journal of Volcanology and Geothermal Research*, 178, 331-348. Doi: 10.1016/j.volgeores.2008.04.024
- Civetta L., Orsi G., Scandone P., Pepe R. (1978) - Eastwards migration of the Tuscan anatetic magmatism due to anticlockwise rotation of the Apennines. *Nature*, 276, 604-606.
- Civetta L., Cornette Y., Crisci G., Gillot P.Y., Orsi G., Requejos C.S. (1984) - Geology, geochronology and chemical evolution of the Island of Pantelleria. *Geological Magazine*, 121, 541-668.
- Civetta L., Cornette Y., Gillot P.Y., Orsi G. (1988) - The eruptive history of Pantelleria (Sicily Channel) in the last 50 ka. *Bulletin of Volcanology*, 50, 47-57.
- Civetta L., Carluccio E., Innocenti F., Sbrana A., Taddeucci G. (1991a) - Magma chamber evolution under Phlegraean Fields during the last 10 ka: trace element and isotope data. *European Journal of Mineralogy*, 3, 415-428.
- Civetta L., Gallo G., Orsi G. (1991b) - Sr- and Nd-isotope and trace-element constraints on the chemical evolution of the magmatic system of Ischia (Italy) in the last 55 ka. *Journal of Volcanology and Geothermal Research*, 46, 213-230. Doi: 10.1016/0377-0273(91)90084-D
- Civetta L., Orsi G., Pappalardo L., Fisher R.V., Heiken G.H., Ort M. (1997) - Geochemical zoning, mixing, eruptive dynamics and depositional processes - the Campanian Ignimbrite, Campi Flegrei, Italy. *Journal of Volcanology and Geothermal Research*, 75, 183-219.
- Civile D., Lodolo E., Tortorici L., Lanzafame G., Brancolini G. (2006) - Relationships between magmatism and tectonics in a continental rift: The Pantelleria Island region (Sicily Channel, Italy). *Marine Geology*, 251, 32-46.
- Ciocchiali R., Condoini M., Guenot N., Tanguy J.C. (2004) - Magma changes at Mount Etna: the 2001 and 2002-2003 eruptions. *Earth and Planetary Science Letters*, 226, 397-414.
- Cole P.D., Scarpati C. (1993) - A facies interpretation of the eruption and emplacement mechanisms of the upper part of the Neapolitan Yellow Tuff, Campi Flegrei, southern Italy. *Bulletin of Volcanology*, 55, 311-326.
- Cole P.D., Guest J.E., Duncan A.M. (1993) - The emplacement of intermediate volume ignimbrites: a case study from Roccamonfina Volcano, Southern Italy. *Bulletin of Volcanology*, 55, 467-480.
- Cole J.W., Milner D.M., Spinks K.D. (2005) - Calderas and caldera structures: A review. *Earth-Science Reviews*, 69, 1-26.
- Cotelli M., Del Carlo P., Vezzoli L. (1998) - The discovery of a Plinian basaltic eruption of Roman age at Mt. Etna volcano (Italy). *Geology*, 26, 1095-1098.
- Cotelli M., Del Carlo P., Vezzoli L. (2000) - Stratigraphic constraints for explosive activity in the last 100 ka at Etna volcano, Italy. *International Journal of Earth Sciences*, 89, 665-677.
- Cotelli M., Cavallaro D., D'Anna G., D'Alessandro A., Graessa F., Mangano G., Patanè D., Greco S. (2016) - Exploring the submarine Graham Bank in the Sicily Channel. *Annals of Geophysics*, 59, 2. Doi: 10.4401/ag-6929
- Constantinescu R., Robertson R., Lindsay J.M., Tonini R., Sandri L., Rouwet D., Smith P., Stewart R. (2016) - Application of the probabilistic model BET_UNREST during a volcanic unrest simulation exercise in Dominica, Lesser Antilles. *Geochemistry Geophysics and Geosystems*, 17, 4438-4456. Doi: 10.1002/2016GC006485
- Conte A.M., Dotti D. (2002) - Petrological and geochemi-

- cal characteristics of Plio-Pleistocene volcanics from Ponza Island (Tyrrenian Sea, Italy). *Mineralogy and Petrology*, 74, 75-94.
- Conte A.M., Dotti D., Gaeta M., Misiti V., Mollo S., Perinelli C. (2009) - Experimental constraints on evolution of leucite-basanite magma at 1 and 10-4 GPa: Implications for parental compositions of Roman high-potassium magmas. *European Journal of Mineralogy*, 21, 763-782.
- Conte A.M., Martorelli E., Calarco M., Sposato A., Perinelli C., Coltellini M., Chiocci F.L. (2014) - The 1891 submarine eruption offshore Pantelleria Island (Sicily Channel, Italy): Identification of the vent and characterization of products and eruptive style. *Geochemistry, Geophysics, Geosystems*, 15, 6, 25555-2574
DOI: 10.1002/2014GC005238
- Conte A.M., Perinelli C., Bianchini G., Natali C., Martorelli E., Chiocci F.L. (2016) - New insights on the petrology of submarine volcanoes from the western Pontine Archipelago (Tyrrenian Sea, Italy). *Journal of Volcanology and Geothermal Research*, 327, 223-239.
- Conte A.M., Perinelli C., Bosman A., Castorina F., Conti A., Cuffaro M., Di Vincenzo G., Martorelli E., Bigi S. (2020) - Tectonics, dynamics and Plio-Pleistocene magmatism in central Tyrrenian Sea: insights from the transitional submarine basalts of the Ventotene volcanic ridge (Pontine Islands, Italy). *Geochemistry, Geophysics, Geosystems*, 21 (12).
DOI: 10.1029/2020GC009346
- Conticelli S. (1998) - Effects of Crustal Contamination on Ultrapotassic Magmas with Lamproitic Affinity: Mineralogical, Geochemical and Isotope data from the Torre Alfina Lavas and Xenoliths, Central Italy. *Chemical Geology*, 149, 51-81.
- Conticelli S., Peccerillo A. (1990) - Petrological significance of High-Pressure ultramafic xenoliths from ultrapotassic rocks of Central Italy. *Lithos*, 24, 305-322.
DOI: 10.1016/0024-4937(89)90050-9
- Conticelli S., Peccerillo A. (1992) - Potassic and ultrapotassic volcanism from central Italy: compositional characteristics, petrogenesis and inferences on the evolution of the mantle source. *Lithos*, 28, 221-240.
DOI: 10.1016/0024-4937(92)90008-M
- Conticelli S., Francalanci L., Manetti P., Peccerillo A. (1987) - Evolution of Latara Volcano: evidence from major and trace element chemistry of pyroclastic rocks. *Periodico di Mineralogia*, 56, 175-199.
- Conticelli S., Francalanci L., Santo A.P. (1991) - Petrology of the final stage Latara lavas: Mineralogical, Geochemical and Sr-isotopic data and their bearing on the genesis of some potassic magmas in Central Italy. *Journal Volcanology and Geothermal Research*, 46, 187-212.
DOI: 10.1016/0377-0273(91)90063-C
- Conticelli S., Manetti P., Menichetti S. (1992) - Petrology, Chemistry, Mineralogy and Sr-isotopic features of Pliocene Orendites from South Tuscany: implications on their genesis and evolutions. *European Journal of Mineralogy*, 4, 1359-1375.
- Conticelli S., Francalanci L., Manetti P., Cioni R., Sbrana A. (1997) - Petrology and Geochemistry of the ultrapotassic rocks from the Sabatini Volcanic District, Central Italy: the role of evolutionary processes in the genesis of variably enriched alkaline magmas. *Journal of Volcanology and Geothermal Research*, 75, 107-136.
- Conticelli S., Bortolotti V., Principi G., Laurenzi M.A., Vagelli G., D'Antonio M. (2001) - Petrology, mineralogy and geochemistry of a mafic dyke from Monte Castello, Elba Island, Italy. *Ophioliti* 26, 249-262.
- Conticelli S., D'Antonio M., Pirarelli L., Civetta L. (2002) - Source contamination and mantle heterogeneity in the genesis of Italian potassic and ultrapotassic volcanic rocks: Sr-Nd-Pb Isotope data from Roman Province and Southern Tuscany. *Mineralogy and Petrology*, 74, 189-222.
DOI: 10.1007/s007100200004
- Conticelli S., Melluso L., Pierini G., Avanzinelli R., Boari E. (2004) - Petrologic, geochemical, and isotopic characteristics of potassic and ultrapotassic magmatism in Central-Southern Italy: inferences on its genesis and on the nature of mantle sources. *Periodico di Mineralogia*, 73, 153-164.
- Conticelli S., Guarneri L., Farinelli A., Mattioli M., Avanzinelli R., Bianchini G., Boari E., Tommasini S., Tiepolo M., Prelević D., Venturelli G. (2009a) - Trace elements and Sr-Nd-Pb isotopes of K-rich, shoshonitic, and calc-alkaline magmatism of the Western Mediterranean Region: genesis of ultrapotassio to calc-alkaline magmatic associations in a post-collisional geodynamic setting. *Lithos*, 107, 68-92.
- Conticelli S., Marchionni S., Rosa D., Giordano G., Boari E., Avanzinelli R. (2009b) - Shoshonite and sub-alkaline magmas from an ultrapotassic volcano: Sr-Nd-Pb isotope data on the Roccamonfina volcanic rocks, Roman Magmatic Province, Southern Italy. *Contribution to Mineralogy and Petrology*, 157, 41-63.
- Conticelli S., Lauretti M.A., Giordano G., Mattioli M., Avanzinelli R., Melluso L., Tommasini S., Boari E., Cifelli F., Pierini G. (2010) - Leucite-bearing (kamafugitic/leucitic) and-free (lamproitic) ultrapotassio rocks and associated shoshonites from Italy: constraints on petrogenesis and geodynamics. *Journal of Virtual Explorer*, 36, 1-95.
- Conticelli S., Avanzinelli R., Marchionni S., Tommasini S., Melluso L. (2011) - Sr-Nd-Pb isotopes from the Radicofani Volcano, Central Italy: constraints on heterogeneities in a veined mantle responsible for the shift from ultrapotassic shoshonite to basaltic andesite magmas in a post-collisional setting. *Mineralogy and Petrology*, 103, 123-148.
- Conticelli S., Avanzinelli R., Poli G., Braschi E., Giordano G. (2013) - Shift from lamproite-like to leucitic rocks: Sr-Nd-Pb isotope data from the Monte Cimino volcanic complex vs. the Vico stratovolcano, Central Italy. *Chemical Geology*, 353, 246-266.
- Conticelli S., Avanzinelli R., Ammannati E., Casalini M. (2015a) - The role of carbon from recycled sediments in the origin of ultrapotassic igneous rocks in the Central Mediterranean. *Lithos*, 232, 174-196.
- Conticelli S., Boari E., Burlamacchi L., Cifelli F., Moscardi F., Laurenzi M.A., Ferrari Pedraglio L., Francalanci L., Bervenuti M.G., Braschi E., Manetti P. (2015b) -

- Geochemistry and Sr-Nd-Pb isotopes of Monte Amiata Volcano, Central Italy: evidence for magma mixing between high-K calc-alkaline and leucitic mantle-derived magmas. *Italian Journal of Geosciences*, 134, 266-290.
- Corsaro R.A., Cristofolini R. (1997) - Geology, geochemistry and mineral chemistry of tholeiitic to transitional Etnaean magmas. *Acta Vulcanologica*, 9, 55-66.
- Corsaro R.A., Cristofolini R. (2000) - Subaqueous volcanism in the Etnaean area: evidence for hydromagmatic activity and regional uplift inferred from the Castle Rock of Acicastello. *Journal of Volcanology and Geothermal Research*, 95, 209-225.
- Corsaro R.A., Cristofolini R., Patanè L. (1996) - The 1669 eruption at Mount Etna: chronology, petrology and geochemistry, with inferences on the magma sources and ascent mechanisms. *Bulletin of Volcanology*, 58, 348-358.
- Corsaro R.A., Andronico D., Behncke B., Branca S., Caltabiano T., Ciancittà F., Cristaldi A., De Beni E., La Spina A., Lodato L., Miraglia L., Neri M., Salemo G., Scollo S., Spata G. (2017) - Monitoring the December 2015 summit eruptions of Mt. Etna (Italy): Implications on eruptive dynamics. *Journal of Volcanology and Geothermal Research*, 341, 53-69.
- Cortini M., Scandone R. (1982) - Feeding system of Vesuvius between 1754 and 1944. *Journal of Volcanology and Geothermal Research*, 12, 393-400.
- Costa A., Dell'Erba F., Di Vito M.A., Isala R., Macedonio G., Orsi G., Pfeiffer T. (2009) - Tephra fallout hazard assessment at the Campi Flegrei caldera (Italy). *Bulletin of Volcanology*, 71, 259-273. doi: 10.1007/s00445-008-0220-3
- Costa G., Meru L., Prestifilippo M., Scollo S., Viccaro M. (2023) - Modeling the trajectories of ballistics in the summit area of Mt. Etna (Italy) during the 2020-2022 sequence of lava fountains. *Geosciences*, 13 (5), 145.
- Crisci G., de Fino M., La Volpe L., Rapisardi L. (1983) - Pleistocene ignimbrites of Monte Vulture (Basilicata, Southern Italy). *Neues Jahrbuch für Geologie und Paläontologie*, 12, 731-746.
- Crisci G.M., De Francesco A.M., Mazzulli R., Poli G., Stanzione O. (1989) - Geochemistry of recent volcanics of Ischia Island, Italy: evidences of fractional crystallization and magma mixing. *Chemical Geology*, 78, 15-33.
- Crocitti M., Sulpizio R., Insinga D.D., De Rosa R., Donato P., Iorio M., Zanchetta G., Barca D., Lubritto C. (2019) - On ash dispersal from moderately explosive volcanic eruptions: Examples from Holocene and Late Pleistocene eruptions of Italian volcanoes. *Journal of Volcanology and Geothermal Research* 385, 198-221.
- Cristofolini (1986) - Le manifestazioni eruttive basiche del Trias superiore nel sottosuolo di Ragusa. *Periodico di Mineralogia*, 35, 1-38.
- Cristofolini R., Ghisetti F., Scarpa R., Vezzani L. (1985) - Character of the stress field in the Calabrian arc and southern Apennines (Italy) as deduced by geological, seismological and volcanological information. *Tectonophysics*, 117, 39-58.
- Cross J.K., Tomlinson E.L., Giordano G., Smith V.C., De Benedetti A.A., Roerge J., Manning C.J., Wulf S., Menzies M.A. (2014) - High level triggers for explosive mafic volcanism: Albano Maar, Italy. *Lithos*, 190, 137-153.
- Cuffaro M., Martorelli E., Bosman A., Conti A., Bigi S., Muccini F., Cochi L., Ugli M., Bortoluzzi G., Scrocca D., Canese S., Chiocci F.L., Conte A.M., Doglioni D., Perinelli C. (2016) - The Ventotene Volcanic Ridge: Newly explored complex in the central Tyrrhenian Sea (Italy). *Bulletin of Volcanology*, 78, 86.
- D'Alessandro A., Mangano G., D'Anna G. (2012) - Evidence of persistent seismo-volcanic activity at Marsili seamount. *Annals of Geophysics*, 55, 213-214.
- D'Ambrosio E., Giaccio B., Lombardi L., Marra F., Rofo M.F., Sposato A. (2010) - L'attività recente del centro eruttivo di Albano tra scienza e mito: un'analisi critica del rapporto tra il vulcano laziale e la storia dell'area albana - Lazio e Sabina, *Sesto Incontro di Studi sul Lazio e Sabina*, Roma, 125-136.
- D'Antonio M., Civetta L., Orsi G., Pappalardo L., Piochi M., Carandente A., De Vita S., Di Vito M.A., Isala R., Southon J. (1999) - The present state of the magmatic system of the Campi Flegrei caldera based on the reconstruction of its behaviour in the past 12 ka. *Journal of Volcanology and Geothermal Research*, 91 (2-4), 247-268.
- D'Antonio M., Tonarini S., Arienzo I., Civetta L., Di Renzo V. (2007) - Components and processes in the magma genesis of the Phlegraean Volcanic District, southern Italy. In: Beccaluva L., Bianchini G., Wilson M. (Eds.) *Cenozoic Volcanism in the Mediterranean Area*. Geological Society of America Special Paper, 418, 203-220. doi: 10.1130/2007.2418(10)
- D'Argenio B., Pescatore T., Scandone P. (1973) - Schema geologico dell'Appennino Meridionale. Atti del Convegno "Moderne vedute sulla geologia dell'Appennino". Accademia Nazionale dei Lincei, 183, 49-72.
- D'Antonio M., Arienzo I., Brown R.J., Petrosino P., Petullo C., Giaccio B. (2021) - Petrography and mineral chemistry of Monte Epomeo Green Tuff, Ischia Island, south Italy: Constraints for identification of the Y-7 tephrostratigraphic marker in distal sequences of the central Mediterranean. *Minerals* 11, 9.
- De Astis G., Ventura G., Vilardo G. (2003) - Geodynamic significance of the Aeolian volcanism (Southern Tyrrhenian Sea, Italy) in light of structural, seismological and geochemical data. *Tectonics*, 22, 1040-1057.
- De Astis G., Pappalardo L., Piochi M. (2004) - Procida volcanic history: new insights into the evolution of the Phlegraean Volcanic District (Campania region, Italy). *Bulletin of Volcanology*, 66, 622-641.
- De Astis G., Lucchi F., Dellino P., La Volpe L., Tranne C.A., Frezzotti M.L., Peccerillo, A. (2013) - Geology, volcanic history and petrology of Vulcano (central Aeolian archipelago). *Geological Society, London, Memoirs*, 37, 281-348.
- De Benedetti A.A., Funiciello R., Giordano G., Diana G., Caprilli E., Paterne M. (2008) - Volcanology, history and myths of the Lake Albano maar (Colli Albani volcano, Italy). *Journal of Volcanology and Geo-*

- thermal Research, 176(3), 387-406.
- De Beni E., Branca S., Coltell M., Groppelli G., Wijbrans J. (2011) - $^{39}\text{Ar}/^{40}\text{Ar}$ isotopic dating of Etna volcanic succession. *Italian Journal of Geosciences*, 130, 3, 292-305.
- De Fino M., La Volpe L., Piccarreta G. (1982) - Magma evolution at Mt. Vulture (Southern Italy). *Bulletin of Volcanology*, 45, 115-126.
- De Fiore O. (1922) - Vulcano (Isole Eolie). *Rivista Vulcanologica I*. Friedlaender 3, 1-383.
- Deino A.L., Orsi G., Piochi M., de Vita S. (2004) - The age of the Neapolitan Yellow Tuff caldera-forming eruption (Campi Flegrei caldera - Italy) assessed by $40\text{Ar}/39\text{Ar}$ dating method. *Journal of Volcanology and Geothermal Research*, 133, 157-170.
- Del Carlo P., Branca S., D'Orlando C. (2017) - New findings of Etna pumice fall deposits in NE Sicily and implications for distal tephra correlations in the Mediterranean area. *Bulletin of Volcanology*, 75, 50.
- Del Negro C., Cappello A., Neri M., Bliotta G., Herault A., Ganci G. (2013) - Lava flow hazards at Mount Etna: Constraints imposed by eruptive history and numerical simulations. *Scientific Reports*, 3, 3493.
- de Rita D., Giordano G. (1996) - Volcanological and structural evolution of Roccamontfina volcano (Italy): origin of the summit caldera. *Geological Society of London Special Publications*, 110(1), 209-224.
- de Rita D., Funiciello R., Pantosti D., Salvini F., Sposato A., Velona M. (1988) - Geological and structural characteristics of the Pontine Islands (Italy) and implications with the evolution of the Tyrrhenian margin. *Memorie della Società Geologica Italiana*, 36, 55-65.
- de Rita D., Funiciello R., Parotto M. (1988) - Geological Map of the Colli Albani Volcanic Complex. Progetto Finalizzato Geodinamica C.N.R., Rome, Italy.
- de Rita D., Funiciello R., Corda L., Sposato A., Rossi U. (1993) - Volcanic units. In: Di Filippo M. (Ed.), Sabatini Volcanic Complex. Quad. Ric. Sci., 114, 33-79. Progetto Finalizzato Geodinamica C.N.R., Roma.
- de Rita D., Faccenna C., Funiciello R., Rosa C. (1995a) - Stratigraphy and Volcano-Tectonics. In: Trigila, R. (Ed.), The Volcano of the Alban Hills. Università degli Studi di Roma "La Sapienza", Rome, Italy, 33-71.
- de Rita D., Giordano G., Rosa C., Sheridan M.F. (1995) - Volcanic risk at the Alban Hills volcano and numerical simulations. *The Volcano of the Alban Hills*. Tipografia SGS, Roma, 267-283.
- de Rita D., Di Filippo M., Rosa C. (1996) - Structural evolution of the Bracciano volcano-tectonic depression, Sabatini Volcanic District, Italy. In: McGuire W.J., Jones A.P., Neuberg J. (Eds.), Volcano instability on the Earth and other planets. *Geological Society of London Special Publications*, 225-236.
- de Rita D., Giordano G., Cecili A. (2001) - A model for submarine rhyolite dome growth: Ponza Island (central Italy). *Journal of Volcanology and Geothermal Research*, 107, 221-239.
- De Rosa R., Mazzuoli R., Scribano V., Trua T. (1991) - Nuovi dati petrologici sulle vulcaniti dei Monti Iblei (Sicilia sud-orientale): implicazioni genetiche e geotettoniche. *Mineralogica et Petrografia Acta*, 34, 133-151.
- de Vita S., Laurenzi M.A., Orsi G., Voltaggio M. (1998) - Application of $^{40}\text{Ar}/^{39}\text{Ar}$ and ^{230}Th dating methods to the chronostratigraphy of quaternary basaltic volcanic areas: the Ustica Island case history. *Quaternary International*, 4 (48), 117-127.
- de Vita S., Sansivero F., Orsi G., Marotta E. (2006) - Cyclical slope instability and volcanism related to volcano-tectonism in resurgent calderas: the Ischia Island (Italy) case study. *Engineering Geology*, 86, 148-165.
- de Vita S., Sansivero F., Orsi G., Marotta E., Piochi M. (2010) - Volcanological and structural evolution of the Ischia resurgent caldera (Italy) over the past 10 ky. In: Groppelli G., Vierck-Goelet L. (Eds.), Stratigraphy and geology of volcanic areas. *Geological Society of America Special Papers*, 464, 193-241. doi: 10.1130/2010.2464
- de Vita S., Di Vito M., Gilanella C., Sansivero F. (2013) - The impact of the Ischia Porto tephra eruption (Italy) on the Greek colony of Pithekoussai. *Quaternary International*, 303, 142-52.
- De Vivo B., Rolandi G., Gans P.B., Calvert A., Bohrson W.A., Spera F.J., Belkin H.E. (2001) - New constraints on the pyroclastic eruptive history of the Campanian volcanic Plain (Italy). *Mineralogy and Petrology*, 73, 47-65.
- Di Girolamo P., Ghiara M.R., Lirer L., Munno R., Rolandi G., Stanzone D. (1984) - Vulcanologia e petrologia dei Campi Flegrei. *Bullettino della Società Geologica Italiana*, 103, 349-413.
- Di Capua A., Vezzoli G., Groppelli G. (2016) - Climatic, tectonic and volcanic controls of sediment supply to an Oligocene foredeep basin: The Val d'Aveto Formation (Northern Italian Apennines). *Sedimentary Geology*, 332, 68-84.
- Di Renzo V., Civetta L., D'Antonio M., Tonarini S., Di Vito M.A., Orsi G. (2011) - The magmatic feeding system of the Campi Flegrei caldera: architecture and temporal evolution. *Chemical Geology*, 281, 227-241. doi: 10.1016/j.chemgeo.2010.12.010
- Di Rita F., Scotti G. (2019) - Pollen analysis and tephrochronology of a MIS 13 lacustrine succession from Eastern Sabatini Volcanic District (Pignano Flaminio, central Italy). *Quaternary Science Reviews*, 204, 78-93.
- Di Rocco T., Freda C., Gaeta M., Mollo S., Dalai L. (2012) - Magma chambers emplaced in carbonate substrate: Petrogenesis of skarn and cumulate rocks and implications for CO₂ degassing in volcanic areas. *Journal of Petrology*, 53, 2307-2332.
- Disperati L., Liotta D. (1998) - Estimating uplift of clay-filled extensional basins through the porosity depth curve: The case of the Radiocarbon Basin (Italy). *Annals of Tectonics*, 12, 162-176.
- Di Vito M.A., Isaia R., Orsi G., Southon J., de Vita S., D'Antonio M., Pappalardo L., Piochi M. (1999) - Volcanic and deformation history of the Campi Flegrei caldera in the past 12 ka. *Journal of Volcanology and Geothermal Research*, 91, 221-246.
- Di Vito M.A., Acocella V., Aiello G., Barra D., Battaglia

- M., Carandente A., Del Gaudio C., de Vita S., Ricciardi G.P., Ricco C., Scandone R., Terrasi F. (2016) - Magma transfer at Campi Flegrei caldera (Italy) before the 1538 AD eruption, *Scientific Reports*, 6, 32245, 1-9.
DOI: doi.org/10.1038/srep32245
- Dolfi D., Conte A.M. (2018) - VII-Caratteri mineralogici e petrochimici delle vulcaniti delle Isole Pontine Occidentali (Ponza, Zannone, Palmarola). In: Note illustrative della Carta geologica d'Italia alla scala 1:50.000. Foglio 413, Borgo Grappa Isole Poniane. ISPRA- Dipartimento per il Servizio Geologico d'Italia, 2018.
- D'Orazio M., Laurenzi M.A., Villa I.M. (1991) - $^{40}\text{Ar}/^{39}\text{Ar}$ dating of a shoshonitic lava flow of the Radiofani volcanic center (Southern Tuscany). *Acta Vulcanologica*, 1, 63-67.
- D'Orlando C., Poggianti E., Bertagnini A., Cioni R., Landi P., Polacci M., Rosi M. (2005) - Changes in eruptive style during the A.D. 1538 Monte Nuovo eruption (Phlegraean Fields, Italy): the role of syn-eruptive crystallization. *Bulletin of Volcanology*, 67, 601-621.
- D'Orazio M., Innocenti F., Tonarini S., Doglioni C. (2006) - Reply to the discussion of "Carbonatites in a subduction system: the Pleistocene Alvikites from Mt. Vulture (Southern Italy)" by D'Orazio M., Innocenti F., Tonarini S., Doglioni C. (*Lithos* 98, 313-334) by Stoppa F., Principe C., Giannandrea P., *Lithos*, 103, 557-561.
- Doronzio D.M., Di Vito M.A., Arienza I., Bini M., Calusi B., Cerninara M., Corradini S., de Vita S., Giaccio B., Gurioli L., Mannella G., Ricciardi G.P., Ruco I., Spario D., Todesco M., Trasatti E., Zanchetta G. (2022) - The 79 CE eruption of Vesuvius: A lesson from the past and the need of a multidisciplinary approach for developments in volcanology. *Earth-Science Reviews* 231, 104072.
- Downes H., Kostoula T., Jones A.P., Beard A.D., Thirlwall M.F., Bodnar J.-L. (2002) - Geochemistry and Sr-Nd isotopic compositions of mantle xenoliths from the Mount Vulture carbonatite-melilitite volcano, central southern Italy. Contribution to Mineralogy and Petrology, 144, 78-92.
- Evernden J.F., Curtis G.H. (1965) - The potassium-argon dating of late Cenozoic rocks in East Africa and Italy. *Current Anthropology*, 6, 343-385.
- Faccenna C., Becker T.W., Lucente F.P., Jolivet L., Rossetti F. (2001a) - History of subduction and back arc extension in the Central Mediterranean. *Geophysical Journal International*, 145(3), 809-820.
- Faccenna C., Funiciello F., Giardini D., Lucente P. (2001b) - Episodic back-arc extension during restricted mantle convection in the Central Mediterranean. *Earth and Planetary Science Letters*, 187, 105-118.
- Faccenna C., Piromallo C., Crespo-Blanc A., Jolivet L., Rossetti F. (2004) - Lateral slab deformation and the origin of the western Mediterranean arcs. *Tectonics*, 23, TC1012.
DOI:10.1029/2002TC001488, 2004
- Fazzini P., Gelmini R., Mantovani M.P., Pellegrini M. (1972) - Geologia dei Monti della Tolfa (Lazio Settentrionale, province di Viterbo e Roma). *Memorie Società Geologica Italiana*, 11, 65-144.
- Fedele L., Scarpati C., Lanphere M., Melluso L., Morra V., Perrotta A., Ricci G. (2008) - The Breccia Museo formation, Campi Flegrei, southern Italy: geochronology, chemostratigraphy and relationship with the Campanian Ignimbrite eruption. *Bulletin of Volcanology*, 70, 1189-1219.
- Fedele L., Morra V., Perrotta A., Scarpati C., Sbrana A., Putignano M. L., Orrù P., Schiattarella M., Aleo G., D'Arganio B., Conforti A. (2012) - Note Illustrative della Carta geologica dell'Isola di Procida alla scala 1: 10.000-Foglio 465-464.
- Fertlito C., Viocaro M., Nicotra E., Cristofolini R. (2010) - Relationship between the flank sliding of the South East Crater (Mt. Etna, Italy) and the paroxysmal event of November 16, 2006. *Bulletin of Volcanology*, 72, 1179-1190.
- Fertlito C., Viocaro M., Nicotra E., Cristofolini R. (2012) - Regimes of magma recharge and their control on the eruptive behaviour during the 2001-2006 period at Mt. Etna (Italy). *Bulletin of Volcanology*, 74, 533-543.
- Ferrari L., Conticelli S., Burlamacchi L., Manetti P. (1996) - Volcanological Evolution of the Monte Amiata Volcanic Center, Southern Tuscany, Central Italy: New Geological and Petrochemical data. *Acta Vulcanologica*, 8, 41-56.
- Finetti I., Morelli C. (1974) - Esplorazione sismica a rilievo del Golfo di Napoli e Pozzuoli. *Bollettino di Geofisica Teorica e Applicata*, 16, 175-222.
- Fisher R.V., Orsi G., Ort M., Heiken G. (1993) - Mobility of a large-volume pyroclastic flow: emplacement of the Campanian Ignimbrite, Italy. *Journal of Volcanology and Geothermal Research*, 56, 205-220.
- Folch A., Sulpizio R. (2010) - Evaluating long-range volcanic ash hazard using supercomputing facilities: application to Somma-Vesuvius (Italy), and consequences for civil aviation over the Central Mediterranean Area. *Bulletin of Volcanology*, 79, 1039-1059.
- Forni F., Lucchi F., Pacciarino A., Tranne C.A., Rossi P.L., Frezzotti M.L. (2013) - Stratigraphy and geological evolution of the Lipari volcanic complex (central Aeolian archipelago). *Geological Society, London, Memoirs*, 37, 213-279.
- Forni F., Petricca E., Bachmann O., Mollo S., De Astis G., Piochi M. (2018) - The role of magma mixing/mingling and cumulate melting in the Neapolitan Yellow Tuff caldera-forming eruption (Campi Flegrei, Southern Italy). *Contribution to Mineralogy and Petrology*, 173, 45.
- Fournieraux C., Metrich N., Bertagnini A., Rosi M. (2012) - Crystal fractionation, magma step ascent, and syn-eruptive mingling: the Averno 2 eruption (Phlegraean Fields, Italy). *Contribution to Mineralogy and Petrology*, 163, 6, 1121-1137.
- Fowler J., Spera F., Bohrson W., Belkin H.E., De Vivo B. (2007) - Phase equilibria constraints on the chemical and physical evolution of the Campanian Ignimbrite. *Journal of Petrology*, 48, 459-493.
- Francalanci L., Lucchi F., Keller J., De Astis G., Tranne C.A. (2013) - Eruptive, volcano-tectonic and magmatic history of the Stromboli volcano (north-eastern Aeolian archipelago). *Geological Society,*

- London, Memoirs, 37, 395-469.
- Freda C., Gaeta M., Palladino D.M., Trigila R. (1997) - The Villa Senni eruption (Alban Hills, Central Italy): the role of H₂O and CO₂ on the magma chamber evolution and on the eruptive scenario. *Journal of Volcanology and Geothermal Research*, 78, 103-120.
- Freda C., Gaeta M., Karner D.B., Marra F., Renne P.R., Taddeucci J., Scarlato P., Christensen J.N., Dallai L. (2006) - Eruptive history and petrologic evolution of the Albano multiple maar (Alban Hills, Central Italy). *Bulletin of Volcanology*, 68, 567-591.
- Freda C., Gaeta M., Misiti V., Mollo S., Dolfi D., Scarlato P. (2008) - Magma-carbonate interaction: an experimental study on ultrapotassic rocks from Alban Hills (Central Italy). *Lithos*, 101, 397-415.
- Freda C., Gaeta M., Giacco B., Marra F., Palladino D.M., Scarlato P., Sottili G. (2011) - CO₂-driven large mafic explosive eruptions: a case study from the Colli Albani (central Italy). *Bulletin of Volcanology*, 73, 241-256.
- Funiciello R., Giordano G., de Rita D. (2003) - The Albano maar lake (Colli Albani Volcano, Italy): recent volcanic activity and evidence of pre-Roman Age catastrophic lahar events. *Journal of Volcanology and Geothermal Research*, 123, 43-61.
- Gaeta M. (1998) - Petrogenetic implications of Basanidine in the Lionato Tuff (Colli Albani Volcanic District, Central Italy). *Mineralogical Magazine*, 62, 697-701.
- Gaeta M., Freda C. (2001) - Strontian fluoromagnesiohastingsite in Alban Hills lavas (Central Italy): constraints on crystallization conditions. *Mineralogical Magazine*, 65, 787-795.
- Gaeta M., Fabbrizio G., Cavarretta G. (2000) - F-philoxopites in the Alban Hills Volcanic District (Central Italy): indications regarding the role of volatiles in magmatic crystallization. *Journal of Volcanology and Geothermal Research*, 99, 179-193.
- Gaeta M., Freda C., Christensen J.N., Dallai L., Marra F., Karner D.B., Scarlato P. (2006) - Time-dependent geochemistry of clinopyroxene from the Alban Hills (Central Italy): Clues to the source and evolution of ultrapotassic magmas. *Lithos*, 86, 330-346.
- Gaeta M., Di Rocca T., Freda C. (2009) - Carbonate assimilation in open magmatic systems: The role of melt-bearing skarns and cumulate-forming processes. *Journal of Petrology*, 50, 361-385.
- Gaeta M., Freda C., Marra F., Arienzo I., Gozzi F., Jicha B., Di Rocca T. (2016) - Paleozoic metasomatism at the origin of Mediterranean ultrapotassic magmas: constraints from time-dependent geochemistry of Colli Albani volcanic products (Central Italy). *Lithos*, 244, 151-64.
- Gaeta M., Bonechi B., Marra F., Perinelli C. (2021) - Uncommon K-felditic magmas: The case study of Tufo del Palatino (Colli Albani Volcanic District, Italy). *Lithos*, 396-397 (106): 239.
- Galli P., Giacco B., Messina P., Peronace E., Amato V., Naso J., Nomade S., Pereira A., Piccetti S., Bellanova J., Billi A., Blamart D., Galderisi A., Giacoli A., Stabile T., Thil F. (2017) - Middle to Late Pleistocene activity of the northern Matese fault system (southern Apennines, Italy). *Tectonophysics*, 699, 61-81.
- Gasparini P., Adams J.A.S. (1967) - K-Ar dating of Italian Plio-Pleistocene volcanic rocks. *Earth and Planetary Science Letters*, 6, 225-230.
- Gasparini M., Rosenbaum G., Wijbrans J., Manetti P. (2009) - The transition from subduction arc to slab tearing: Evidence from Capraia Island, northern Tyrrhenian Sea. *Journal of Geodynamics*, 47, 30-38.
- Ghinassi M., Lazzarotto A. (2005) - Different source areas feeding the Early Pliocene marine Radiofani Basin (Siena): geological and paleogeographical implications. *Boletino della Società Geologica Italiana*, 3, 15-28.
- Giacco B., Sposato A., Gaeta M., Marra F., Palladino D.M., Taddeucci J., Barbieri M., Messina P., Rollo M.F. (2007) - Mid-distal occurrences of the Albano Maar pyroclastic deposits and their relevance for reassessing the eruptive scenarios of the most recent activity at the Colli Albani Volcanic District, Central Italy. *Quaternary International*, 171(172), 160-178.
- Giacco B., Isaia R., Fedele F.G., Di Canzio E., Hoffecker J., Ronchetti A., Sinitsov A.A., Anikovich M., Lisitsyn S.N., Popov V.V. (2008) - The Campanian Ignimbrite and codola tephra layers: two temporal/stratigraphic markers for the early upper palaeolithic in southern Italy and eastern Europe. *Journal of Volcanology and Geothermal Research*, 177 (1), 208-226.
- Doi: 10.1016/j.volgeores.2007.10.007
- Giacco B., Marra F., Hajdas I., Karner D.B., Renne P.R., Sposato A. (2009) - ⁴⁰Ar/³⁹Ar and ¹⁴C geochronology of the Albano maar deposits: implications for defining the age and eruptive style of the most recent explosive activity at the Albano Hills Volcanic District, Italy. *Journal of Volcanology and Geothermal Research*, 185, 203-213.
- Giacco B., Nomade S., Wulf S., Isaia R., Sottili G., Cavuoto G., Galli P., Messina P., Sposato A., Sulipizio R., Zanchetta G. (2012) - The late MIS 5 Mediterranean tephra markers: a reappraisal from peninsular Italy terrestrial records. *Quaternary Science Reviews*, 56, 31-45.
- Giacco B., Arienzo I., Sottili G., Castorina F., Gaeta M., Nomade S., Galli P., Messina P. (2013a) - Isotopic (Sr-Nd) and major element fingerprinting of distal tephras: An application to the Middle-Late Pleistocene markers from the Colli Albani volcano, central Italy. *Quaternary Science Reviews*, 67, 190-206.
- Giacco B., Castorina F., Nomade S., Scardia G., Voltaggio M., Sagnotti L. (2013b) - Revised Chronology of the Sulmona Lacustrine Succession, Central Italy. *Journal of Quaternary Science*, 28, 545-551.
- Giacco B., Galli P., Peronace E., Arienzo I., Nomade S., Cavinato G.P., Mandini M., Messina P., Sottili G. (2014) - A 560-440 ka tephra record from the Mercurio Basin, Southern Italy: volcanological and tephrostratigraphic implications. *Journal of Quaternary Science*, 29, 232-248.
- Giacco B., Regattieri E., Zanchetta G., Nomade S., Renne P.R., Sprain C.J., Drysdale R.N., Tzedakis P.C., Messina P., Scardia G., Sposato A., Bassinot

- F. (2015) - Duration and dynamics of the best orbital analogue to the present interglacial. *Geology*, 43, 603-606.
- Giaccio B., Niespolo E.M., Pereira A., Nomade S., Renne P.R., Albert P.G., Arienzo I., Regattieri E., Wagner B., Zanchetta G., Gaeta M., Galli P., Mannella G., Peronace E., Sottili G., Florindo F., Leicher N., Marra F., Tomlinson E.L. (2017a) - First integrated tephrochronological record for the last ca. 190 kyr from the Fucino Quaternary lacustrine succession, central Italy. *Quaternary Science Reviews*, 158, 211-234.
- Giaccio B., Hajdas I., Isala R., Deino A., Nomade S. (2017b) - High-precision ^{14}C dating and $^{40}\text{Ar}/^{39}\text{Ar}$ dating of the Campanian Ignimbrite (Y-5) reconciles the time-scales of climatic-cultural processes at 40 ka. *Scientific Reports*, 7, 45940. doi: 10.1038/srep45940.
- Giaccio B., Leicher N., Mannella G., Monaco L., Regattieri E., Wagner B., Zanchetta G., Gaeta M., Marra F., Nomade S., Palladino D.M., Pereira A., Scheidt S., Sottili G., Wonik T., Wulf S., Zeeden C., Ariztegui D., Cavatorta G.P., Dean J.R., Florindo F., Leng M.J., Macri P., Niespolo E., Renne P.R., Rolf C., Sadori L., Thomas C., Tzedakis P.C. (2019) - Extending the tephra and palaeoenvironmental record of the Central Mediterranean back to 430 ka: A new core from Fucino Basin, central Italy. *Quaternary Science Reviews*, 225, 106003.
- Giannandrea P., La Volpe L., Principe C., Schiattarella M. (2006) - Unità stratigrafiche a limiti inconformi e storia evolutiva del vulcano medio-pleistocenico di Monte Vulture (Appennino meridionale, Italia). *Bullettino della Società Geologica Italiana*, 125, 67-92.
- Giannetti B. (1996a) - The geology of the Yellow Trachytic Tuff, Roccamonfina Volcano. *Journal of Volcanology and Geothermal Research*, 71 (1), 53-72.
- Giannetti B. (1996b) - Volcanology of trachytic and associated basaltic pyroclastic deposits at Roccamonfina volcano, Roman Region, Italy. *Journal of Volcanology and Geothermal Research*, 71 (2-4), 229-248.
- Giannetti B. (2001) - Origin of the calderas and evolution of Roccamonfina volcano (Roman Region, Italy). *Journal of Volcanology and Geothermal Research*, 106(3-4), 301-319.
- Giannetti B., De Casa G. (2000) - Stratigraphy, chronostratigraphy, and sedimentology of ignimbrites from the white trachytic tuff, Roccamonfina Volcano, Italy. *Journal of Volcanology and Geothermal Research*, 96(3-4), 243-255.
- Giano S.I., Pescatore E., Biscione M., Masini E., Bentivenga M. (2022) - Geo- and Archaeo-heritage in the Mount Vulture Area: list, data management, communication, and dissemination. A Preliminary note. *Geoheritage*, 14, 10.
- Gillet P.-Y., Keller J. (1993) - Radiochronological dating of Stromboli. *Acta Vulcanologica*, 3, 69-77.
- Gillet P.Y., Kieffer G., Romano R. (1994) - The evolution of Mount Etna in the light of potassium-argon dating. *Acta Vulcanologica* 5, 81-87.
- Giordano G. (1998a) - Facies characteristics and magma-water interaction of the White Trachytic Tufts (Roccamonfina Volcano, southern Italy). *Bulletin of Volcanology*, 60, 10-26.
- Giordano G. (1998b) - The effect of paleotopography on lithic distribution and facies associations of small volume ignimbrites: the WTT Cupa (Roccamonfina volcano, Italy). *Journal of Volcanology and Geothermal Research*, 87(1-4), 255-273.
- Giordano G., CARG Team (2010) - Stratigraphy, volcano tectonics and evolution of the Colli Albani volcanic field. In: R. Funiciello, G. Giordano (Eds.), *The Colli Albani Volcano*. Special Publications of IAVCEI, 3, Geological Society of London, London, 43-98.
- Giordano G., De Astis G. (2021) - The summer 2019 basaltic Vulcanian eruptions (paroxysms) of Stromboli. *Bulletin of Volcanology*, 83, 1. doi: 10.1007/s00445-020-01423-2
- Giordano G., Caricchi L. (2022) - Determining the state of activity of transcrustal magmatic systems and their volcanoes. *Annual Review Earth and Planetary Science*, 50, 231-259.
- Giordano G., de Rita D., Cas R., Rodani S. (2002) - Valley pond and ignimbrite veneer deposits in the small-volume phreatomagmatic 'Peperino Albano' basic ignimbrite, Lago Albano maar, Colli Albani volcano, Italy: influence of topography. *Journal of Volcanology and Geothermal Research*, 118 (1-2), 131-144.
- Giordano G., De Benedetti A.A., Diana A., Diana G., Gaudiose F., Marasco F., Miceli M., Mollo S., Cas R.A.F., Funiciello R. (2006) - The Colli Albani mafic caldera (Roma, Italy): Stratigraphy, structure and petrology. *Journal of Volcanology and Geothermal Research*, 155, 49-80.
- Giuffrida M., Viccaro M. (2017) - Three years (2011-2013) of eruptive activity at Mt. Etna: Working modes and timescales of the modern volcano plumbing system from microanalytical studies of crystals. *Earth Science Reviews*, 171, 289-322.
- Giuffrida M., Viccaro M., Ottolini L. (2018) - Ultrafast syn-eruptive degassing and ascent trigger high-energy basic eruptions. *Scientific Reports*, 8, 147.
- Giuffrida M., Nicotra E., Viccaro M. (2020) - Changing modes and rates of mafic magma supply at Pantelleria (Sicily Channel, Southern Italy): new perspectives on the volcano factory drawn upon olivine records. *Journal of Petrology*, 61 (5), egaa051.
- Giuffrida G., Scandura M., Costa G., Zuccarello F., Sciotto M., Cannata A., Viccaro M. (2021) - Tracking the summit activity of Mt. Etna volcano between July 2019 and January 2020 by integrating petrological and geophysical data. *Journal of Volcanology and Geothermal Research*, 418, 107350.
- Gozzi F., Gaeta M., Freda C., Mollo S., Di Rocca T., Marra F., Dallai L., Pack A. (2014) - Primary magmatic calcite reveals origin from crustal carbonate. *Lithos*, 190-191, 191-203.
- Granieri D., Carapezza M.L., Chiodini G., Avino R., Caliro S., Ranaldi M., Ricci T., Tarchini L. (2006) - Correlated increase in CO_2 fumareolic content and diffuse emission from La Fossa crater (Vulcano, Italy): Evidence of volcanic unrest or increasing gas release from a stationary deep magma body? *Geophysical Research Letters*, 33, L13316.
- Groppelli G., Vierck-Goette L. (Eds.) (2010) - *Stratigraphy and geology of volcanic areas*. The Geological

- Society of America Special Paper, pp. 464. ISBN 978-0-8137-2464-5
- Günther J., Prelević D., Mertz D., Rocholl A., Mertz-Kraus R., Conticelli S. (2023) - Subduction-legacy and Olivine Monitoring for Mantle-heterogeneities of the Sources of Ultrapotassic Magmas: the Italian case study. *Geochemistry, Geophysics, Geosystems*, 24. doi: 10.1029/2022GC010709
- Gurioli L., Sulpizio R., Cioni R., Sbrana A., Santacroce R., Luperini W., Andronico D. (2010) - Pyroclastic flow hazard assessment at Somma-Vesuvius based on the geological record. *Bulletin of Volcanology*, 72, 1021-1038. doi: 10.1007/s00445-010-0379-2
- Harris A.J.L., Steffka A., Calvari S., Spampinato L. (2011) - Thirty years of satellite-derived lava discharge rates at Etna: Implications for steady volumetric output. *Journal of Geophysical Research*, 116, B06204.
- Hieke Merlin O. (1964) - Le vulcaniti del settore orientale del Monte Vulture (Lucania). Memorie dell'Istituto di Geologia e Mineralogia dell'Università di Padova, XXVI, 3-67.
- Hieke Merlin O. (1967) - I prodotti vulcanici del Monte Vulture (Lucania). Memorie dell'Istituto di Geologia e Mineralogia dell'Università di Padova, 28, 1-67.
- Hornig-Kjarsgaard I., Keller J., Koberski U., Städlebauer E., Francalanci L., Lenhart R. (1993) - Geology, stratigraphy and volcanological evolution of the island of Stromboli, Aeolian arc, Italy. *Acta Vulcanologica*, 3, 21-68.
- Iezzi G., Caso C., Ventura G., Vallefuoco M., Cavallo A., Behrens H., Mollo S., Paltrinieri D., Signanini P., Vetere F. (2014) - First documented deep submarine explosive eruptions at the Marsili Seamount (Tyrrenian Sea, Italy): A case of historical volcanism in the Mediterranean Sea. *Gondwana Research*, 25, 764-774.
- Ippolito F., Ortolani F., Russo M. (1973) - Struttura marginale tirrenica dell'Appennino Campano: reinterpretazione di dati di ricerche di idrocarburi. *Memorie della Società Geologica Italiana*, 12, 229-249.
- Iurino D.A., Mecozzi B., Iannucoli A., Moscarella A., Strani F., Bona F., Gaeta M., Sardella R. (2022) - A middle Pleistocene wolf from central Italy provides insights on the first occurrence of Canis lupus in Europe. *Scientific Reports*, 12(1), 2882.
- Jolivet L., Faccenna C., D'Agostino N., Fournier M., Worrall D. (1999) - The kinematics of back-arc basins, examples from the Tyrrhenian, Aegean and Japan Seas. Geological Society, London, Special Publications, 164(1), 21-53.
- Jones A.P., Kostoula T., Stoppa F., Wolley A.R. (2000) - Petrography and mineral chemistry of mantle xenoliths in a carbonate-rich melilitic tuff from Mt. Vulture volcano, southern Italy. *Mineralogical Magazine*, 64, 341-361.
- Jordan N.J., Rotolo S.G., Williams R., Speranza F., McIntosh WC., Branney MJ., Scaillet S. (2018) - Explosive eruptive history of Pantelleria, Italy: repeated caldera collapse and ignimbrite formation at a peralkaline volcano. *Journal of Volcanology and Geothermal Research*, 349, 67-73.
- Jorgenson C., Caricchi L., Bouvier A.-S., Giordano G., Weber G., Marzer F. (2020) - Consequences of CO₂-magma interaction for explosive volcanism at Colli Albani (Italy). In AGU Fall Meeting Abstracts, V038-0002. 2020.
- Kahl M., Viccaro M., Ubide T., Morgan D., Dingwell D.B. (2017) - A branched magma feeder system during the 1659 eruption of Mt. Etna: evidence from a time-integrated study of zoned olivine phenocryst populations. *Journal of Petrology*, 58, 443-472.
- Kerner D.B., Marra F., Renne P.R. (2001) - The History of the Monti Sabatini and Alban Hills Volcanoes: Groundwork for Assessing Volcanic-Tectonic Hazards for Rome. *Journal of Volcanology and Geothermal Research*, 107, 185-219.
- Keller J. (1980) - The Island of Vulcano. *Rendiconti della Società Italiana di Mineralogia e Petrologia*, 36, 369-414.
- Keller J., Ryan W.B.F., Ninkovich D., Altherr R. (1978) - Explosive volcanic activity in the Mediterranean over the past 200,000 yr as recorded in deep-sea sediments. *Geological Society of America Bulletin*, 89, 591-604.
- Kelly T.J., Carey S., Pistolesi M., Rossi M., Croft-Bell K.L.C., Roman C., Marani M. (2014) - Exploration of the 1891 Foerstner submarine vent site (Pantelleria, Italy): Insights into the formation of basaltic balloons. *Bulletin of Volcanology*, 76, 844. doi: doi.org/10.1007/s00445-014-0844-4
- La Volpe L., Principe C. (1989a) - Stratigrafia e storia eruttiva del Monte Vulture - Revisione ed aggiornamenti. *Bullettino Gruppo Nazionale per la Vulcanologia*, 5, 889-902.
- LaBerge R.D., Giordano G., Cas R.A.F., Allières L. (2006) - Syn-depositional substrate deformation produced by the shear force of a pyroclastic density current: an example from the Pleistocene ignimbrite at Monte Cimino, northern Lazio, Italia. *Journal of Volcanology and Geothermal Research*, 158, 307-320.
- LaBerge R.D., Porreca M., Mattei M., Giordano G., Cas R.A.F. (2008) - Meandering flow of a pyroclastic density current documented by the anisotropy of magnetic susceptibility (AMS) in the quartz latite ignimbrite of the Pleistocene Monte Cimino volcanic centre (central Italy). *Tectonophysics*. doi: 10.1016/j.tecto.2008.09.009
- Lane C.S., Andric M., Culen V.L., Blockley S.P.E. (2011) - The occurrence of distal Icelandic and Italian tephra in the Lateglacial of lake Bled, Slovenia. *Quaternary Science Reviews*, 30, 1013-1018.
- Lane C.S., Brauer A., Martin-Puertas C., Blockley S.P.E., Smith V.C., Tomlinson E.L. (2015) - The Late Quaternary tephrostratigraphy of annually laminated sediments from Meerfelder Maar, Germany. *Quaternary Science Reviews*, 122, 192-206.
- Lardini D., Nappi G. (1987) - I cicli eruttivi del complesso vulcanico Cimino. *Rendiconti della Società Italiana di Mineralogia e Petrologia*, 42, 141-153.
- Laurenzi M.A., Villa I.M. (1987) - ⁴⁰Ar/³⁹Ar chronostratigraphy of Vico ignimbrites. *Periodico di Mineralogia*, 56, 285-293.
- Laurenzi M., Braschi E., Casalini M., Conticelli S. (2015) - New ⁴⁰Ar/³⁹Ar dating and revision of the Geochro-

- nology of the Monte Amiata volcano, Central Italy. *Italian Journal of Geosciences*, 134, 255-265.
- Le Maître R.L., Streckeisen A., Zanettin B. (2002) - Igneous rocks. A classification and glossary of terms. Recommendations of the IUGS Subcommission on the Systematics of Igneous Rocks.
- Leicher N., Giaccio B., Zanchetta G., Wagner B., Francke A., Palladino D.M., Sulpizio R., Albert P.G., Tomlinson E.L. (2019) - Central Mediterranean explosive volcanism and tephrochronology during the last 630 ka based on the sediment record from Lake Ohrid. *Quaternary Science Reviews*, 226, 106021.
- Leicher N., Giaccio B., Zanchetta G., Sulpizio R., Albert P.G., Tomlinson E. L., Lagos M., Francke A., Wagner B. (2021) - Lake Ohrid's tephrochronological dataset reveals 1.36 Ma of Mediterranean explosive volcanic activity. *Scientific Data*, 8, 231. doi: 10.1038/s41597-021-01013-7
- Leicher N., Giaccio B., Pereira A., Nomade S., Monaco L., Mannella G., Gelli P., Peronance E., Palladino D.M., Sottili G., Zanchetta G., Wagner B. (2023) - Central Mediterranean tephrochronology between 313 and 366 ka: New insights from the Fucino paleolake sediment succession. *Boreas*. doi: 10.1111/bor.12610
- Lentini F. (1982) - The geology of the Mt. Etna basement. *Memorie della Società Geologica Italiana*, 23, 7-25.
- Leocat E. (2011) - Histoire eruptive des volcans du secteur occidental des îles Éoliennes (Sud de la Mer Tyrrhénienne, Italie). PhD thesis, University of Paris 11, Orsay.
- Liotta D. (1996) - Analisi del settore centro-meridionale del bacino pliocenico di Radicofani (Toscana Meridionale). *Bollettino della Società Geologica Italiana*, 115, 115-143.
- Lirer L., Rolandi G., Rubin M. (1991) - 14C age of the "Museum Breccia" (Campi Flegrei) and its relevance for the origin of the Campanian Ignimbrite. *Journal of Volcanology and Geothermal Research* 48, 223-227.
- Lirer L., Petrosino P., Alberico I. (2010) - Hazard and risk assessment in a complex multi-source volcanic area: the example of the Campania Region, Italy. *Bulletin of Volcanology*, 72, 411-429. doi: 10.1007/s00445-009-0334-2
- Lisicki L.E., Raymo M.E. (2005) - A Pliocene-Pleistocene stack of 57 globally distributed benthic $\delta^{18}\text{O}$ records. *Paleoceanography*, 20, PA1003. doi: 10.1029/2004PA001071
- Locardi E. (1965) - Tipi di ignimbriti di magmi mediterranei; le ignimbriti del vulcano di Vico. *Atti della Società Toscana di Scienze Naturali*, 72, 55-174.
- Lombardi G., Nicoletti M., Petruciani C. (1974) - Età delle vulcaniti acide dei complessi Tolfa-Tarquinia-Cerreto-Manziale (Lazio NordOccidentale). *Periodico Mineralogia*, 43, 181-204.
- Loreto M.F., Zitellini N., Ranero C., Palmiotto C., Manel P. (2020) - Extensional tectonics during the Tyrrhenian back-arc basin formation and a new morphotectonic map. *Basin Research*, 33, 138-158.
- Lucchi F., Keller J., Tranne C.A. (2013a) - Regional stratigraphic correlations across the Aeolian archipelago (southern Italy). *Geological Society, London, Memoirs*, 37, 55-81
- Lucchi F., Gertisser R., Keller J., Formi F., De Astis G., Tranne C.A. (2013b) - Eruptive history and magmatic evolution of the island of Salina (central Aeolian archipelago). *Geological Society, London, Memoirs*, 37, 155-211.
- Lucchi F., Tranne C.A., Peccerillo A., Keller J., Rossi P.L. (2013c) - Geological history of the Panarea volcanic group (eastern Aeolian archipelago). *Geological Society, London, Memoirs*, 37, 349-393.
- Lucchi F., Peccerillo A., Tranne C.A., Rossi P.L., Frezzotti M.L., Donati C. (2013d) - Volcanism, calderas, and magmas of the Alicudi composite volcano (western Aeolian archipelago). *Geological Society, London, Memoirs*, 37, 83-111.
- Lucchi F., Samo A., Tranne C.A., Peccerillo A., Keller J. (2013e) - Volcanism, magmatism, volcanotectonics and sea-level fluctuations in the geological history of Filicudi (western Aeolian archipelago). *Geological Society, London, Memoirs*, 37, 113-153.
- Lucchi F., Francalanci L., De Astis G., Tranne C.A., Braschi E., Klaver M. (2018) - Geological evidence for recurrent collapse-driven phreatomagmatic pyroclastic density currents in the Holocene activity of Stromboli volcano, Italy. *Journal of Volcanology and Geothermal Research*. doi: 10.1016/j.volgeores.2018.10.024
- Luhr J.F., Giannetti B. (1987) - The Brown Leucitic Tuff of Roccamonfina Volcano (Roman Region, Italy). *Contribution to Mineralogy and Petrology*, 95, 420-436.
- Lupton J., De Ronde C., Sprovieri M., Baker E.T., Bruno P.P., Italiano F., Walker S., Faure K., Leybourne M., Britton K., Greene R. (2011) - Active hydrothermal discharge on the submarine Aeolian Arc. *Journal of Geophysical Research B: Solid Earth*, 116, B02102.
- Lustrino M., Duggen S., Rosenberg C.L. (2011) - The Central-Western Mediterranean: Anomalous igneous activity in an anomalous collisional tectonic setting. *Earth-Science Reviews*, 104(1-3), 1-40.
- Macedonio G., Costa A., Folch A. (2008) - Ash fallout scenarios at Vesuvius: Numerical simulations and implications for hazard assessment. *Journal of Volcanology and Geothermal Research*, 178, 368-377.
- Macedonio G., Costa A., Scollo S., Neri A. (2016) - Effects of eruption source parameter variation and meteorological dataset on tephra fallout hazard assessment: example from Vesuvius (Italy). *Journal of Applied Volcanology*, 5, 5. doi: 10.1186/s13617-016-0045-2
- Mahood G.A., Hildreth W. (1986) - Geology of the peralkaline volcano at Pantelleria, Strait of Sicily. *Bulletin of Volcanology*, 48, 143-172.
- Malaguti A.B., Rosi M., Pistolesi M., Speranza F., Menzies M. (2022) - The contribution of palaeomagnetism, tephrochronology and radiocarbon dating to refine the last 1100 years of eruptive activity at Vulcano (Italy). *Bulletin of Volcanology*, 84, 12.
- Malaguti A.B., Branca S., Speranza F., Coltellini M., Del Carlo P., Renzulli A. (2023) - Age of the Valle del Bove formation and chronology of the post-collapse

- flank eruptions, Etna volcano (Italy). *Journal of Volcanology and Geothermal Research*, 434, 107752.
- Malinverno A., Ryan W.B.F. (1986) - Extension in the Tyrrhenian Sea and shortening in the Apennines as result of arc migration driven by sinking of the lithosphere. *Tectonics*, 5, 227-245.
- Mandarano M., Paonita A., Martelli M., Viccaro M., Nicotra E., Millar I.L. (2016) - Revealing magma degassing below closed-conduit active volcanoes: Geochemical features of volcanic rocks versus fumarolic fluids at Vulcano (Aeolian Islands, Italy). *Lithos*, 248-251, 272-287.
- Mangiacapra A., Moretti R., Rutherford M., Civetta L., Orsi G., Papale P. (2008) - The deep magmatic system of the Campi Flegrei caldera (Italy). *Geophysical Research Letters*, 35, L21304. doi: 10.1029/2008GL035550
- Marani M.P., Gamberi F. (2004) - Distribution and nature of submarine volcanic landforms in the Tyrrhenian Sea: The arc vs the back-arc. *Memorie Descrittive della Carta Geologica Italiana*, LXIV, 109-126.
- Marra F., Karner D.B. (2005) - The Albano Maar (Alban hills volcanic district, Italy): active or dormant volcano? *Il Quaternario*, 18 (2), 173-185.
- Marra F., Freda C., Scariato P., Taddeucci J., Karner D.B., Renne P.R., Gaeta M., Palladino D.M., Trigila R., Cavarretta G. (2003) - ^{40}Ar - ^{39}Ar Geochronology of the recent phase of activity of the Alban Hills Volcanic District (Rome, Italy): implications for seismic and volcanic hazards. *Bulletin of Volcanology*, 65, 227-247.
- Marra F., Taddeucci J., Freda C., Marzocchi W., Scariato P. (2004) - Recurrence of volcanic activity along the Roman Cormagmatic Province (Tyrrhenian margin of Italy) and its tectonic significance. *Tectonics*, 23, TC4013.
- Marra F., Karner D.B., Freda C., Gaeta M., Renne P.R. (2009) - Large mafic eruptions at the Alban Hills Volcanic District (Central Italy): chronostratigraphy, petrography and eruptive behavior. *Journal of Volcanology and Geothermal Research*, 179, 217-232.
- Marra F., Deocampo D., Jackson M. D., Ventura G. (2011) - The Alban Hills and Monti Sabatini volcanic products used in ancient Roman masonry (Italy): An integrated stratigraphic, archaeological, environmental and geochemical approach. *Earth-Science Reviews*, 108(3-4), 115-138.
- Marra F., Sottili G., Gaeta M., Giaccio B., Jicha B., Masotta M., Palladino D.M., Deocampo D.M. (2014) - Major explosive activity in the Sabatini Volcanic District (central Italy) over the 800-390 ka interval: geochronological - geochemical overview and tephrostratigraphic implications. *Quaternary Science Reviews*, 94, 74-101.
- Marra F., Gaeta M., Giaccio B., Jicha B., Palladino D.M., Polcari M., Sottili G., Taddeucci J., Florindo F., Stramondo S. (2016) - Assessing the volcanic hazard for Rome: ^{39}Ar - ^{37}Ar and In-SAR constraints on the most recent eruptive activity and present-day uplift at Colli Albani Volcanic District. *Geophysical Research Letters*, 43, 6898-6906.
- Marra F., Costantini L., Di Buduo G.M., Florindo F., Jicha B., Monaco L., Palladino D.M., Sottili G. (2019) - Combined glacio-eustatic forcing and volcanotectonic uplift: geomorphological and geochronological constraints on the Tiber River terraces in the eastern Vulcini Volcanic District (central Italy). *Global and Planetary Changes*, 182, 103009.
- Marra F., Jicha B., Palladino D.M., Gaeta M., Costantini L., Di Buduo G.M. (2020a) - ^{40}Ar - ^{39}Ar single crystal dates from pyroclastic deposits provide a detailed record of the 580-240 ka eruptive period at the Vulcini Volcanic District (central Italy). *Journal of Volcanology and Geothermal Research*, 398, 106904. doi: 10.1016/j.jvolgeores.2020.106904
- Marra F., Castellano C., Cuccia L., Florindo F., Gaeta M., Jicha B., Palladino D.M., Sottili G., Tertulliani A., Tolomei C. (2020b) - Monti Sabatini and Colli Albani: the dormant twin volcanoes at the gates of Rome. *Scientific Reports*, 10, 8666. doi: 10.1038/s41598-02-65394-2
- Marra F., Cardello G.L., Gaeta M., Jicha B.R., Montone P., Niespolo E.M., Nomade S., Palladino D.M., Pereira A., De Luca G., Florindo F., Frepoli A., Renne P.R., Sottili G. (2021) - The Volsi Volcanic Field (central Italy): eruptive history, magma system and implications on continental subduction processes. *International Journal of Earth Sciences*, 110, 689-718.
- Maroni M., Moratti G., Costantini A., Conticelli S., Benvenuti M.G., Pandolfi L., Bonini M., Cornamusini G., Laurenzi M.A. (2015) - Geology of the Monte Amiata region, southern Tuscany, central Italy. *Italian Journal of Geosciences*, 134(2), 171-199.
- Marzocchi W., Sandri L., Selva J. (2010) - BET_VH: a probabilistic tool for long-term volcanic hazard assessment. *Bulletin of Volcanology*, 72, 705-716.
- Mascle G. H., Tricart P., Torelli L., Bouillou J.P., Rolfo F., Lapierre H., Monié P., Depardon S., Mascle J., Peis D. (2001) - Evolution of the Sardinia Channel (Western Mediterranean): new constraints from a diving survey on Cornacina seamount off SE Sardinia. *Marine Geology*, 179, 179-201.
- Mascletta M., Gaeta M., Gozzi F., Marra F., Palladino D.M., Sottili G. (2010) - H_2O - and temperature-zoning in magma chambers: the example of the Tufo Giallo della Via Tiberina eruptions (Sabatini Volcanic District, central Italy). *Lithos*, 118(1-2), 119-130.
- Massaro S., Stocchi M., Martinez Montesinos B., Sandri L., Selva J., Sulpizio R., Giaccio B., Moscatelli M., Peronace E., Nocentini M., Isala R., Titos Luzón M., Dellino P., Naso G., Costa A. (2023) - Assessing long-term tephra fallout hazard in Southern Italy from Neapolitan volcanoes. *Natural Hazards and Earth System Sciences Discussions*, 1-35.
- Meletti M., Conticelli S., Giordano G. (2010) - The Tyrrhenian margin geological setting: from the Apennine orogeny to the K-rich volcanism. The Colli Albani Volcano, Funiciello R., Giordano G. (Eds.), Special Publication of IAVCEI, 3, The Geological Society, London, 7-27.
- Meluso L., Morra V., Di Girolamo P. (1996) - The Monte Vulture volcanic complex (Italy): evidence for distinct parental magmas and for residual melts with melilite. *Mineralogy and Petrology*, 56, 225-250.

- Mercalli G., Silvestri O. (1891) - Le eruzioni dell'isola di Vulcano, incominciate il 3 Agosto 1888 e terminate il 22 Marzo 1890. Annali dell'Ufficio Centrale di Meteorologia e Geodinamica, 10 (4), 1-213.
- Muschietti S., Albert P.G., Lucchi F., Sulpizio R., Smith V.C., Kearne R., Tranne C.A. (2020) - Frequent activity on Vulcano (Italy) spanning the last 80 ky: New insights from the chemo-stratigraphy of the Brown Tufts. Journal of Volcanology and Geothermal Research, 406.
- Métrich N., Santacroce R., Savelli C. (1988) - Ventotene, a potassic quaternary volcano in central Tyrrhenian Sea. Rendiconti Società Italiana Mineralogia Petrologia, 43, 1195-1213.
- Métrich N., Allard P., Spillaert N., Andronico D., Burton M. (2004) - 2001 flank eruption of the alkali- and volatile-rich primitive basalt responsible for Mount Etna's evolution in the last three decades. Earth and Planetary Science Letters, 228, 1-17.
- Métrich N., Bertagnini A., Pistolesi M. (2021) - Paroxysms at Stromboli Volcano (Italy): Source, Genesis and Dynamics. Frontiers in Earth Sciences. DOI: 10.3389/feart.2021.593339
- Milano G., Passaro S., Sprovieri M. (2012) - Present-day knowledge on the Palinuro Seamount (south-eastern Tyrrhenian Sea). Bollettino di Geofisica Teorica ed Applicata, 53, 403-416.
- Milia A., Giordano F., Nardi G. (1998) - Stratigraphic and structural evolution of Naples Harbour over the last 12 ka. Giornale di Geologia, 60, 41-52.
- Mixon E.E., Jicha B.R., Tootell D., Singer B.S. (2022) - Optimizing $^{40}\text{Ar}/^{39}\text{Ar}$ analyses using an Isotope NGX-600 mass spectrometer. Chemical Geology, 593, 120753. DOI: 10.1016/j.chemgeo.2022.120753
- Monaco L. (2022) - Tempo and dynamics of the peri-Tyrrhenian Quaternary explosive volcanism inferred from distal archives. PhD Dissertation, XXXIV Doctorate Cycle, Ph.D. School "Vito Volterra", Earth Sciences Curriculum, Sapienza-University of Rome, pp. 267.
- Monaco L., Palladino D.M., Gaeta M., Marra F., Sottili G., Leicher N., Mannella G., Nomade S., Pereira A., Regattieri E., Wagner B., Zanchetta G., Albert P.G., Ariozzo I., D'Antonio M., Petrosino P., Manning C.J., Giaccio B. (2021) - Mediterranean tephrostratigraphy and peri-Tyrrhenian explosive activity re-evaluated in light of the 430-365 ka record from Fucino Basin (central Italy). Earth-Science Reviews, 220, 103706.
- Monaco L., Leicher N., Palladino D.M., Ariozzo I., Marra F., Petrelli M., Nomade S., Pereira A., Sottili G., Conticelli S., D'Antonio M., Fabbrizio A., Jicha B.R., Mannella G., Petrosino P., Regattieri E., Tzedakis P.C., Wagner B., Zanchetta G., Giaccio B. (2022a) - The Fucino 250-170 ka tephra record: New insights on peri-Tyrrhenian explosive volcanism, central mediterranean tephrochronology, and timing of the MIS 8-6 climate variability. Quaternary Science Reviews, 296, 107797.
- Monaco L., Palladino D.M., Albert P.G., Ariozzo I., Conticelli S., Di Vito M., Fabbrizio A., D'Antonio M., Isaia R., Manning C.J., Nomade S., Pereira A., Petrosino P., Sottili G., Sulpizio R., Zanchetta G., Giaccio B. (2022b) - Linking the Mediterranean MIS 5 tephra markers to Campi Flegrei (southern Italy) 109-92 ka explosive activity and refining the chronology of MIS 5c-d millennial-scale climate variability. Global and Planetary Change, 211, 103785.
- Montesinos B.M., Luzón M.T., Sandri L., Rudy O., Cheptsov A., Macedonio G., Folch A., Barsotti S., Selva J., Costa A. (2022) - On the feasibility and usefulness of high-performance computing in probabilistic volcanic hazard assessment: An application to tephra hazard from Campi Flegrei. Frontiers in Earth Science, 10:941789. DOI: 10.3389/feart.2022.941789
- Nappi G. (1969) - Genesi ed evoluzione della Caldera di Latera: Bollettino del Servizio Geologico Italiano, 90, 61-81.
- Nappi G., Marini A. (1986) - I cicli eruttivi dei Vulsini orientali nell'ambito della vulcanotettonica del Compresso. Memorie della Società Geologica Italiana, 35, 679-687.
- Nappi G., Renzulli A., Santi P. (1991) - Evidence of incremental growth in the Vulsinian calderas (central Italy). In: Verma-Surendra P. (Ed.), Calderas: Genesis, Structure and Unrest. Journal of Volcanology and Geothermal Research, 47, 13-31.
- Nappi G., Capaccioni B., Mattioli M., Mancini E., Valentini L. (1994) - Plinian fall deposits from Vulsini volcanic district (central Italy). Bulletin of Volcanology, 56, 502-515.
- Nappi G., Renzulli A., Santi P., Gillot Y. (1995) - Geological evolution and geochronology of the Vulsini volcanic district (central Italy). Bollettino della Società Geologica Italiana, 114, 599-613.
- Nappi G., Chiocchini U., Bonomo R., Madonna S., Mattioli M., Ricci V., Vita L. (2022) - Note illustrative della carta geologica d'Italia alla scala 1:50.000, F. 345 - Viterbo. ISPRA - Servizio Geologico d'Italia, Roma, pp. 273. DOI: 10.15161/car.it/75774
- Nazzaro A. (1997) - Il Vesuvio: storia eruttiva e teorie vulcanologiche. Liguori ed., Napoli, Italy, pp. 374.
- Neri M., Rivalta E., Maccaferri F., Acciolla V., Cirrincione R. (2018) - Etnean and Hyblean volcanism shifted away from the Malta Escarpment by crustal stresses. Earth and Planetary Science Letters, 486, 15-22.
- Nicoletti M. (1969) - Datazioni Ar-K di alcune vulcaniti delle Regioni vulcaniche Cimina e Vicaria. Periodico di Mineralogia, 38, 1-20.
- Nicoletti M., Petruccioli C., Pro M., Triglia R. (1981) - Nuove datazioni vulsine per uno schema di evoluzione dell'attività vulcanica. Nota II il quadrante sud-occidentale. Periodico di Mineralogia, 50, 141-169.
- Nicotra E., Viccaro M. (2012) - Unusual magma storage conditions at Mt. Etna (Southern Italy) as evidenced by plagioclase megacryst-bearing lavas: implications for the plumbing system geometry and summit caldera collapse. Bulletin of Volcanology, 74, 795-815.
- Nicotra E., Viccaro M., De Rosa R., Sapienza M. (2014) - Volcanological evolution of the Rivi-Capo volcanic complex at Salina, Aeolian Islands: magma storage processes and ascent dynamics. Bulletin of Volcanology, 76, 1001-1018.

- ology, 75, 840.
- Nicotra E., Giuffrida M., Viccaro M., Donato P., D'Orlano C., Paonita A., De Rosa R. (2018) - Timescales of pre-eruptive magmatic processes at Vulcano (Aeolian Islands, Italy) during the last 1000 years. *Lithos*, 316-317, 347-365.
- Nicotra E., Minniti M., Donato P., De Rosa R. (2020) - Insights into the Eruptive Dynamics of Small Caldera-Forming Eruptions: The Case Study of the Welded Scoriae of Vulcano (Aeolian Islands, Italy). *Frontiers in Earth Science*, 8, 223. DOI: 10.3389/feart.2020.00223
- Orlando A., Conticelli S., Marietti P., Vaggelli G. (1994) - The basement of northern Vulcanian volcanic district as inferred from the study of crustal xenoliths from the Torre Alfina lavas, Viterbo, central Italy. *Memoire della Società Geologica Italiana*, 48, 681-688.
- Ori G., Gallo G., Zanchi A. (1991) - Simple-shearing block resurgence in caldera depressions. A model from Pantelleria and Ischia. *Journal of Volcanology and Geothermal Research*, 47, 1-11. DOI: 10.1016/0377-0273(91)90097
- Ori G., D'Antonio M., de Vita S., Gallo G. (1992) - The Neapolitan Yellow Tuff, a large-magnitude trachytic phreatoplinarian eruption: eruptive dynamics, magma withdrawal and caldera collapse. *Journal of Volcanology and Geothermal Research*, 53, 275-287.
- Ori G., Civetta L., D'Antonio M., Di Girolamo P., Picchi M. (1995) - Step-filling and development of a three-layers magma chamber: the Neapolitan Yellow Tuff case history. *Journal of Volcanology and Geothermal Research*, 67, 291-312.
- Ori G., de Vita S., Di Vito M. (1996) - The restless, resurgent Campi Flegrei nested caldera (Italy): constraints on its evolution and configuration. *Journal of Volcanology and Geothermal Research*, 74, 179-214.
- Ori G., Civetta L., Del Gaudio C., de Vita S., Di Vito, Isaia R.M., Petruzzuoli S., Ricciardi G., Ricco C. (1999) - Short term ground deformations and seismicity in the nested Campi Flegrei caldera Italy: an example of active block resurgence in a densely populated area. *Journal of Volcanology and Geothermal Research*, 91, 415-451.
- Ori M., Ori G., Pappalardo L., Fisher R.V. (2003) - Anisotropy of magnetic susceptibility studies of depositional processes in the Campanian Ignimbrite, Italy. *Bulletin of Volcanology*, 65, 55-72.
- Pabet S., Werner G., Civetta L., Tesoro R. (2008) - Magma chamber evolution prior to the Campanian Ignimbrite and Neapolitan Yellow Tuff eruptions (Campi Flegrei, Italy). *Bulletin of Volcanology*, 70, 961-976. DOI: 10.1007/s00445-007-0180-z
- Palano M., Viccaro M., Zuccarello F., Gresta S. (2017) - Magma transport and storage at Mt. Etna (Italy): a review of geodetic and petrological data for the 2002-03, 2004 and 2006 eruptions. *Journal of Volcanology and Geothermal Research*, 347, 149-164.
- Palladino D.M., Agoeta E. (1997) - Pumice fall deposits of the Western Vulsini Volcanoes (Central Italy). *Journal of Volcanology and Geothermal Research*, 78, 77-102.
- Palladino D.M., Giordano G. (2019) - On the mobility of pyroclastic currents in light of deposit thickness and clast size trends. *Journal of Volcanology and Geothermal Research*, 384, 64-74.
- Palladino D.M., Pettini M. (2020) - Source- vs topographic-forcing in pyroclastic currents: the case of the Orvieto-Bagnoregio Ignimbrite, Vulsini, central Italy. *Periodico di Mineralogia*, 89, 217-229.
- Palladino D.M., Simeti S. (2005) - Eruptive dynamics and caldera-collapse during the Orvieto eruption, Vulsini, Italy. *Bulletin of Volcanology*, 67, 423-440.
- Palladino D.M., Valentine G.A. (1995) - Coarse-tail vertical and lateral grading in pyroclastic flow deposits of the Latara Volcanic Complex (Vulsini, Central Italy): origin and implications for flow dynamics. *Journal of Volcanology and Geothermal Research*, 69, 343-364.
- Palladino D.M., Gaeta M., Marra F. (2001) - A large K-foiditic hydromagmatic eruption from the early activity of the Alban Hills Volcanic District, Italy. *Bulletin of Volcanology*, 63, 345-359.
- Palladino D.M., Simeti S., Sottilli G., Triglia R. (2010) - Integrated approach for the reconstruction of stratigraphy and geology of Quaternary volcanic terrains: an application to the Vulsini Volcanoes (central Italy). In: Groppelli G., Viereck L. (Eds.), "Stratigraphy and geology in volcanic areas", Geological Society of America, Special Papers, 464, 66-84.
- Palladino D.M., Gaeta M., Giacchio B., Sottilli G. (2014) - On the anatomy of magma chamber and caldera collapse: the example of trachy-phonolitic explosive eruptions of the Roman Province (central Italy). *Journal of Volcanology and Geothermal Research*, 281, 12-28.
- Paone A. (2013) - Petrogenesis of trachyte and rhyolite magmas on Ponza Island (Italy) and its relationship to the Campanian magmatism. *Journal of Volcanology and Geothermal Research*, 267, 15-29.
- Paonita A., Federico C., Bonfanti P., Capasso G., Inguaggiato S., Italiano F., Madonia P., Pecoraro G., Sortino F. (2013) - The Episodic and Abrupt Geochemical Changes at La Foosa Fumeroles (Vulcano Island, Italy) and Related Constraints on the Dynamics, Structure, and Compositions of the Magmatic System. *Geochimica et Cosmochimica Acta*, 120, 158-178.
- Pappalardo L., Civetta L., D'Antonio M., Deino A., Di Vito M.A., Ori G., Carandente A., de Vita S., Isaia R., Picchi M. (1999) - Chemical and Sr - isotopical evolution of the Phlegraean magmatic system before the Campanian Ignimbrite (37 ka) and the Neapolitan Yellow Tuff (12 ka) eruptions. *Journal of Volcanology and Geothermal Research*, 91, 141-168.
- Pappalardo L., Civetta L., de Vita S., Di Vito M.A., Ori G., Carandente A., Fisher R.V. (2002) - Timing of magma extraction during the Campanian Ignimbrite eruption (Campi Flegrei caldera). *Journal of Volcanology and Geothermal Research*, 114, 479-497.
- Pasquare G., Chiesa S., Vezzoli L., Zanchi A. (1983) - Evoluzione paleogeografica e strutturale di parte del Toceana meridionale a partire dal Miocene superiore. *Memoire della Società Geologica Italiana*, 25, 145-157.

- Pasquare G., Seri G., Vezzoli L. (1985) - Carta geologica dell'area della Media Valle Latina. Scala 1:50 000. Progetto finalizzato Geodinamica. Sottoprogetto: Sorveglianza dei vulcani e rischio vulcanico. In: Carte tematiche sul vulcanismo recente. Istituto di Geologia, University of Milan, Italy.
- Paterno M., Guichard F., Duplessy J.C., Siani G., Sulpizio R., Labeyrie J. (2008) - A 90,000-200,000 yrs marine tephra record of Italian volcanic activity in the Central Mediterranean Sea. *Journal of Volcanology and Geothermal Research*, 177, 187-196.
- Peccerillo A. (2005) - Plio-Quaternary Volcanism in Italy. Springer-Verlag, Berlin Heidelberg, pp. 365.
- Peccerillo A. (2017) - Cenozoic Volcanism in the Tyrrhenian Sea Region. IAVCEI, Advances in Volcanology, Springer (ed.), Barcelona, Spain, pp. 399.
- Peccerillo A., Conticelli S., Manetti P. (1987) - Petrological characteristics and the genesis of the recent magmatism of South Tuscany and North Latium. *Periodico di Mineralogia*, 56, 157-173.
- Peccerillo A., De Astis G., Faraone D., Fomi F., Frezzotti M.L. (2013) - Compositional variations of magmas in the Aeolian arc: Implications for petrogenesis and geodynamics. *Geological Society of London Memoirs*, 37, 1, 491-510. doi:10.1144/M37.15
- Pereira A., Monaco L., Marras F., Nomade S., Gaeta M., Leicher N., Palladino D.M., Scotti G., Guillou H., Sciau V., Giacomo B. (2020) - Tephrochronology of the central Mediterranean MIS 11c interglacial (~425-395 ka): New constraints from the Vico volcano and Tiber delta, central Italy. *Quaternary Sciences Reviews*, 243, 106470.
- Perini G. (1997) - Evoluzione magmatologica del vulcano di Vico. Ph.D. thesis, Università degli Studi di Firenze.
- Perini G., Conticelli S. (2002) - Crystallization conditions of Leucite-Bearing magmas and their implications on the magmatological evolution of ultrapotassic magmas: The Vico Volcano, Central Italy. *Mineralogy and Petrology*, 74, 253-276.
- Perini G., Conticelli S., Francalanci L. (1997) - Inferences on the volcanological history of the Vico volcano, Roman Magmatic Province, Central Italy: stratigraphic, petrographic and geochemical data. *Mineralogica et Petrographica Acta*, 40, 67-93.
- Perini G., Conticelli S., Francalanci L., Davidson J.P. (2000) - The relationship between potassic and calc-alkaline post-orogenic magmatism at Vico volcano, central Italy. *Journal of Volcanology and Geothermal Research*, 95, 247-272.
- Perini G., Tepley F.J. III, Davidson J.P., Conticelli S. (2003) - The origin of K-feldspar megacrysts hosted in alkali-potassic rocks: track for low scale mantle heterogeneity. *Lithos*, 66, 223-240.
- Perini G., Francalanci L., Davidson J.P., Conticelli S. (2004) - Evolution and Genesis of Magmas from Vico Volcano, Central Italy: Multiple Differentiation Pathways and Variable Parental Magmas. *Journal of Petrology*, 45(1), 139-182.
- Pescatore T., Rolandi G. (1981) - Preliminary observations on stratigraphy of volcanoclastic deposits of the SW sector of Campi Flegrei. *Bollettino della Società Geologica Italiana*, 100, 233-254.
- Petrosino P., Jicha B.R., Mazzeo F.C., Russo Ermoli E. (2014) - A high-resolution tephrochronological record of MIS 14-12 in the Southern Apennines (Acerno Basin, Italy). *Journal of Volcanology and Geothermal Research*, 274, 34-50.
- Petrosino P., Jicha B.R., Mazzeo F.C., Claranti N., Girone A., Malorano P., Marino M. (2015) - The Montalbano Jonico marine succession: an archive for distal tephra layers at the Early-Middle Pleistocene boundary in southern Italy. *Quaternary International*, 383, 89-103. doi:10.1016/j.quaint.2014.10.049
- Petrosino P., Arienza I., Mazzeo F.C., Natale J., Petrelli M., Milia A., Perugini D., D'Antonio M. (2018) - The San Gregorio Magno lacustrine basin (Campania, southern Italy): improved characterization of the tephrostratigraphic markers based on trace elements and isotopic data. *J. Quat. Sci.* 34, 393-404.
- Piochi M., Bruno P.P., De Astis G. (2005) - Relative roles of rifting tectonics and magma uprising processes: inferences from geophysical, structural and geochemical data of the Neapolitan volcanic region (southern Italy). *Geochemistry Geophysics and Geosystems*, 6, Q07006. doi:10.1029/2004GC000885
- Pistolesi M., Rosi M., Malaguti A.B., Lucchi F., Tranne C.A., Speranza F., Albert P.G., Smith V.C., Di Roberto A., Billotta E. (2021) - Chrono-stratigraphy of the youngest (last 1500 years) rhyolitic eruptions of Lipari (Aeolian Islands, Southern Italy) and implications for distal tephra correlations. *Journal of Volcanology and Geothermal Research*, 420, 107397.
- Poli S., Chiesa S., Gillot P.Y., Gregnanin A., Guichard F., Stella R. (1987) - Major and trace element variation versus time in the volcanic products of Ischia (Gulf of Naples), evidence of successive magmatic cycles. *Contributions to Mineralogy and Petrology*, 95, 322-335.
- Poli S., Chiesa S., Gillot P.Y., Guichard F., Vezzoli L. (1989) - Time dimension in the geochemical approach and hazard estimates of a volcanic area: The Isle of Ischia case (Italy). *Journal of Volcanology and Geothermal Research*, 36, 327-335.
- Principe C., Giannandrea P. (2006) - Storia evolutiva del Monte Vulture. In C. Principe ed. La geologia del Monte Vulture, 49-54.
- Principe C., La Volpe L. (1989) - Guida all'escursione sul Monte Vulture del 18-19 ottobre 1989. Convegno scientifico "Genesi e aspetti geodinamici del vulcanismo potassico e ultrapotassico: stato dell'arte e prospettive di ricerca". Bari, 17-18 ottobre 1989, 1-33.
- Principe C., Tanguy J.C., Arrighi S., Paiotti A., Le Goff M., Zoppi U. (2004) - Chronology of Vesuvius' activity from A.D. 79 to 1631 based on archeomagnetism of lavas and historical sources. *Bulletin of Volcanology*, 66, 703-724.
- Putignano M.L., Schiattarella M. (2010) - Geomorfologia strutturale e domini di frattura dei fondali marini pericostieri dell'Isola di Procida (Campi Flegrei insulari, Italia meridionale). *Alpine and Mediterranean Quaternary*, 23(2), 229-242.
- Putignano M.L., Orrù P.E., Schiattarella M. (2014) - Ho-

- Iocene coastline evolution of Procida Island, Bay of Naples, Italy. *Quaternary International*, 332, 115-125.
- Puxeddu M. (1971) - Studio chimico-petrografico delle vulcaniti del M. Cimino (Viterbo). Atti della Società Toscana di Scienze Naturali, Memorie, Serie A, 7B, 329-394.
- Radicati di Brozolo F., Di Girolamo P., Turi B., Oddone M. (1988) - ^{40}Ar - ^{39}Ar and K-Ar dating of K-rich rocks from the Roccamonfina Volcano, Roman Comagmatic Region, Italy. *Geochimica et Cosmochimica Acta*, 52, 1435-1441.
- Ripepe M., Lacanna G., Pistolesi M., Silengo M. C., Aluppa A., Laiola M., Massimetti F., Innocenti L., Della Schiava M., Bitetto M., La Monica F. P., Nishimura T., Rosi M., Mangione D., Ricciardi A., Genco R., Coppola D., Marchetti E., Delle Donne D. (2021) - Ground deformation reveals the scale-invariant conduit dynamics driving explosive basaltic eruptions. *Nature Communications*, 12, 1683. Doi: 10.1038/s41467-021-21722-2
- Pisica G., Speranza F., Giordano G., De Astis G., Lucchi F. (2019) - Palaeomagnetic dating of the Neostromboli succession. *Journal of Volcanology and Geothermal Research*, 371. Doi: 10.1016/j.jvoigeores.2018.12.009
- Roland G., Bellucci F., Cortini M. (2004) - A new model for the formation of the Somma Caldera. *Mineralogy and Petrology*, 80, 27-44.
- Romagnoli C. (2013) - Characteristics and morphological evolution of the Aeolian volcanoes from the study of submarine portions. Geological Society, London, Memoirs, 37, 13-26.
- Romagnoli C., Kokelaar P., Casalbore D., Chiocci F.L. (2009) - Lateral collapses and active sedimentary processes on the northwestern flank of Stromboli volcano. *Marine Geology*, 265, 101-119.
- Romagnoli C., Casalbore D., Bartoluzzi G., Bosman A., Chiocci F.L., D'Orlano F., Gamberi F., Ugi M., Marani M. (2013) - Bathymorphological setting of the Aeolian islands. Geological Society, London, Memoirs, 37, 27-36.
- Romagnoli C., Belvisi V., Intriangi S., Di Martino G., Tonielli (2020) - New insights on the evolution of the Linosa volcano (Sicily Channel) from the study of its submarine portions. *Marine Geology*, 419, 106060.
- Romano R., Sturiale C., Lentini F. (1979) - Carta Geologica del Monte Etna, Consiglio Nazionale delle Ricerche Progetto Finalizzato Geodinamica, 1:50,000-scale, Catania, Sicily.
- Rosatelli G., Stoppa F., Jones A.P. (2000) - Intrusive calcitecarbonatite occurrence from Mt. Vulture volcano, southern Italy. *Mineralogical Magazine*, 64.
- Rosi M., Santacroce R. (1983) - The A.D. 472 "Pollena" eruption: volcanological and petrological data for this poorly-known, Plinian-type event at Vesuvius. *Journal of Volcanology and Geothermal Research*, 17 (1-4), 249-271.
- Rosi M., Sbrana A. (eds) (1987) - Phlegraean Fields. Quaderni de "La Ricerca Scientifica" CNR, Rome, 114, 1-175.
- Rosi M., Principe C., Vecchi R. (1993) - The 1631 Vesuvius eruption. A reconstruction based on historical and stratigraphical data. *Journal of Volcanology and Geothermal Research*, 58(1-4), 151-182. Doi: 10.1016/0377-0273(93)90106-2
- Rosi M., Vezzoli L., Aleotti P., De Censi M. (1996) - Interaction between caldera collapse and eruptive dynamics during the Campanian Ignimbrite eruption, Phlegraean Fields, Italy. *Bulletin of Volcanology*, 57, 541-554.
- Rosi M., Vezzoli L., Castelmenzano A., Greco G. (1999) - Plinian pumice fall deposit of the Campanian Ignimbrite eruption (Phlegraean Fields Italy). *Journal of Volcanology and Geothermal Research*, 91, 179-198.
- Rosi M., Pistolesi M., Bertagnini A., Landi P., Pompilio M., Di Roberto A. (2013) - Stromboli volcano, Aeolian Islands (Italy): present eruptive activity and hazards. *Geological Society, London, Memoirs*, 37, 473-490.
- Rossi P.L., Tranne C. A., Calanchi N., Lant E. (1996) - Geology, stratigraphy and volcanological evolution of the island of Linosa (Sicily Channel). *Acta Vulcanologica*, 8, 73-90.
- Rotolo S.G., Castorina F., Cellura D., Pompilio M. (2006) - Petrology and geochemistry of submarine volcanism in the Sicily Channel Rift. *Journal of Geology*, 114 (3), 355-365.
- Rotolo S.G., La Felice S., Mangalaviti A., Landi P. (2007) - Geology and petrochemistry of the recent (<25 ka) silicic volcanism at Pantelleria Island. *Bollettino della Società Geologica Italiana*, 126, 191-208.
- Rotolo S.G., Scaillet S., La Felice S., Vita-Scaillet G. (2013) - A revision of the structure and stratigraphy of pre-Green Tuff ignimbrites at Pantelleria (Strait of Sicily). *Journal of Volcanology and Geothermal Research*, 250, 61-74.
- Rotolo S.G., Scaillet S., Speranza F., White J.C., Williams R., Jordan N.J. (2021) - Volcanological evolution of Pantelleria Island (Strait of Sicily) peralkaline volcano: a review. *Comptes Rendus Geosciences*, 353, 111-132.
- Rouchon V., Gillet P.Y., Quideleur X., Chiesa S., Floris B. (2008) - Temporal evolution of the Roccamonfina volcanic complex (Pleistocene), Central Italy. *Journal of Volcanology and Geothermal Research*, 177, 500-514.
- Russo Ermoli E., Aucelli P.P.C., Di Rollo A., Mattei M., Petrosino P., Porreca M., Reeskopf C.M. (2010) - An integrated stratigraphical approach to the Middle Pleistocene succession of the Sessano Basin (Molise, Italy). *Quaternary International*, 225, 114-127.
- Sandri L., Jolly G., Lindsay J., Howe T., Marzocchi W. (2012) - Combining long-and short-term probabilistic volcanic hazard assessment with cost-benefit analysis to support decision making in a volcanic crisis from the Auckland Volcanic Field, New Zealand. *Bulletin of Volcanology*, 74, 705-723. Doi: 10.1007/s00445-011-0556-y
- Sandri L., Thouret J.-C., Constantinescu R., Blass S., Tonini R. (2014) - Long-term multi-hazard assessment for El Misti volcano (Peru). *Bulletin of Volcanology*, 76, 771. Doi: 10.1007/s00445-013-0771-9
- Sandri L., Coeta A., Selva J., Tonini R., Macedonio G.,

- Folch A., Sulpizio R. (2016) - Beyond eruptive scenarios: assessing tephra fallout hazard from neapolitan volcanoes. *Scientific Reports*, 6, 24271-24313.
Doi: 10.1038/srep24271
- Santacroce R. ed. (1987) - Somma Vesuvius. CNR, Quaderni della Ricerca Scientifica, 114 (8), pp. 220.
- Santacroce R., Cioni R., Marianelli P., Sbrana A., Sulpizio R., Zanchetta G., Donahue D.J., Joron J. L. (2008) - Age and whole rock-glass compositions of proximal pyroclastics from the major explosive eruptions of Somma-Vesuvius: A review as a tool for distal tephrostratigraphy. *Journal of Volcanology and Geothermal Research*, 177, 1-18.
Doi: 10.1016/j.volgeores.2008.06.009
- Sartori R. (2003) - The Tyrrenian back-arc basin and subduction of the Ionian lithosphere. *Episodes*, 26 (3), 217-221.
Doi: 10.18814/epilugs/2003/v26i3/011
- Savelli C. (1987) - K/Ar ages and chemical data of volcanism in the western Pontine Islands (Tyrrenian Sea). *Bollettino della Società Geologica Italiana*, 106, 536-546.
- Savelli C. (1988) - Late Oligocene to recent episodes of magmatism in and around the Tyrrenian Sea: implications for the processes of opening in a young inter-arc basin of intra-orogenic (Mediterranean) type. *Tectonophysics*, 146, 163-181.
- Savelli C., Ligi M. (2017) - An updated reconstruction of basaltic crust emplacement in Tyrrenian sea, Italy. *Scientific Reports*, 7, 18024.
- Sbrana A., Toccaceli R.M. (2011) - Isola di Ischia. Foglio 464, Carta Geologica della Regione Campania, scala 1:10.000. Note illustrative. Regione Campania, Assessorato difesa del Suolo, pp. 216.
- Sbrana A., Marianelli P., Pasquini G. (2018) - Volcanology of Ischia (Italy). *Journal of Maps*, 14 (2), 494-503.
Doi: 10.1080/17445647.2018.1496811
- Sbrana A., Cioni R., Marianelli P., Sulpizio R., Andronico D., Pasquini G. (2020) - Volcanic evolution of the Somma-Vesuvius Complex (Italy). *Journal of Maps*, 16(2), 127-147.
Doi: 10.1080/17445647.2019.1706653
- Scaillet S., Vita-Scaillet G., Guillou H. (2008) - Oldest human footprints dated by Ar/Ar. *Earth and Planetary Science Letters*, 275, 320-325.
- Scaillet S., Rotolo S.G., La Felice S., Vita G. (2011) - High resolution $^{40}\text{Ar}/^{39}\text{Ar}$ chronosestratigraphy of the post-caldera (< 20 ka) volcanic activity at Pantelleria, Sicily Strait. *Earth Planetary Science Letters*, 309, 280-290.
- Scarpelli C., Cole P., Perrotta A. (1993) - The Neapolitan Yellow Tuff-A large volume multiphase eruption from Campi Flegrei. Southern Italy. *Bulletin of Volcanology*, 55, 343-356.
- Scarpelli C., Perrotta A., Lepore S., Calvert A. (2013) - Eruptive history of Neapolitan volcanoes: constraints from $^{40}\text{Ar}/^{39}\text{Ar}$ dating. *Geological Magazine*, 150, 03, 412-426.
- Schiattarella M., Beneduce P. (2006) - Caratteri geomorfologici, assetto strutturale ed evoluzione morfotettonica del Monte Vulture e delle aree contigue. In C. Principe ed. *La geologia del Monte Vulture*, 17-24.
- Schiattarella M., Beneduce P., Di Leo P., Giano S.I., Giannandrea P., Principe C. (2005) - Assetto strutturale ed evoluzione morfotettonica quaternaria del vulcano del Monte Vulture (Appennino Lucano). *Bollettino della Società Geologica Italiana*, 124, 543-562.
- Schmincke H.U., Behncke B., Grasso M., Raffi S. (1997) - Evolution of the Northwestern Iblean Mountains, Sicily: uplift, Pliocene/Pleistocene sea-level changes, paleoenvironment, and volcanism. *Geologische Rundschau*, 86, 637-669.
- Scollo S., Cottelli M., Bonadonna C., Del Carlo P. (2013) - Tephra hazard assessment at Mt. Etna (Italy). *Natural Hazards and Earth System Sciences*, 13, 3221-3233.
- Scribano V. (1987a) - Deep-seated xenoliths in alkaline volcanic rocks from the Hyblean Plateau (SE-Sicily). *Memorie della Società Geologica Italiana*, 38, 475-482.
- Scribano V. (1987b) - The ultramafic and mafic nodules suite in a tuff-breccia pipe from Cozzo Molino (Hyblean Plateau, Sicily). *Rendiconti della Società Italiana di Mineralogia e Petrologia*, 42, 203-217.
- Scribano V., Viccaro M., Cristofolini R., Ottolini L. (2009) - Metasomatic events recorded in ultramafic xenoliths from the Hyblean area (Southeastern Sicily, Italy). *Mineralogy and Petrology*, 95, 235-250.
- Selva J., Costa A., Marzocchi W., Sandri L. (2010) - BET_VH: exploring the influence of natural uncertainties on long-term hazard from tephra fallout at Campi Flegrei (Italy). *Bulletin of Volcanology*, 72, 717-733.
- Selva J., Costa A., Sandri L., Macedonio G., Marzocchi W. (2014) - Probabilistic short-term volcanic hazard in phases of unrest: A case study for tephra fallout. *Journal of Geophysical Research Solid Earth*, 119, 8805-8826.
Doi: 10.1002/2014jb011252
- Selva J., Costa A., De Natale G., Di Vito M. A., Isaia R., Macedonio G. (2018) - Sensitivity test and ensemble hazard assessment for tephra fallout at Campi Flegrei, Italy. *Journal of Volcanology and Geothermal Research*, 351, 1-28.
Doi: 10.1016/j.volgeores.2017.11.024
- Selva J., Acocella V., Bisson M., Caliro S., Costa A., Della Seta M., De Martino P., de Vita S., Federico C., Giordano G., Martino S., Cardaci C. (2019) - Multiple natural hazards at volcanic islands: a review for the Ischia volcano (Italy). *Journal of Applied Volcanology*, 8, 5.
Doi: 10.1186/s13617-019-0086-4
- Selva J., Bonadonna C., Branca S., De Astis G., Gambino S., Paonita A., Pistolesi M., Ricci T., Sulpizio R., Tibaldi A., Ricciardi A. (2020) - Multiple hazards and paths to eruptions: A review of the volcanic system of Vulcano (Aeolian Islands, Italy). *Earth-Science Reviews*, 207, 103188.
- Selva J., Sandri L., Taroni M., Sulpizio R., Tierz P., Costa A. (2022) - A simple two-state model interprets temporal modulations in eruptive activity and enhances multivolcano hazard quantification. *Science*

- Advances, 8(44), eabq4415.
- Serri G., Innocenti F., Manetti P. (2001) - Magmatism from Mesozoic to Present: petrogenesis, time-space distribution and geodynamics implications. Val G.B., Martini I.P. (Eds.), *Anatomy of an Orogen: The Apennines and Adjacent Mediterranean Basins*, Kluwer Academic Publisher, Great Britain, 77-104.
- Shackleton N.J. (1995) - New data on the evolution of Pliocene climatic variability. In: Vrba E.S., Denton G., Partridge H.T.C., Burklie L.H. (Eds.), *Paleoclimate and Evolution, with Emphasis on Human Origins*, New Haven, Connecticut, Yale University Press, 242-248.
- Shackleton N.J., Berger A., Peltier W.A. (1990) - An alternative astronomical calibration of the lower Pleistocene timescale based on ODP Site 677. Transactions of the Royal Society of Edinburgh: Earth Sciences, 81, 251-261.
- Smith V.C., Isaia R., Pearce N.J.G. (2011) - Tephrostratigraphy and glass compositions of post-15 kyr Campi Flegrei eruptions: implications for eruption history and chronostratigraphic markers. Quaternary Science Reviews, 30, 3638-3660.
- Soligo M.L., Tuclmei P. (2010) - Geochronology of Colli Albani volcano. In: R. Funiciello, G. Giordano (Eds.), *The Colli Albani Volcano*, Special Publications of IAVCEI, 3, Geological Society London, London (2010), 99-106.
- Solevartti F. (1983) - Geologic, Volcanologic and tectonic of the Vico-Cimino Area, Italy. Journal of Volcanology and Geothermal Research, 17, 203-217.
- Sottilli G., Palladino D.M., Zanon V. (2004) - Plinian activity during the early eruptive history of the Sabatini Volcanic District, Central Italy. Journal of Volcanology and Geothermal Research, 135, 361-379.
- Sottilli G., Palladino D.M., Marra F., Jicha B., Kamer D.B., Renne P. (2010a) - Geochronology of the most recent activity in the Sabatini Volcanic District, Roman Province, central Italy. Journal of Volcanology and Geothermal Research, 196, 20-30.
- Sottilli G., Taddeucci J., Palladino D.M. (2010b) - Constraints on magma-wall rock thermal interaction during explosive eruptions from textural analysis of cored bombs. Journal of Volcanology and Geothermal Research, 198, 27-34.
- Sottilli G., Palladino D.M., Gaeta M., Masotta M. (2012) - Origins and energetics of maar volcanoes: examples from the ultrapotassic Sabatini Volcanic District (Roman province, Central Italy). Bulletin of Volcanology, 74, 163-186.
- Sottilli G., Arienzo I., Castorina F., Gaeta M., Giaccio B., Marra F., Palladino D.M. (2019) - Time-dependent Sr and Nd isotope variations during the evolution of the ultrapotassic Sabatini Volcanic District (Roman province, Central Italy). Bulletin of Volcanology, 81, 67.
- Sparks R.S.J. (1975) - Stratigraphy and geology of the ignimbrites of Vulsini volcano, central Italy. Geologische Rundschau, 64, 497-523.
- Speranza F., Landi P., D'Alelio Caracciolo F., Pignatelli A. (2010) - Paleomagnetic dating of the most recent silicic eruptive activity at Pantelleria (Strait of Sicily). Bulletin of Volcanology, 72, 847-858.
- Speranza F., Di Chiara A., Rotolo S.G. (2012) - Correlation of welded ignimbrites on Pantelleria, using paleomagnetism. Bulletin of Volcanology, 74, 341-357.
- Stoppa F., Principe C. (1997) - Eruption style and petrology of a new carbonatitic suite from the Mt. Vulture Southern Italy: The Monticchio Lakes Formation. Journal of Volcanology and Geothermal Research, 78, 251-265.
- Stoppa F., Principe C., Giannandrea P. (2008) - Comments on: Carbonatites in a subduction system: The Pleistocene alkaliites from Mt. Vulture (southern Italy) by d'Orazio et al., (2007). Lithos, 103, 550-556.
- Stothers R.B., Rampino M.R. (1983) - Volcanic eruptions in the Mediterranean before A.D. 630 from written and archaeological sources. Journal of Geophysical Research, 88 (B8), 6357-6371.
- Suiting I., Schmincke H.U. (2010) - Iblean diatremes 2: shallow marine volcanism in the Central Mediterranean at the onset of the Messinian Salinity Crisis (Iblean Mountains, SE-Sicily) - a multidisciplinary approach. International Journal of Earth Sciences, 99, 1917-1940.
- Suiting I., Schmincke H.U. (2012) - Iblean diatremes 3: volcanic processes on a Miocene carbonate platform (Iblean Mountains, SE-Sicily): a comparison of deep vs. shallow marine eruptive processes. Bulletin of Volcanology, 74, 207-230.
- Sulli A. (2000) - Structural framework and crustal characteristics of the Sardinia Channel Alpine transect in the central Mediterranean. Tectonophysics, 324, 321-336.
- Sulpizio R., Mele D., Dellino P., La Volpe L. (2005) - A complex, subplinian-type eruption from low-viscosity, phonolitic to tephri-phonolitic: The AD 472 (Pollena) eruption of Somma-Vesuvius, Italy. Bulletin of Volcanology, 67, 743-767.
DOI: 10.1007/s00445-005-0414-x
- Sulpizio R., Bonasia R., Dellino P., Di Vito M.A., La Volpe L., Mele D., Zanchetta G., Sadori L. (2008) - Discriminating the long distance dispersal of fine ash from sustained columns or near ground ash clouds: the example of the Pomicci di Avellino eruption (Somma-Vesuvius, Italy). Journal of Volcanology and Geothermal Research, 177, 263-276.
- Sulpizio R., Folch A., Costa A., Scaini C., Dellino P. (2012) - Hazard assessment of far-range volcanic ash dispersal from a violent Strombolian eruption at Somma-Vesuvius volcano, Naples, Italy: Implications on civil aviation. Bulletin of Volcanology, 74, 2205-2218.
- Sulpizio R., Lucchi F., Forni F., Massaro S., Tranne C.A. (2016) - Unravelling the effusive-explosive transitions and the construction of a volcanic cone from geological data: The example of Monte dei Pomi, Salina Island (Italy). Journal of Volcanology and Geothermal Research, 327, 1-22.
- Tanguy J.C. (1976) - Tholeiitic basalt magmatism of Mount Etna and its relations with the alkaline series. Contributions to Mineralogy and Petrology, 66, 51-67.
- Tanguy J.C., Condomines M., Kieffer G. (1997) - Evolution of Mount Etna magma: constraints on the pre-

- sent feeding system and eruptive mechanism. *Journal of Volcanology and Geothermal Research*, 75, 221-250.
- Thompson M.A., Lindsay J.M., Gallard J.C. (2015) - The influence of probabilistic volcanic hazard map properties on hazard communication. *Journal of Applied Volcanology*, 4, 6. Doi: 10.1186/s13617-015-0023-0
- Tibaldi A. (2001) - Multiple sector collapses at Stromboli volcano, Italy: how they work. *Bulletin of Volcanology*, 63, 112-125.
- Tibaldi A., Vezzoli L. (2004) - A new type of volcano flank failure: the resurgent caldera sector collapse, Ischia, Italy. *Geophysical Research Letters*, 31(14), L14605.
- Tierz P., Sandri L., Costa A., Zaccarelli L., Di Vito M. A., Sulpizio R., Marzocchi W. (2016) - Suitability of energy cone for probabilistic volcanic hazard assessment: Validation tests at Somma-Vesuvius and Campi Flegrei (Italy). *Bulletin of Volcanology*, 78 (11), 79.
- Tinti S., Manucci A., Pagnoni G., Armigliato A., Zaniboni F. (2005) - The 30 December 2002 landslide-induced tsunamis in Stromboli: sequence of the events reconstructed from the eyewitness accounts. *Natural Hazards and Earth System Sciences*, 5, 763-775.
- Titos M., Martínez Montésinos B., Barsotti S., Sandri L., Folch A., Mingari L., Macdonald G., Costa A. (2022) - Long-term hazard assessment of explosive eruptions at Jan Mayen (Norway) and implications for air traffic in the North Atlantic. *Natural Hazards and Earth Systems Science* 22, 139-163. Doi: 10.5194/nhess-22-139-2022
- Tomlinson E.L., Arlenzo I., Wulf S., Smith V.C., Carandente A., Civetta L., Hardiman M., Orsi G., Rosi M., Thirlwall M., Muller W., Menzies M.A. (2012) - Campi Flegrei Italy: geochemistry of the proximal sources for major Mediterranean tephras (C-1, C-2, Y-3 & Y-5). *Geochimica et Cosmochimica Acta*, 93, 102-128. Doi: 10.1016/j.gca.2012.05.043
- Tomlinson E.L., Albert P.G., Wulf S., Brown R.J., Smith V.C., Keller J., Orsi G., Bourne A.J., Menzies M.A. (2014) - Age and geochemistry of tephra layers from Ischia, Italy: constraints from proximal-distal correlations with Lago Grande di Monticchio. *Journal of Volcanology and Geothermal Research*, 287, 22-39.
- Tonarini S., Armienti P., D'Orazio M., Innocenti F., Pompilio M., Petrini R. (1996) - Geochemical and isotopic monitoring of Mt. Etna 1989 - 1993 eruptive activity: bearing on the shallow feeding system. *Journal of Volcanology and Geothermal Research*, 64, 95-115.
- Tonarini S., D'Orazio M., Armienti P., Innocenti F., Scribano V. (1996) - Geochemical features of eastern Sicily lithosphere as probed by Hyblean xenoliths and lavas. *European Journal of Mineralogy*, 8, 1153-1173.
- Tonarini S., Armienti P., D'Orazio M., Innocenti F. (2001) - Subduction-like fluids in the genesis of Mt. Etna magmas: evidence from boron isotopes and fluid mobile elements. *Earth and Planetary Science Letters*, 192, 471-483.
- Tonarini S., Leeman W.P., Civetta L., D'Antonio M., Ferrara G., Necco A. (2004) - B/Nb and ^{11}B systematics in the Phleorean Volcanic District (PVD). *Journal of Volcanology and Geothermal Research*, 113, 123-139.
- Tonarini S., D'Antonio M., Di Vito M.A., Orsi G., Carandente A. (2009) - Geochemical and isotopical (B, Sr, Nd) evidence for mixing and mingling processes in the magmatic system feeding the Astromi volcano (4.1-3.8 ka) within the Campi Flegrei caldera (South Italy). *Lithos*, 107, 135-151. Doi: 10.1016/j.lithos.2008.09.012
- Trigila R., Agoeta E., Currado C., De Benedetti A.A., Freda C., Gaeta M., Palladino D.M., Rosa G. (1995) - Petrology. In: Trigila R. (Ed.), *The Volcano of the Alban Hills*. Università degli Studi di Roma "La Sapienza", Rome, Italy, 95-165.
- Trua T., Esperança S., Mazzuoli R. (1998) - The evolution of the lithospheric mantle along the N. African plate: geochemical and isotopic evidence from the tholeiitic and alkaline volcanic rocks of the Hyblean Plateau, Italy. *Contributions to Mineralogy and Petrology*, 131, 307-322.
- Trua T., Serri G., Rossi P.L. (2004) - Coexistence of IAB-type and OIB-type magmas in the southern Tyrrhenian back-arc basin: evidence from recent seafloor sampling and geodynamic implications. In: From seafloor to deep mantle: architecture of the Tyrrhenian backarc basin. *Memorie descrittive della Carta Geologica d'Italia*, LXIV, B3-96.
- Trua T., Serri G., Marani M. (2007) - Geochemical features and geodynamic significance of the southern Tyrrhenian backarc basin. *Geological Society of America, Special Paper* 418, 221-233.
- Valentine G.A., Sottili G., Palladino D.M., Taddeucci J. (2015) - Tephra ring interpretation in light of evolving maar-diatreme concepts: Stracciaccappa maar (Central Italy). *Journal of Volcanology and Geothermal Research*, 308, 19-29.
- Valentine G.A., Palladino D.M., DiemKaye K., Fletcher C. (2019) - Lithic-rich and lithic-poor ignimbrites and their basal deposits: Sovana and Sorano formations (Latera caldera, Italy). *Bulletin of Volcanology*, 81, 29.
- Ventura G. (2013) - Kinematics of the Aeolian volcanism (Southern Tyrrhenian Sea) from geophysical and geological data. *Geological Society, London, Memoirs*, 37, 3-11.
- Ventura G., Viardo G., Bruno P.P. (1999) - The role of flank failure in modifying the shallow plumbing system of volcanoes: an example from Somma-Vesuvius, Italy. *Geophysical Research Letters*, 26, 3681-3684.
- Ventura G., Milano G., Passaro S., Sprovieri M. (2013) - The Marsili ridge (Southern Tyrrhenian Sea, Italy): An island-arc volcanic complex emplaced on a 'relief' back-arc basin. *Earth Science Reviews*, 116, 85-94.
- Vezzoli L. (1988) - Island of Ischia. *Quaderni de 'La ricerca scientifica' del CNR - Progetto Finalizzato Geodinamica. Monografie finali*, 10, 134.
- Vezzoli L., Conticelli S., Innocenti F., Landi P., Manetti P., Palladino D.M., Trigila R. (1987) - Stratigraphy

- of the Latara Volcanic Complex: proposals for a new nomenclature. *Periodico di Mineralogia*, 56, 89-110.
- Vezzoli L., Principe C., Mallatti J., Arrighi S., Tanguy J.-C., Le Goff M. (2009) - Modes and times of caldera resurgence: The <10 ka evolution of Ischia Caldera, Italy, from high-precision archaeomagnetic dating. *Journal of Volcanology and Geothermal Research*, 186, 305-319.
- Viccaro M., Cristofolini R. (2008) - Nature of mantle heterogeneity and its role in the short-term geochemical and volcanological evolution of Mt. Etna (Italy). *Lithos*, 105, 272-288.
- Viccaro M., Zuccarello F. (2017) - Mantle ingredients for making the fingerprint of Etna alkaline magmas: implications for shallow partial melting within the complex geodynamic framework of Eastern Sicily. *Journal of Geodynamics*, 109, 10-23.
- Viccaro M., Nicotra E., Millar I.L., Cristofolini R. (2011) - The magma source at Mount Etna volcano: perspectives from the Hf isotope composition of historic and recent lavas. *Chemical Geology*, 281, 343-351.
- Viccaro M., Giuffrida M., Zuccarello F., Scandura M., Palano M., Gresta S. (2019) - Violent paroxysmal activity drives self-feeding magma replenishment at Mt. Etna. *Scientific Reports*, 9, 6717.
- Viccaro M., Cannata A., Cannavò F., De Rosa R., Giuffrida M., Nicotra E., Petrelli M., Sacco G. (2021) - Shallow conduit dynamics fuel the unexpected paroxysms of Stromboli volcano during the summer 2019. *Scientific Reports*, 11(1), 266. doi: 10.1038/s41598-020-79556-7
- Villa I.M., Buettner A. (2009) - Chronostratigraphy of Monte Vulture volcano (southern Italy): secondary mineral microtextures and ^{39}Ar - ^{40}Ar systematics. *Bulletin of Volcanology*, 71, 1195-1208.
- Villa I.M., Giuliani O., De Grandis G., Cioni R. (1989) - Datazioni K/Ar dei vulcani di Tolfa e Marziana (in Italian). *Bollettino Gruppo Nazionale di Vulcanologia*, 5, 1025-1026.
- Villa I.M., Calanchi I., Dinelli E., Luochini F. (1999) - Age and Evolution of the Albano Crater Lake (Roman Volcanic Province). *Acta Vulcanologica*, 11, 305-310.
- Virkler A.P., Cashman K., Giordano G., Groppe G. (2012) - Evolution of the male Villa Senni caldera-forming eruption at Colli Albani volcano, Italy, indicated by textural analysis of juvenile fragments. *Journal of Volcanology and Geothermal Research*, 235-236, 37-54.
- Walker G.P.L. (1984) - Down sag calderas, ring faults, caldera sizes, and incremental caldera growth. *Journal of Geophysical Research: Solid Earth*, 89 (B10), 8407-8416.
- Washington H.S. (1906) - The Roman comagmatic region. Carnegie Institution Washington Publications, 57, p.p. 199.
- Webster J.D., Raia F., Tappin C., De Vivo B. (2003) - Pre eruptive geochemistry of the Ignimbrite-forming magmas of the Campanian Volcanic Zone, Southern Italy, determined from silicate melt inclusions. *Mineralogy and Petrology*, 79, 99-125.
- White J.C., Neave D.A., Rotolo S.G., Parker D.F. (2020) - Geochemical constraints on basalt petrogenesis in the Strait of Sicily Rift Zone (Italy): Insights into the importance of short lengthscale mantle heterogeneity. *Chemical Geology*, 545, 1-18.
- Wohletz K., Orsi G., de Vita S. (1995) - Eruptive mechanisms of the Neapolitan Yellow Tuff interpreted from stratigraphic, chemical and granulometric data. *Journal of Volcanology and Geothermal Research*, 67, 263-290.
- Wulf S., Kraml M., Keller J., Negandank J.F.W. (2004) - Tephrochronology of the 100 ka lacustrine sediment record of Lago Grande di Monticchio (southern Italy). *Quaternary International*, 122, 7-30.
- Wulf S., Kraml M., Keller J. (2008) - Towards a detailed distal tephrostratigraphy in the Central Mediterranean: the last 20,000 yrs record of Lago Grande di Monticchio. *Journal of Volcanology and Geothermal Research*, 177, 118-132.
- Wulf S., Keller J., Paterne M., Mingrami J., Lauterbach S., Opitz S., Sottili G., Giacino B., Albert P.G., Salow C., Tomlinson E.L., Viccaro M., Brauer A. (2012) - The 100-133 ka record of Italian explosive volcanism and revised tephrochronology of Lago Grande di Monticchio. *Quaternary Science Reviews*, 58, 104-123.
- Zanchetta G., Sulpizio R., Roberts N., Cioni R., Eastwood W.J., Siani G., Caron B., Paterne M., Santacroce R. (2011) - Tephrostratigraphy, chronology and climatic events of the Mediterranean basin during the Holocene: an overview. *Holocene*, 21, 33-52.
- Zuccarello F., Schiavi F., Viccaro M. (2021) - Magma dehydration controls the energy of recent eruptions at Mt. Etna volcano. *Terra Nova*, 33, 423-429.
- Zuccarello F., Schiavi F., Viccaro M. (2022) - The eruption run-up at Mt. Etna volcano: constraining magma decompression rates and their relationships with the final eruptive energy. *Earth and Planetary Science Letters*, 597, 11782.

Ms. received: March 7, 2023
Accepted: May 24, 2023

Revised: May 20, 2023
Available online: June 29, 2023

