



children

IMPACT
FACTOR
2.835

Indexed in:
PubMed

Article

Body Composition in Adolescent PKU Patients: Beyond Fat Mass

Albina Tummolo, Rosa Carella, Giulia Paterno, Nicola Bartolomeo, Massimo Giotta, Annamaria Dicintio, Donatella De Giovanni and Rita Fischetto

Special Issue

Management of Metabolic Disorders in Adolescence Update

Edited by





Dr. Albina Tummolo



<https://doi.org/10.3390/children9091353>

Article

Body Composition in Adolescent PKU Patients: Beyond Fat Mass

Albina Tummolo ^{1,*}, Rosa Carella ¹, Giulia Paterno ¹, Nicola Bartolomeo ², Massimo Giotta ², Annamaria Dicintio ¹, Donatella De Giovanni ¹ and Rita Fischetto ¹

¹ Department of Metabolic Diseases and Clinical Genetics, Giovanni XXIII Children Hospital, Azienda Ospedaliero-Universitaria Consorziale, 70126 Bari, Italy

² Interdisciplinary Department of Medicine, University of Bari Aldo Moro, 70124 Bari, Italy

* Correspondence: albina.tummolo@policlinico.ba.it

Abstract: There is a lack of evidence on the impact on body composition of high protein intake and types of protein substitutes in PKU patients—particularly in adolescents, who are more inclined to dietary transgressions. In this observational, cross-sectional study, PKU patients were observed during prepubertal age (p) or after the pubertal spurt (P), assessing body composition and bone quality and correlating these parameters with dietary compliance and types of protein substitutes. Anthropometric and dietary data were evaluated together with bioelectrical impedance analysis (BIA), quantitative ultrasound (QUS) and branched-chain amino acids (BCAAs). A total of 36 patients (16 males, 17 prepubertal and 19 post-pubertal; mean \pm SD age 11.4 ± 3.9 years) were included. A higher BMI was observed in adolescents (p -value: 0.018). The BIA revealed a significant increase in total body water (TBW) and muscle mass (MM) in P subjects either compliant (p -value: 0.001) or non-compliant with the diet (p -value: 0.001). MM content correlated with increased Phe intake ($r = 0.63$; $p < 0.001$). In the subgroup of five patients taking L-AAAs and glycomacropeptides (GMPs), BCAA values tended to be lower than those taking only L-AA mixtures, with a significant trend for valine. Maintenance of body composition parameters within the normal range—for both fat and muscle mass—and levels of BCAAs can be helpful in reducing the risk of becoming overweight in adulthood. Further studies are needed to confirm these findings.

Keywords: PKU; adolescence; bioelectrical impedance analysis; muscle mass; branched-chain amino acids; quantitative ultrasound



Citation: Tummolo, A.; Carella, R.; Paterno, G.; Bartolomeo, N.; Giotta, M.; Dicintio, A.; De Giovanni, D.; Fischetto, R. Body Composition in Adolescent PKU Patients: Beyond Fat Mass. *Children* **2022**, *9*, 1353. <https://doi.org/10.3390/children9091353>

Academic Editor: Stella Stabouli

Received: 21 July 2022

Accepted: 31 August 2022

Published: 4 September 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Phenylketonuria (PKU) is a rare autosomal recessive inborn error of phenylalanine (Phe) metabolism caused by the deficient activity of Phe hydroxylase (PAH), which is needed to convert Phe into tyrosine. When left untreated, patients with PKU develop severe and irreversible neurological damage, growth deficiency, eczema and skin and hair hypopigmentation [1].

The mainstay of treatment involves a Phe-restricted diet, composed of strict control of natural protein intake, administration of protein substitutes (PE) derived from Phe-free amino acids (L-AAAs), large neutral amino acids (LNAA) or low-Phe glycomacropeptide (GMP) [2,3] and the use of special low-protein foods. The primary aim is to maintain blood Phe levels within a therapeutic target range in order to prevent neurological sequelae [1] while ensuring nutritional requirements are met for normal growth development [2,4,5]. To obtain these endpoints, the daily protein intake of individuals with PKU exceeds the WHO/FAO/UNU protein intakes of the general population [6].

L-AAAs are special formulas containing synthetic mixtures of amino acids devoid of Phe and supplemented with tyrosine—the amino acid that is lacking in PKU. A variety of commercial products are available to meet patients' preferences for taste and palatability [3].

GMP is a natural Phe-free protein source derived from whey that contains small amounts of Phe. However—given the low Phe content—GMP has been used for PKU patients, and

many studies have identified no adverse events and have demonstrated its efficacy in maintaining serum levels of Phe—mainly thanks to the palatability of the formulations [3,7].

The impact of a restricted diet regimen on both short- and long-term growth [8] requires ongoing evaluation [9,10]. Although Body Mass Index (BMI) is widely used as the outcome measure of growth, it has clear limitations. The relative contribution of lean and fat mass to body weight cannot be assessed. Therefore, based on BMI, some patients may appear to be relatively weight-stable, despite actually changing in terms of their underlying fat/lean ratio.

Bioelectrical Impedance Analysis (BIA) offers a rapid, simple, noninvasive and cheap way of assessing body composition in children and adolescents [11].

Adolescence, in particular, is a period of growth and maturation, characterized by changes in body composition—mainly in the location and amount of body fat and changes in anabolic effects, with reduced insulin sensitivity [12,13]. In PKU patients, many studies [8,14,15] have emphasized the risk of higher fat mass percentages related to diet content and lifestyle—being particularly evident in females. PKU adolescents may be particularly exposed to body composition deviations because of their diet [16–18] and their higher frequency of diet transgressions [7,19,20]. Some studies have found that PKU subjects with poorer metabolic control due to low compliance to diets are more likely to be overweight than others, but the relative contribution of fat and muscle to their being overweight has not been assessed [16,21,22].

The role on muscle body mass of a high protein diet in PKU patients has been poorly investigated, as well as the effects of the different protein substitutes used to meet their daily protein requirements on their body composition and bone mineralization.

In our study, we aimed at assessing body composition and bone quality in adolescent PKU patients, comparing them with prepubertal PKU patients. We also correlated body composition and bone quality parameters with compliance to diet and type of protein substitutes: L-AAs with or without GMPs.

2. Materials and Methods

2.1. Patients

We performed an observational, cross-sectional study on a cohort of patients affected by PKU. The inclusion criteria were diagnosis by neonatal screening and confirmation by molecular analysis. The study was conducted between May 2017 and January 2022. The exclusion criteria were concomitant chronic disorders altering bone mineralization and/or nutritional conditions and/or timing of puberty (precocious or delayed puberty were excluded). The type of PKU was assigned based on the PKU European Guidelines [1]. Written informed consent was obtained from parents. All procedures were in agreement with the guidelines of the Declaration of Helsinki on Human Experimentation.

Patients were assessed either during prepubertal age (prepubertal group hereafter referred as group “p”) or after the pubertal spurt (pubertal group referred as group “P”). The Tanner stage (prepubertal, Tanner 1; pubertal, Tanner 2 to 5) was used to determine pubertal progression by physical examination [23].

A subgroup of 5 patients was observed during both periods.

Anthropometric parameters were assessed with the same balance and altimeter weighting, measuring patients in light clothing. BMI was calculated as the weight/height² ratio; normal values of the BMI Z-score range between +1.0 and −1.0, according to the World Health Organization (WHO) charts [24].

All patients were treated with a diet regimen restricted in Phe and supplemented with L-AAs, and in five cases, even took GMP supplementation. A trained metabolic dietitian analyzed the dietary data from each patient using Winfood Pro software (version 3.0.0, 2011, Medimatica Srl, Teramo, Italy). Dietary protein intake, expressed in grams per kilograms per day, was extrapolated from FAO/WHO/UNU recommended safe levels [6], with an extra percentage of total protein intake necessary to support growth, differing by age [25,26]. Energy intake was expressed as kcal/Kg/day and, for PKU subjects, was comparable with

the estimated average requirements of the general population [27]. Compliance to the diet was evaluated through the mean Phe levels resulting from the weekly dried blood spot (DBS) tests performed over the previous 6 months; values were directly retrieved from the laboratory database. Target levels were: 120–360 $\mu\text{mol/l}$ for patients aged <12 years and 120–600 $\mu\text{mol/l}$ for those older than 12 years of age, according to the European Guidelines [1]. Dietary compliance was defined as poor if mean DBS Phe levels were higher than the target level or good if mean dried blood spot Phe levels fell within the target Phe level. All our patients took the entire prescribed quantity of synthetic proteins even if they were poorly compliant to the diet. The daily amount of GMP was calculated as 70% of the total protein equivalents (PE), expressed in grams of the product used [28]. Additionally, the percentage of GMPs—expressed as g/kg/day—over total PE intake (g/kg/day) was calculated. Serum samples were assayed for branched-chain amino acids (BCAA) by liquid chromatography (LC)–MS/MS [29].

2.2. Body Composition

BIA measurements were performed by a trained study assistant. A multi-frequency (20 kHz and 100 kHz) BIA device using an eight-point tactile electrode system (Inbody 230, Biospace Corp., Seoul, Korea) referred to as BIA8MF was used. The performance characteristics of the instrument and the procedures used have been already described [30]. Body composition parameters, calculated by the manufacturer's software (Lookin'Body 120, Biospace Corp., Seoul, Korea) were: fat mass (FM; Kg), percentage of body fat (PBF), total body water (TBW; Kg), waist–hip ratio (WHR) and muscle mass (MM; Kg). The machine provided normal ranges of BIA parameters for the general population according to gender and age.

2.3. QUS (Quantitative Ultrasound Scan)

A calcaneus QUS SONOST 3000 (Osteosys, Korea) was used to calculate the bone quality index (BQI) and the BQI SDS. The BQI was derived from the broadband ultrasound attenuation (BUA, measured in dB/MHz) and the speed of sound (SOS, measured in m/s). The BUA reflects bone structure and is directly related to temperature; the SOS reflects bone mineral density and is inversely related to temperature. Correlation coefficients (α , β) combined with BUA and SOS allow one to obtain the BQI ($\text{BQI} = \alpha \times \text{SOS} + \beta \times \text{BUA}$) [31].

2.4. Statistical Analysis

Thirty-six patients were recruited during the observation period (17 prepubertal and 19 pubertal).

For a two-sample pooled *t* test of a normal mean difference with a two-sided significance level of 0.05, assuming a common standard deviation of 0.3 g/kg, group sample sizes of 17 and 19 have a power of 0.82 to detect a mean difference of 0.3 g/kg in total protein intake [30] between the Prepubertal and Pubertal group.

Continuous variables were expressed as the mean and standard deviation (SD) for normally distributed parameters or the median and interquartile range (IQR) in the case of skewed data distribution. Shapiro–Wilk statistics were used to test normality. Differences in continuous variables between groups were compared using a Student's *t*-test for normally distributed parameters or a nonparametric Mann–Whitney U Test otherwise.

Categorical data were expressed as frequency and percentage, and the Chi-square test or Fisher's exact test was used to compare the two groups.

Multivariable generalized linear models were applied to evaluate the effects of puberty and compliance to diet—adjusted for sex—on the BIA and QUS parameters and on levels of BCAA. The results are shown as least square means with their standard errors; pairwise multiple comparisons were adjusted according to the Tukey correction.

All tests of statistical significance were two-tailed, and *p*-values less than 0.05 were considered statistically significant. Statistical analysis was performed using SAS/STAT[®] Statistics, Version 9.4 (2013), SAS Institute Inc., Cary, NC, USA.

3. Results

3.1. Comparison of BIA and QUS Parameters in Different Subgroups

A total of 36 patients (17 p and 19 P, mean \pm SD age 11.4 ± 3.9 years) were included in the study. Of them, five were observed over both periods, for a total of 41 observations. The demographic and dietary characteristics of the entire group and of the two subgroups are shown in Table 1. Nine out of thirty-six patients (25%) were considered non-compliant with the diet, based on the mean of the Phe values of the previous six months: 18% in the p group and 32% in the P group (p -value: 0.451).

Table 1. Demographic and dietary characteristics of the patients.

	Total (n = 36)	p Group (n = 17)	P Group (n = 19)	p -Value
Males (%)	16 (44%)	6 (35%)	10 (53%)	0.296
Age (years)	11.4 ± 3.9	8.1 ± 2.2	14.4 ± 2.4	<0.001
Phe restriction + LAA mixtures + GMP	5 (14%)	3 (18%)	2 (11%)	0.650
Total protein intake (g/Kg/day)	1.24 ± 0.30	1.39 ± 0.31	1.10 ± 0.21	0.002
Natural protein intake (g/kg/day)	0.39 [0.22–0.75]	0.41 [0.29–0.79]	0.32 [0.19–0.70]	0.288
PE intake (g/kg/day)	0.77 ± 0.32	0.86 ± 0.28	0.68 ± 0.34	0.094
GMPs intake (g/day) *	12.4 ± 2.2	12.1 ± 2.9	13 ± 1.4	0.711
GMPs (g/kg/day)/total PE (g/kg/day) (%) *	57 ± 32.3	54 ± 13.3	61 ± 34.9	0.473
Total caloric intake (Kcal/Kg/day)	47.1 ± 17.1	58.7 ± 17.2	36.8 ± 8.0	<0.001
Phe intake (mg/kg/day)	13.4 [7.6–20.6]	16.9 [10.76–21]	9 [5.4–17.7]	0.034
Phe levels	389.7 ± 199	313.5 ± 106.8	457.9 ± 237.5	0.013
Patients non-compliant with diet (%)	9/36 (25%)	3/17 (18%)	6/19 (32%)	0.451

Data expressed as mean \pm standard deviation or median [interquartile range]. * 5 observations (3 p and 2 P).

Regardless of dietary compliance, considering the whole group of patients, a significant increase in BMI was observed in the P group compared to p (18.06 ± 2.8 vs. 22.47 ± 3.78 (p -value: <0.001)). However, in adolescents, the PBF was significantly reduced (28.46 ± 14.62 vs. 19.17 ± 12.93 , p -value: 0.049), whereas the BQI was significantly increased (53.73 ± 7.27 vs. 71.5 ± 17.49 , p -value: 0.033; data not shown).

We performed a comparison of BIA and QUS parameters between males and females, regardless of their age (Table S1); no statistically significant difference was found in any parameters. Considering dietary compliance, a multivariate analysis adjusted for sex showed a significant increase in BMI between the prepubertal and post-pubertal group (p -value: 0.018; Table 2, a vs. c). No statistically significant differences in the amount and percentage of fat mass was observed in any of the subgroups analyzed (Table 2). The same was observed for the BQI and SDS-BQI. Within the prepubertal group, being compliant or non-compliant to the diet did not produce statistically significant differences in the QUS and BIA parameters (Table 2, a vs. b). On the other hand, in the P group, following or not following the diet resulted in a statistically significant difference in TBW (p -value: 0.001) and MM (p -value: 0.001; Table 2, c vs. d). These values are also higher than the upper limit of the mean normal values for prepubertal and post-pubertal children (Table 2).

Table 2. Parameter averages estimated in the pre/post-pubertal and compliant/non-compliant groups through multivariate analysis, adjusting for sex.

Parameter	p-Group Compliant (a)	p-Group Non-Compliant (b)	P-Group Compliant (c)	P-Group Non-Compliant (d)	(a) vs. (b)	(c) vs. (d)	(a) vs. (c)	(b) vs. (d)
	LS. Means ± Standard Error				Adjusted <i>p</i> -Value *			
BMI	17.4 ± 0.9	19.3 ± 1.9	21.3 ± 0.8	25.2 ± 1.2	0.818	0.067	0.018	0.059
sds-BMI	−0.1 ± 0.3	0.7 ± 0.7	0.2 ± 0.3	0.8 ± 0.4	0.761	0.652	0.911	0.997
FM n.v. 5.51–9.51 <i>n.v. 10.01–20.07</i>	7.5 ± 2.4	12.9 ± 4.6	12.2 ± 2.2	11.9 ± 3.4	0.765	1.000	0.492	0.998
PBF% n.v. 11.75–21.75 <i>n.v. 13.89–23.89</i>	27.2 ± 4.4	28.9 ± 8.5	21.8 ± 4	13.5 ± 6.2	0.998	0.687	0.798	0.463
TBW n.v.14.41–17.63 <i>n.v. 35.38–43.26</i>	15 ± 2	18.3 ± 3.7	30.9 ± 1.8	45.6 ± 2.7	0.881	0.001	<0.001	<0.001
WHR n.v.0.76–0.86 <i>n.v. 0.78–0.88</i>	0.74 ± 0.02	0.8 ± 0.05	0.85 ± 0.02	0.83 ± 0.03	0.710	0.930	0.010	0.957
MM n.v. 9.71–11.87 <i>n.v. 26.77–32.78</i>	10 ± 1.6	12.8 ± 3.1	23 ± 1.4	35.3 ± 2.3	0.873	0.001	<0.001	<0.001
sds-BQI	−1.6 ± 0.4	−1.9 ± 0.9	−1.8 ± 0.3	−0.9 ± 0.5	0.984	0.496	0.978	0.755
BQI	54.8 ± 4.3	50.1 ± 10.6	69.6 ± 4	75.4 ± 5.9	0.978	0.850	0.081	0.182
Leucine	135 ± 11	106 ± 27	141 ± 10	147 ± 15	0.770	0.991	0.980	0.573
Isoleucine	69.9 ± 4.7	54.6 ± 12	76.6 ± 4.7	65.9 ± 6.7	0.667	0.570	0.755	0.847
Valine	227 ± 12	205 ± 30	243 ± 12	231 ± 17	0.908	0.940	0.795	0.874

* Adjusted for multiple comparison by Tukey. Block letters: range of normal values for prepubertal children. Italics: range of normal values for post-pubertal children.

Concerning those compliant to the diet, the comparison between the two groups showed an increase in TBW (*p*-value: <0.001), WHR (*p*-value: 0.01) and MM (*p*-value: <0.001) in the P group (Table 2, a vs. c).

Similarly, among those non-compliant to the diet, TBW and MM were significantly lower in the p than in the P group (*p*-value: <0.001), highlighting the change in the amount of water and muscle tissue after puberty, regardless to the adherence to diet (Table 2, b vs. d). No difference among the subgroups for BCAA values were found. For all parameters, the “sex” adjustment factor was not significant.

3.2. Comparison of BIA and QUS Parameters before and after the Pubertal Spurt

An analysis of five patients before and after the pubertal spurt showed an increase in BMI in all cases. Within the BIA parameters, FM remained almost stable in Patient 1 and Patient 4, while an increase in FM occurred in the others. The PBF presented a variable trend. The parameters that increased in all cases were TBW and MM, which were found to always be higher after puberty. The WHR was also higher in all patients in the post-pubertal period. For QUS parameters, SDS-BQI remained almost stable in two cases (Patients 4 and 5), but decreased in the remaining three patients (Table 3).

Table 3. Parameters of the five patients analyzed before and after puberty.

General Characteristics								BIA Parameters								QUS Parameters									
Pts	Sex	Age	Tanner Stage	Phe Values (Mean SD)	BMI	SDS BMI	FM	PBF	TBW	WHR	MM	BQI	SDS-BQI												
1	F	9.3	11.5	I	III	217	148	22.15	26.21	1.31	1.62	20.70	21.80	53.70%	40.60%	13.20	23.50	0.75	0.87	8.60	16.80	54.26	75	0.90	-0.30
2	F	10.7	12.9	I	III	312	368	15.67	19.34	-1.22	-0.31	4.8	9.2	15.30%	20.9%	19.7	25.7	0.79	0.83	13.9	18.7	61.39	66.49	-1.70	-2.90
3	M	9.1	12.8	I	III	364	708	21.40	27.60	1.23	1.74	21.90	29.10	42.30%	42.80%	21.90	28.50	0.86	0.98	15.70	21.10	61.97	59.15	-1.00	-2.20
4	F	10.7	12.9	I	II	355	645	21.23	22.60	0.76	0.68	15.10	15.60	35.80%	29.8%	19.9	26.9	0.79	0.82	14	19.60	71.40	81	-1.40	-1.30
5	M	12.1	14.8	I	II	350	487	18.43	23.4	-0.39	0.66	13.5	19	39.10%	36.50%	15.5	24.3	0.76	0.85	10.6	17.5	52	65	-2.4	-2.3

Grey columns: prepubertal period. White columns: post-pubertal period.

3.3. Comparison of BIA and QUS Parameters According to Type of Protein Substitute

An evaluation of BIA and QUS parameters and BCAA levels in relation to the type of synthetic protein consumed—L-AAs alone (31 patients) or with GMP (five patients)—showed no significant differences in BIA and QUS parameters, although a trend toward a higher BQI and a lower FM and MM was observed in the second subgroup. The group of patients taking GMP with L-AAs tended to also have lower branched-chain amino acid values, with statistical significance for valine (*p*-value: 0.01; Figure 1).




Figure 1. Comparison of parameters according to type of protein substitute.

3.4. Correlation of BIA and QUS Parameters with Protein Intake and Compliance to Diet

The only significant correlations (p -value < 0.05) identified were: TBW and MM with dietary Phe intake (respectively r : 0.47; 0.46) and WHR with increasing mean Phe values (r : 0.43). No significant correlations with the other BIA parameters or with the BQI were retrieved (data not shown). The increase in MM correlated significantly with average Phe values (Figure 2).




Figure 2. Correlation between mean plasma Phe levels and muscle mass (MM). $r = 0.63$ $p < 0.001$; Analysis only possible for 32 out of 36 patients because of missing BIA values.

4. Discussion

Our study showed a higher BMI in post-pubertal than prepubertal PKU patients, due to significant increases in TBW and MM in both compliant and non-compliant to diet subjects—in this last case, also exceeding to the normal range for the general population. Muscle mass content correlated with increased plasma Phe levels and therefore with diet compliance.

In the subgroup of patients taking L-AAs and GMPs, we found no statistically significant differences in BIA and QUS values compared to the group taking only L-AA mixtures; however, the branched-chain amino acid values tended to be lower in the GMP group than in those who took only L-AA mixtures, with a significant trend for valine.

Body composition changes in the human body with age and is also gender-specific [32]. This is secondary to reproductive and endocrine changes in the body [33]. During puberty, hormonal fluctuations as well as a rapid growth in body size are accompanied by marked changes in body composition [34]. This situation often correlates with the development of excessive weight and obesity in children [35], although even the MM increases after puberty [36]—but this is generally not considered a harbinger of later-life metabolic disturbances.

In our cohort of patients, we did not find an increase in FM after puberty—in contrast with what has been reported by other studies [13–15], with no differences between males and females.

It can be hypothesized that the influence of diet on the body composition of these subjects overtakes the effects of sexual hormones that are responsible for the different changes in body composition between the two sexes after puberty [37–40].

In our sample, the increased BMI in post-pubertal patients was sustained by a higher muscle mass and improved bone quality index. In this context, BIA provided a more accurate assessment than BMI on the muscle/fat ratio.

This increase is explained by studies reporting greater protein retention at this age. After puberty, selective stimulation of whole-body protein synthesis by GH, IGF-1 and androgenic hormones may explain the higher protein gain during this period [41].

Significant evidence from large, high-quality randomized control trials demonstrates that the higher protein content of formula milk, more than breast milk, is associated with adverse infant and child outcomes [42,43]. The role of protein intake on adiposity beyond the weaning period has also been investigated [37–39]; Gunther [44] and Assmann [45] reported that higher protein intake exceeding physiologic requirements, particularly between the age of 9 and 15 years, is associated with higher fat-free mass (FFM). The growth hormone/insulin-like factor 1 axis may be stimulated by excessive protein intake and drive the early differentiation and proliferation of adipocytes [46].

PKU patients can be considered a vulnerable group for the above metabolic abnormalities. In fact, to ensure normal growth, PKU children are prescribed a total protein intake that exceeds the WHO/FAO/UNU 2007 safe levels of proteins [1].

Special attention should be paid to older children and adolescents, who have autonomy in dietary choices and are often inclined to dietary transgressions, which in case of normal protein substitute intake, is associated with an even higher daily protein intake.

GH, IGF-1 and sex steroids all markedly increase during puberty and their actions are amplified mutually as they control not only increases in muscle mass but also affect the mineralization of the skeleton. The synergistic actions of these anabolic hormones appear to be most significant during the finite years of puberty [47,48].

An increase in bone density during the adolescent period has also been reported, with both genders reaching peak bone gain after puberty [49]. This higher protein retention may explain the increasing bone quality observed by us in the present and in a previous study by our group [31], which has also been confirmed by other authors [50].

In classical PKU, the protein substitutes L-AAs and/or GMP provide up to 80% of dietary protein requirements and are essential to ensure metabolic stability and growth. Protein substitutes, while meeting the protein requirements for cellular function and growth, have several pharmacological and physiological functions [51,52]. The amino acid profile is different for each protein substitute, with variations in amino acid patterns as well as in the amount of essential and non-essential amino acids per 100 g of PE. Additionally, their relative absorption rates and bioavailability are not fully understood [53].

In currently available GMP formulas, the amount of Leucine on average exceeds that of the most common L-AAs formulas (196 vs. 121 mg/g PE), while the amount of Isoleucine is comparable (71 vs. 77 mg/g PE) and there is a lower amount of Valine (60 vs. 89 mg/g PE). Modifications in GMP composition, however, do not determine statistically significant differences in BCAA plasma levels compared to the average values of BCAAs in patients taking L-AAs—except for a decrease in Val levels [54], as confirmed by our study. However, this seems to be a beneficial effect of GMPs, since recent studies associate high levels of Valine with increased oxidative stress and the onset of Type 2 Diabetes [55].

In this context, other studies have reported that elevated levels of plasma BCAAs may be associated with obesity [56] and insulin resistance [57–59]. However, the relationship between BCAA and detailed parameters of body composition has yet to be reported.

Acute increases in plasma amino acids, linked to the absorption profile of L-AAs, worsen insulin sensitivity. L-AAs are unable to replicate the physiological actions of

whole proteins, being directly absorbed from the small intestine [60]. Substantial evidence suggests that the ingestion of large doses of L-AAs increases amino acid oxidation and nitrogen excretion, decreasing their availability for cellular functioning—whereas GMP lowers the rate of amino acid absorption and improves nitrogen retention [61].

Ney et al. [62] and van Calcar et al. [61] suggested that GMP may induce a slower and more sustained release of amino acids, leading to a more physiological availability of amino acids, leading to improved growth and body composition.

There have been few studies investigating changes in body composition in PKU patients taking L-AA and/or GMP. A 3-year prospective study by Daly et al. [21] compared the effect of GMP vs. L-AAs on body composition and growth in 48 PKU children between 5 and 16 years of age. Results showed that children in the GMP group had improved lean body mass (LBM), with decreased fat mass and % body fat—possibly secondary to the high content of BCAA in GMP. A higher LBM content may be supposed; however, this study is not based on the new GMP compositions and does not report information about MM content and the measurement of plasma BCAA.

In another study, Huemer et al. [63] measured growth and body composition over 12 months in 34 children with classical PKU. Total protein intake was 124% of the German recommended daily allowance. A significant correlation was found between LBM and natural protein intake, in keeping with our results—even though the possible implications of a particularly high protein intake in these patients were not discussed.

Limitations of our study include the low number of patients in the GMP group compared with those also administered with L-AAs and the lack of insulin dose as a marker to correlate with BCAA levels. On the other hand, this study has some strengths: The inclusion and exclusion criteria for patient selection and the inclusion of independent variables (covariates) in a multiple regression model [64] reduced the selection bias problems typical of cross-sectional observational studies. This is the first study highlighting the importance of assessing muscle mass content, beyond fat mass content, in helping assessing nutritional risks—particularly during adolescence and under non-compliance conditions. Furthermore, this represents the first evidence for a potential role of GMPs in keeping plasma BCAA levels lower, thanks to their more physiological amino acid absorption profile.

5. Conclusions

In conclusion, BIA, performed regularly, provides more complete elements than BMI for the evaluation of the nutritional status of PKU patients—especially in adolescence. In this period, MM increased compared to FM—especially in subjects with a low compliance to the diet. Maintenance within the normal range of parameters of body composition, for both fat and muscle mass, and levels of branched-chain amino acids can be helpful to reduce the risk of becoming overweight in adulthood. Further studies involving a greater number of patients and a correlation between total protein intake, use of different protein substitutes in PKU diets and the risk of metabolic syndrome in adult life are needed to support these results.

Supplementary Materials: The following supporting information can be downloaded at: <https://www.mdpi.com/article/10.3390/children9091353/s1>, Table S1: Comparison of BIA and QUS parameters according with gender.

Author Contributions: A.T. interpreted data and wrote the manuscript; G.P. and R.C. made contributions to study sample follow-up; N.B. and M.G. performed statistical analyses and interpreted data; D.D.G. and A.D. contributed to data collection; R.F. critically revised the manuscript. All authors have read and agreed to the published version of the manuscript.

Funding: Authors declare no sources of study funding including sponsorship for the described work.

Institutional Review Board Statement: The study was conducted according to the guidelines of the “Declaration of Helsinki” and did not require approval by the Local Ethics Committee.

Informed Consent Statement: Written informed consent was obtained from the participants of the study/their parents.

Data Availability Statement: The data presented in this study are available upon request from the corresponding author.

Conflicts of Interest: No financial benefits have been received or will be received from any party related directly or indirectly to the subject of this article.

References

1. Van Wegberg, A.M.J.; Macdonald, A.; Ahring, K.; BéLanger-Quintana, A.; Blau, N.; Bosch, A.M.; Burlina, A.; Campistol, J.; Feillet, F.; Gizewska, M.; et al. The complete European guidelines on phenylketonuria: Diagnosis and treatment. *Orphanet J. Rare Dis* **2017**, *12*, 162. [[CrossRef](#)] [[PubMed](#)]
2. Rodrigues, C.; Pinto, A.; Faria, A.; Teixeira, D.; van Wegberg, A.M.J.; Ahring, K.; Feillet, F.; Calhau, C.; MacDonald, A.; Moreira-Rosário, A.; et al. Is the Phenylalanine-Restricted Diet a Risk Factor for Overweight or Obesity in Patients with Phenylketonuria (PKU)? A Systematic Review and Meta-Analysis. *Nutrients* **2021**, *13*, 3443. [[CrossRef](#)] [[PubMed](#)]
3. Boyer, S.W.; Barclay, L.J.; Burrage, L.C. Inherited Metabolic Disorders: Aspects of Chronic Nutrition Management. *Nutr. Clin. Pract.* **2015**, *30*, 502–510. [[CrossRef](#)] [[PubMed](#)]
4. Schiergens, K.A.; Weiss, K.J.; Dokoupil, K.; Fleissner, S.; Maier, E.M. Dietary treatment of inborn errors of metabolism—a balancing act between indulgence and therapy. *Bundesgesundheitsblatt Gesundheitsforschung* **2020**, *63*, 864–871. [[CrossRef](#)]
5. Camp, K.M.; Lloyd-Puryear, M.A.; Huntington, K.L. Nutritional treatment for inborn errors of metabolism: Indications, regulations, and availability of medical foods and dietary supplements using phenylketonuria as an example. *Mol. Genet. Metab.* **2012**, *107*, 3–9. [[CrossRef](#)]
6. World Health Organization. Protein and amino acid requirements in human nutrition: Report of a joint FAO/WHO/UNU expert consultation. In *Joint FAO/WHO/UNU Expert Consultation on Protein and Amino Acid Requirements in Human Nutrition*; Food and Agriculture Organization of the United Nations, World Health Organization & United Nations University: Geneva, Switzerland, 2007.
7. Pena, M.J.; Pinto, A.; Daly, A.; MacDonald, A.; Azevedo, L.; Rocha, J.C.; Borges, N. The Use of Glycomacropeptide in Patients with Phenylketonuria: A Systematic Review and Meta-Analysis. *Nutrients* **2018**, *10*, 1794. [[CrossRef](#)]
8. Evans, M.; Truby, H.; Boneh, A. The Relationship between Dietary Intake, Growth, and Body Composition in Inborn Errors of Intermediary Protein Metabolism. *J. Pediatr.* **2017**, *188*, 163–172. [[CrossRef](#)]
9. Gambello, M.J.; Li, H. Current strategies for the treatment of inborn errors of metabolism. *J. Genet. Genom.* **2018**, *45*, 61–70. [[CrossRef](#)]
10. Langeveld, M.; Hollak, C.E.M. Bone health in patients with inborn errors of metabolism. *Rev. Endocr. Metab. Disord.* **2018**, *19*, 81–92. [[CrossRef](#)]
11. López-Sánchez, G.F.; Sgroi, M.; D’Ottavio, S.; Díaz-Suárez, A.; González-Villora, S.; Veronese, N.; Smith, L. Body Composition in Children and Adolescents Residing in Southern Europe: Prevalence of Overweight and Obesity According to Different International References. *Front. Physiol.* **2019**, *10*, 130. [[CrossRef](#)]
12. Stefanaki, C.; Bacopoulou, F.; Peppas, M. Prediabetes and adolescence—Trends, causes, effects, and screening. *US Endocrinol.* **2016**, *12*, 94–98. [[CrossRef](#)]
13. Obesity and Overweight—WHO. Available online: <https://www.who.int/news-room/fact-sheets/detail/obesity-and-overweight> (accessed on 9 June 2021).
14. Robertson, L.V.; McStravick, N.; Ripley, S.; Weetch, E.; Donald, S.; Adam, S.; Micciche, A.; Boocock, S.; MacDonald, A. Body mass index in adult patients with diet-treated phenylketonuria. *J. Hum. Nutr. Diet.* **2013**, *26*, 1–6. [[CrossRef](#)] [[PubMed](#)]
15. Burrage, L.C.; McConnell, J.; Haesler, R.; O’Riordan, M.A.; Sutton, V.R.; Kerr, D.S.; McCandless, S.E. High prevalence of overweight and obesity in females with phenylketonuria. *Mol. Genet. Metab.* **2012**, *107*, 43–48. [[CrossRef](#)] [[PubMed](#)]
16. Mazzola, P.N.; Nalin, T.; Castro, K.; van Rijn, M.; Derks, T.G.J.; Perry, I.D.S.; Mainieri, A.S.; Schwartz, I.V.D. Analysis of body composition and nutritional status in Brazilian phenylketonuria patients. *Mol. Genet. Metab. Rep.* **2016**, *6*, 16–20. [[CrossRef](#)]
17. Ng, M.; Fleming, T.; Robinson, M.; Thomson, B.; Graetz, N.; Margono, C.; Mullany, E.C.; Biryukov, S.; Abbafati, C.; Abera, S.F.; et al. Global, regional, and national prevalence of overweight and obesity in children and adults during 1980–2013: A systematic analysis for the Global Burden of Disease Study 2013. *Lancet* **2014**, *384*, 766–781. [[CrossRef](#)]
18. Walter, J.H.; White, F.J.; Hall, S.K.; MacDonald, A.; Rylance, G.; Boneh, A.; Francis, D.E.; Shortland, G.J.; Schmidt, M.; Vail, A. How practical are recommendations for dietary control in phenylketonuria? *Lancet* **2002**, *360*, 55–57. [[CrossRef](#)]
19. Walter, J.H.; White, F.J. Blood phenylalanine control in adolescents with phenylketonuria. *Int. J. Adolesc. Med. Health* **2004**, *16*, 41–45. [[CrossRef](#)]
20. Crone, M.R.; van Spronsen, F.J.; Oudshoorn, K.; Bekhof, J.; van Rijn, G.; Verkerk, P.H. Behavioural factors related to metabolic control in patients with phenylketonuria. *J. Inher. Metab. Dis.* **2005**, *28*, 627–637. [[CrossRef](#)]
21. Daly, A.; Högl, W.; Crabtree, N.; Shaw, N.; Evans, S.; Pinto, A.; Jackson, R.; Strauss, B.; Wilcox, G.; Rocha, J.; et al. Growth and Body Composition in PKU Children—A Three-Year Prospective Study Comparing the Effects of L-Amino Acid to Glycomacropeptide Protein Substitutes. *Nutrients* **2021**, *13*, 1323. [[CrossRef](#)]

22. Camatta, G.C.; Kanufre, V.D.C.; Alves, M.R.A.; Soares, R.D.L.; Norton, R.D.C.; De Aguiar, M.J.B.; Starling, A.L.P. Body fat percentage in adolescents with phenylketonuria and associated factors. *Mol. Genet. Metab. Rep.* **2020**, *23*, 100595. [[CrossRef](#)]
23. Tanner, J.M.; Whitehouse, R.H. Clinical longitudinal standards for height, weight, height velocity, weight velocity, and stages of puberty. *Arch. Dis. Child.* **1976**, *51*, 170–179. [[CrossRef](#)] [[PubMed](#)]
24. BMI-for-Age (5–19 Years)–WHO. Available online: <https://www.who.int/tools/growth-reference-data-for-5to19-years/indicators/bmi-for-age> (accessed on 20 July 2021).
25. Vockley, J.; Andersson, H.C.; Antshel, K.M.; Braverman, N.E.; Burton, B.K.; Frazier, D.M.; Mitchell, J.; Smith, W.E.; Thompson, B.H.; Berry, S.A.; et al. Phenylalanine hydroxylase deficiency: Diagnosis and management guideline. *Genet. Med.* **2014**, *16*, 188–200. [[CrossRef](#)] [[PubMed](#)]
26. Singh, R.H.; Rohr, F.; Frazier, D.; Cunningham, A.; Mofidi, S.; Ogata, B.; Splett, P.L.; Moseley, K.; Huntington, K.; Acosta, P.B.; et al. Recommendations for the nutrition management of phenylalanine hydroxylase deficiency. *Genet. Med.* **2014**, *16*, 121–131. [[CrossRef](#)] [[PubMed](#)]
27. Shaw, V. *Clinical Paediatric Dietetics-Fifth Edition*; Wiley Blackwell: Chichester, UK; Hoboken, NJ, USA, 2020.
28. Stroup, B.M.; Sawin, E.A.; Murali, S.G.; Binkley, N.; Hansen, K.E.; Ney, D.M. Amino Acid Medical Foods Provide a High Dietary Acid Load and Increase Urinary Excretion of Renal Net Acid, Calcium, and Magnesium Compared with Glycomacropeptide Medical Foods in Phenylketonuria. *J. Nutr. Metab.* **2017**, *2017*, 1909101. [[CrossRef](#)]
29. Millington, D.S. Tandem Mass Spectrometry in Clinical Diagnosis. In *Physician's Guide to the Laboratory Diagnosis of Metabolic Diseases*; Blau, N., Duran, M., Blaskovics, M.E., Gibson, K.M., Eds.; Springer: Berlin/Heidelberg, Germany, 2003.
30. Tummolo, A.; Arico, M.; Pesce, S.; Fantasia, M.L.; Paterno, G.; Melpignano, L.; Giovanni, D.D.; Faienza, M.F. Body Composition and Bone Mineral Quality in Phenylketonuria: Influence of Pubertal Development. *J. Endocrinol. Disord.* **2021**, *7*, 1045.
31. Manuals and User Guides for OsteoSys SONOST 3000. Available online: <https://www.manualslib.com/products/Osteosys-Sonost-3000-8710830.html> (accessed on 6 July 2017).
32. Karastergiou, K.; Smith, S.R.; Greenberg, A.S.; Fried, S.K. Sex differences in human adipose tissues—The biology of pear shape. *Biol. Sex Differ.* **2012**, *3*, 13. [[CrossRef](#)]
33. Ouellet, V.; Labbé, S.M.; Blondin, D.P.; Phoenix, S.; Guérin, B.; Haman, F.; Turcotte, E.E.; Richard, D.; Carpentier, A.C. Brown adipose tissue oxidative metabolism contributes to energy expenditure during acute cold exposure in humans. *J. Clin. Investig.* **2012**, *122*, 545–552. [[CrossRef](#)]
34. Ravussin, E.; Galgani, J.E. The implication of brown adipose tissue for humans. *Annu. Rev. Nutr.* **2011**, *31*, 33–47. [[CrossRef](#)]
35. Seidell, J.C.; Oosterlee, A.; Deurenberg, P.; Hautvast, J.G.; Ruijs, J.H. Abdominal fat depots measured with computed tomography: Effects of degree of obesity, sex, and age. *Eur. J. Clin. Nutr.* **1988**, *42*, 805–815.
36. Brown, D.C.; Kelnar, C.J.; Wu, F.C. Energy metabolism during male human puberty. I. Changes in energy expenditure during the onset of puberty in boys. *Ann. Hum. Biol.* **1996**, *23*, 273–279. [[CrossRef](#)]
37. Loomba-Albrecht, L.A.; Styne, D.M. Effect of puberty on body composition. *Curr. Opin. Endocrinol. Diabetes Obes.* **2009**, *16*, 10–15. [[CrossRef](#)] [[PubMed](#)]
38. Horlick, M.B.; Rosenbaum, M.; Nicholson, M.; Levine, L.S.; Fedun, B.; Wang, J.; Pierson, R.N., Jr.; Leibel, R.L. Effect of puberty on the relationship between circulating leptin and body composition. *J. Clin. Endocrinol. Metab.* **2000**, *85*, 2509–2518. [[CrossRef](#)]
39. Ausili, E.; Rigante, D.; Salvaggio, E.; Focarelli, B.; Rendeli, C.; Ansuini, V.; Paolucci, V.; Triarico, S.; Martini, L.; Caradonna, P. Determinants of bone mineral density, bone mineral content, and body composition in a cohort of healthy children: Influence of sex, age, puberty, and physical activity. *Rheumatol. Int.* **2012**, *32*, 2737–2743. [[CrossRef](#)] [[PubMed](#)]
40. Garnett, S.P.; Högl, W.; Blades, B.; Baur, L.A.; Peat, J.; Lee, J.; Cowell, C.T. Relation between hormones and body composition, including bone, in prepubertal children. *Am. J. Clin. Nutr.* **2004**, *80*, 966–972. [[CrossRef](#)]
41. Mauras, N.; Rini, A.; Welch, S.; Sager, B.; Murphy, S.P. Synergistic effects of testosterone and growth hormone on protein metabolism and body composition in prepubertal boys. *Metabolism* **2003**, *52*, 964–969. [[CrossRef](#)]
42. Koletzko, B.; Broekaert, I.; Demmelmair, H.; Franke, J.; Hannibal, I.; Oberle, D.; Schiess, S.; Baumann, B.T.; Verwied-Jorky, S. Protein intake in the first year of life: A risk factor for later obesity? The E.U. Childhood Obesity Project. *Adv. Exp. Med. Biol.* **2005**, *569*, 69–79.
43. Koletzko, B.; von Kries, R.; Closa, R.; Escribano, J.; Scaglioni, S.; Giovannini, M.; Beyer, J.; Demmelmair, H.; Gruszfeld, D.; Dobrzanska, A.; et al. Lower protein in infant formula is associated with lower weight up to age 2 y: A randomized clinical trial. *Am. J. Clin. Nutr.* **2009**, *89*, 1836–1845.
44. Günther, A.L.; Buyken, A.E.; Kroke, A. Protein intake during the period of complementary feeding and early childhood and the association with body mass index and percentage body fat at 7 y of age. *Am. J. Clin. Nutr.* **2007**, *85*, 1626–1633. [[CrossRef](#)]
45. Assmann, K.E.; Joslowski, G.; Buyken, A.E.; Cheng, G.; Remer, T.; Kroke, A.; Günther, A.L. Prospective association of protein intake during puberty with body composition in young adulthood. *Obesity* **2013**, *21*, E782–E789. [[CrossRef](#)]
46. Rolland-Cachera, M.F.; Deheeger, M.; Akrouf, M.; Bellisle, F. Influence of macronutrients on adiposity development: A follow up study of nutrition and growth from 10 months to 8 years of age. *Int. J. Obes. Relat. Metab. Disord.* **1995**, *19*, 573–578.
47. van Vught, A.J.; Heitmann, B.L.; Nieuwenhuizen, A.G.; Veldhorst, M.A.; Brummer, R.J.; Westerterp-Plantenga, M.S. Association between dietary protein and change in body composition among children (EYHS). *Clin. Nutr.* **2009**, *28*, 684–688. [[CrossRef](#)] [[PubMed](#)]

48. Mauras, N.; Rogol, A.D.; Haymond, M.W.; Veldhuis, J.D. Sex steroids, growth hormone, insulin-like growth factor-1: Neuroendocrine and metabolic regulation in puberty. *Horm. Res.* **1996**, *45*, 74–80. [[CrossRef](#)] [[PubMed](#)]
49. Zulfarina, M.S.; Sharkawi, A.M.; Aqilah-S.N, Z.-S.; Mokhtar, S.-A.; Nazrun, S.A.; Naina-Mohamed, I. Influence of Adolescents' Physical Activity on Bone Mineral Acquisition: A Systematic Review Article. *Iran. J. Public Health* **2016**, *45*, 1545–1557.
50. Mangano, K.M.; Sahni, S.; Kerstetter, J.E. Dietary protein is beneficial to bone health under conditions of adequate calcium intake: An update on clinical research. *Curr. Opin. Clin. Nutr. Metab. Care* **2014**, *17*, 69–74. [[CrossRef](#)] [[PubMed](#)]
51. MacDonald, A.; Chakrapani, A.; Hendriksz, C.; Daly, A.; Davies, P.; Asplin, D.; Hall, K.; Booth, I.W. Protein substitute dosage in PKU: How much do young patients need? *Arch. Dis. Child.* **2006**, *91*, 588–593. [[CrossRef](#)] [[PubMed](#)]
52. MacDonald, A.; Rylance, G.; Davies, P.; Asplin, D.; Hall, S.K.; Booth, I.W. Administration of protein substitute and quality of control in phenylketonuria: A randomized study. *J. Inherit. Metab. Dis.* **2003**, *26*, 319–326. [[CrossRef](#)] [[PubMed](#)]
53. Macleod, E.L.; Ney, D.M. Nutritional Management of Phenylketonuria. *Ann. Nestle Eng.* **2010**, *68*, 58–69. [[CrossRef](#)]
54. Ney, D.M.; Stroup, B.M.; Clayton, M.K.; Murali, S.G.; Rice, G.M.; Rohr, F.; Levy, H.L. Glycomacropeptide for nutritional management of phenylketonuria: A randomized, controlled, crossover trial. *Am. J. Clin. Nutr.* **2016**, *104*, 334–345. [[CrossRef](#)]
55. Hu, W.; Yang, P.; Fu, Z.; Wang, Y.; Zhou, Y.; Ye, Z.; Gong, Y.; Huang, A.; Sun, L.; Zhao, Y.; et al. High L-Valine Concentrations Associate with Increased Oxidative Stress and Newly-Diagnosed Type 2 Diabetes Mellitus: A Cross-Sectional Study. *Diabetes Metab. Syndr. Obes.* **2022**, *15*, 499–509. [[CrossRef](#)]
56. Felig, P.; Marliss, E.; Cahill, G.F. Plasma amino acid levels and insulin secretion in obesity. *N. Engl. J. Med.* **1969**, *281*, 811–816. [[CrossRef](#)]
57. Adams, S.H. Emerging perspectives on essential amino acid metabolism in obesity and the insulin-resistant state. *Adv. Nutr.* **2011**, *2*, 445–456. [[CrossRef](#)] [[PubMed](#)]
58. Wang, T.J.; Larson, M.G.; Vasan, R.S.; Cheng, S.; Rhee, E.P.; McCabe, E.; Lewis, G.D.; Fox, C.S.; Jacques, P.F.; Fernandez, C.; et al. Metabolite profiles and the risk of developing diabetes. *Nat. Med.* **2011**, *17*, 448–453. [[CrossRef](#)] [[PubMed](#)]
59. Krebs, M.; Krssak, M.; Bernroider, E.; Anderwald, C.; Brehm, A.; Meyerspeer, M.; Nowotny, P.; Roth, E.; Waldhäusl, W.; Roden, M. Mechanism of amino acid-induced skeletal muscle insulin resistance in humans. *Diabetes* **2002**, *51*, 599–605. [[CrossRef](#)]
60. Wu, G. Amino acids: Metabolism, functions, and nutrition. *Amino Acids* **2009**, *37*, 1–17. [[CrossRef](#)]
61. Van Calcar, S.C.; MacLeod, E.L.; Gleason, S.T.; Etzel, M.R.; Clayton, M.K.; Wolff, J.A.; Ney, D.M. Improved nutritional management of phenylketonuria by using a diet containing glycomacropeptide compared with amino acids. *Am. J. Clin. Nutr.* **2009**, *89*, 1068–1077. [[CrossRef](#)]
62. Ney, D.M.; Etzel, M.R. Designing medical foods for inherited metabolic disorders: Why intact protein is superior to amino acids. *Curr. Opin. Biotechnol.* **2017**, *44*, 39–45. [[CrossRef](#)] [[PubMed](#)]
63. Huemer, M.; Huemer, C.; Möslinger, D.; Huter, D.; Stöckler-Ipsiroglu, S. Growth and body composition in children with classical phenylketonuria: Results in 34 patients and review of the literature. *J. Inherit. Metab. Dis.* **2007**, *30*, 694–699. [[CrossRef](#)]
64. Trutschel, D.; Palm, R.; Holle, B.; Simon, M. Methodological approaches in analysing observational data: A practical example on how to address clustering and selection bias. *Int. J. Nurs. Stud.* **2017**, *76*, 36–44. [[CrossRef](#)]