

Mycorrhization and chemical seed priming boost tomato stress tolerance by shifts of primary and defence metabolic pathways

Luca Giovannini^{1^}, Chiara Pagliarani^{1^}, Eva Cañizares², Fabiano Sillo¹, Walter Chitarra³, Silvia De Rose¹, Elisa Zampieri¹, Andreas Ioannou⁴, Alexandros Spanos⁴, Federico Vita⁵, Miguel González-Guzmán², Vasileios Fotopoulos⁴, Vicent Arbona², Raffaella Balestrini^{6*}

[^]These authors share the position as first authors.

¹Institute for Sustainable Plant Protection, National Research Council (CNR-IPSP), Strada delle Cacce 73, Torino 10135, Italy

²Departament de Biologia, Bioquímica i Ciències Naturals, Universitat Jaume I, Castelló de la Plana, Spain

³Research Centre for Viticulture and Enology – Council for Agricultural Research and Economics (CREA-VE), Via XXVIII Aprile 26, Conegliano (TV) 31015, Italy

⁴Department of Agricultural Sciences Biotechnology and Food Science, Cyprus University of Technology, Lemesos 3036, Cyprus

⁵University of Bari ‘Aldo Moro’, Department of Biology, Via Orabona 4, Bari 70124, Italy

⁶Institute of Biosciences and Bioresources, National Research Council (CNR-IBBR), Via Amendola 165/A, Bari 70126, Italy

***Author for correspondence:**

Raffaella Balestrini

Institute of Biosciences and Bioresources, National Research Council, Via Amendola 165/A, Bari 70126, Italy

raffaellamaria.balestrini@cnr.it

Authors' e-mail addresses:

Luca Giovannini: luca.giovannini@ipsp.cnr.it; Chiara Pagliarani: chiara.pagliarani@ipsp.cnr.it; Eva Cañizares: ecanizar@uji.es; Fabiano Sillo: fabiano.sillo@ipsp.cnr.it; Walter Chitarra: walter.chitarra@crea.gov.it; Silvia De Rose: silvia.derose@ipsp.cnr.it; Elisa Zampieri: elisa.zampieri@ipsp.cnr.it; Andreas Ioannou: andrejkjg.ioannou@edu.cut.ac.cy; Alexandros Spanos: al.spanos@edu.cut.ac.cy; Federico Vita: federico.vita@uniba.it; Miguel González-Guzmán: mguzman@uji.es; Vasileios Fotopoulos: vassilis.fotopoulos@cut.ac.cy; Vicent Arbona: vicente.arbona@camn.uji.es; Raffaella Balestrini: raffaellamaria.balestrini@cnr.it

Highlight: Priming with arbuscular mycorrhizal fungi and natural compounds enhances the resilience of tomato plants to water deficit and salinity through transcriptome and metabolome remodelling, boosting antioxidant and osmoprotective responses.

© The Author(s) 2024. Published by Oxford University Press on behalf of the Society for Experimental Biology. All rights reserved. For commercial re-use, please contact reprints@oup.com for reprints and translation rights for reprints. All other permissions can be obtained through our RightsLink service via the Permissions link on the article page on our site—for further information please contact journals.permissions@oup.com.

Abstract

Priming modulates plant stress responses before the stress appears, increasing the ability of the primed plant to endure adverse conditions and thrive. In this context, we investigated the effect of biological (*i.e.*, arbuscular mycorrhizal fungi, AMF) agents and natural compounds (*i.e.*, salicylic acid applied alone or combined with chitosan) against water deficit and salinity on a commercial tomato genotype (*cv.* Moneymaker). Effects of seed treatments on AMF colonization were evaluated, demonstrating the possibility of using them in combination. Responses to water and salt stresses were analysed on primed plants alone or in combination with the AMF inoculum in soil. Trials were conducted on potted plants by subjecting them to water deficit or salt stress. The effectiveness of chemical seed treatments, both alone and in combination with post-germination AM fungal inoculation, was investigated using a multidisciplinary approach that included eco-physiology, biochemistry, transcriptomics, and untargeted metabolomics. Results showed that chemical seed treatment and AM symbiosis modified the tomato response to water deficit and salinity triggering a remodelling of both transcriptome and metabolome, which ultimately elicited the plant antioxidant and osmoprotective machinery. The plant physiological adaptation to both stress conditions improved, confirming the success of the adopted approaches in enhancing stress tolerance.

Keywords: chemical priming, arbuscular mycorrhizal symbiosis, water deficit, salinity, transcriptomics, metabolomics

Abbreviations: T_0 : start of the trial; T_f : end of the trial; AMF: arbuscular mycorrhizal fungi; SA: salicylic acid-treated; CHI+SA: chitosan+salicylic acid-treated; CTRL: untreated; NS: not-stressed; WS: water stress; SS: salt stress; MYC: inoculated; NMYC: not-inoculated; RWC_{soil} : soil relative water content; g_s : stomatal conductance; E : leaf transpiration; VPD_{leaf} : leaf Vapour Pressure Deficit; Ψ_{stem} : stem water potential; A_N : assimilation; CCI: Chlorophyll Content Index; ABA: abscisic acid; JA: jasmonic acid; IAA: indol-3-acetic acid; RNA-seq: RNA sequencing; DEGs: differentially expressed genes; 2D-PLSDs: supervised bi-dimensional analysis.

Accepted Manuscript

Introduction

Intensive agriculture progressively exacerbated the plant's exposure to environmental stressors, adversely impacting both crop productivity and quality yields (Francini and Sebastiani, 2019; Fróna *et al.*, 2021). In the last years, ongoing climate change has been worsening the frequency and severity of heat waves and drought events, accompanied by a general decrease and discontinuity in rainfall on agricultural ecosystems across the planet (Rosenzweig *et al.*, 2014; Horton *et al.*, 2016). A further problem occurring in combination with prolonged drought spells is a high risk of salt stress (Qadir *et al.*, 2014), resulting from reduced irrigation and precipitation and heat-driven evaporation.

The challenge of ensuring yield and quality of crops under this complex and worrying scenario may be tackled by developing and implementing sustainable measures that can endow target plant species with superior stress tolerance, thereby fostering the robustness and resilience of cropping systems. To this purpose, various strategies have been applied, such as traditional breeding and recently gene editing techniques that, however, still encounter applicability restrictions in several European countries (Hu and Xiong, 2014). An alternative and valuable solution is represented by priming agents able to stimulate the innate plant defence system (Mauch-Mani *et al.*, 2017; Alagna *et al.*, 2020). The plant perception of the priming stimulus triggers the establishment of a complex array of signalling routes, involving epigenetic, transcriptional, post-transcriptional, metabolic and physiological modifications (Hilker and Schmülling, 2019), ultimately leading to an overall improvement of the plant resistance/tolerance to stress (Mauch-Mani *et al.*, 2017).

Priming through chemical compounds applied at the seed and seedling level is very promising but further elucidation is needed to understand the mechanisms behind (Savvides *et al.* 2016; Gohari *et al.*, 2020 and 2024; Tripathi *et al.*, 2024). Among these compounds, chitosan (CHI) and salicylic acid (SA) have emerged as seed priming agents able to promote plant tolerance to abiotic stresses (Hidangmayum *et al.*, 2019; Kulak *et al.*, 2021; Chakma *et al.*, 2021). Chitosan is a poly-(D)-glucosamine derived from chitin, and as priming agent, it has been shown to elicit the ability of plants to endure stress conditions, especially against biotic stresses (Iriti and Varoni, 2015). However, its beneficial impact in triggering abiotic stress responses in different plant species has also been reported (Balusamy *et al.*, 2022; Hidangmayum *et al.*, 2023a,b). Chitosan effects on plants subjected to water stress and salinity are associated with the regulation of cellular osmotic pressure by promoting the availability/uptake of water and essential nutrients

(Hidangmayum *et al.*, 2019). On the other hand, salicylic acid (SA), a well-known plant hormone, plays a key role in the control of plant responses to both biotic and abiotic stresses (Rivas-SanVicente and Plasencia, 2011). Salicylic acid is known to regulate diverse physiological processes, including stomatal closure, photosynthesis, and the synthesis of some stress-related proteins and antioxidants (Janda *et al.*, 2014), and it has been proven to be effective against water deficit conditions (Chakma *et al.*, 2021). Together, CHI and SA may offer complementary strategies for priming plants against abiotic stresses. Their combined use as seed priming agents could therefore represent a promising approach for enhancing crop resilience under challenging environmental conditions.

Additionally, mounting experimental evidence showed that biological agents, such as root-associated microorganisms, can lead plants in a priming state that enable them to react quickly and robustly to stress exposure (Balestrini *et al.*, 2018; Alagna *et al.*, 2020). Arbuscular mycorrhizal fungi (AMF), which are obligate biotrophs currently assigned to the subphylum *Glomeromycotina* (Spatafora *et al.*, 2016), establish mutualistic symbioses with the roots of about the 80% of plant species, including the most important crops (Balestrini *et al.*, 2021). AMF play a crucial role in the acquisition and translocation of essential mineral nutrients using their extraradical mycelium, while receiving carbon compounds in return (Smith and Read, 2008). The interaction with AMF can improve plant adaptation to various stress conditions, including water and salt stresses (Balestrini *et al.*, 2018; Bastías *et al.*, 2022). Root symbiosis with AMF positively affects plant physiological parameters, biochemical pathways and gene expression, resulting in an improved tolerance of the host plants (Bastías *et al.*, 2022). Several works investigated transcriptomic changes occurring in different species of agronomic interest exposed to environmental stresses during AM symbiosis (Balestrini *et al.*, 2019; Zhang *et al.*, 2019; Moradi Tarnabi *et al.*, 2020; Recchia *et al.*, 2018). Moreover, untargeted metabolomic approaches have successfully been applied to explore the whole metabolic changes controlling the interaction between the host plant and AMF. Differences in the metabolic profiles of roots and leaves from mycorrhizal crops have emerged, allowing the identification of compounds potentially involved in stress tolerance (Rivero *et al.*, 2018; Tugizimana *et al.*, 2018; Liang *et al.*, 2021).

Tomato (*Solanum lycopersicum* L.) is one of the most important crop species worldwide (Cortina *et al.*, 2005) with a global production of 186.1 million metric tonnes (mT) in 2022 (FAOSTAT database, <http://www.fao.org/faostat/en/#data>). The availability of an updated

reference genome sequence and the recent release of the genomes of many commercial and wild tomato genotypes makes this species an ideal model for physiological and functional genomic studies aimed at identifying stress tolerance mechanisms and stress marker genes for breeding purposes (Wang *et al.*, 2024). Water limitation was reported to have a negative effect on the tomato photosynthetic activity (Chitarra *et al.*, 2016) and nutrient uptake, resulting in an overall decrease in fruit production (Ma *et al.* 2019; Sardans *et al.*, 2020). Soil salinity has also been shown to influence tomato growth and fruit yields (Roşca *et al.*, 2023 and references therein). Therefore, tomato represents a model crop for studying the role of the arbuscular mycorrhizal (AM) symbiosis in the regulation of abiotic stress responses. Several authors reported a positive impact of AM symbiosis in tomato tolerance to water deficit through different mechanisms (Chitarra *et al.*, 2016; Rivero *et al.*, 2018; Balestrini *et al.*, 2019; Leventis *et al.*, 2021; Duc *et al.*, 2023; Zhang *et al.*, 2024). Nonetheless, the application of AMF as stress priming agents is still a challenge in agriculture, mainly due to poorly consistent results (Santos *et al.*, 2019). In this context, the crop genotype, timing and intensity of different single and/or combined stresses represent variables that can significantly affect the interaction of plants with beneficial microbes (Pascale *et al.*, 2020). Furthermore, native AMF of a particular soil or site are often reported to be more effective than non-native ones, probably as they are adapted to soil features such as nutrient concentrations, or to environmental constraints such as drought (Pellegrino *et al.*, 2011). Within this scenario, other poorly explored aspects concern the establishment of stress tolerance signals from combined AMF and abiotic priming treatments.

This study was focused to verify the effectiveness of chemical priming (*i.e.* salicylic acid, SA and chitosan+salicylic acid, CHI+SA) applied to tomato seeds prior to germination, in combination with an AM fungal inoculum in the soil and considering diverse abiotic stress conditions (water deficit and salinity). A multidisciplinary approach, involving physiological measurements, transcriptomic, biochemical and untargeted metabolomic analyses, was followed to deciphering the mechanisms involved in the tomato response to priming and AM symbiosis.

Materials and methods

Plant material and growth conditions

Solanum lycopersicum cv Moneymaker (MM) seeds (Semiorto Sementi, Sarno, Italy) were treated with natural compounds by the Cyprus University of Technology (CUT; Galviz-Fajardo *et al.*

2020). Seeds were treated with chitosan (CHI), salicylic acid (SA) or chitosan+salicylic acid (CHI+SA) as potential priming agents, while untreated seeds were used as control (CTRL).

A homogeneous and clear solution of low molecular weight chitosan (CHI) (CAS Number: 9012-76-4) was prepared by dissolving 0.1 g of CHI in 85 ml of deionized water (dH₂O) along with 100 µl of acetic acid (0.5%), followed by stirring for 1 hour. To achieve a concentration of 50 µM Salicylic acid (SA), 10 ml of 500 µM SA was prepared in dH₂O and added to the homogenous CHI solution while stirring. Tripolyphosphate (TPP) was used to cross-link the CTS nanocarrier, calculated based on the amount of CHI used. In this regard, to form CHI nanoparticles and encapsulate SA, 0.4 g of sodium tripolyphosphate (TPP) (CAS Number: 7758-29-4) was dissolved in 5 ml of dH₂O. The addition of TPP resulted in crosslinking of CHI nanoparticles in the form of coagulation, which was continued by stirring the nano-carriers overnight (Hassanpour *et al.*, 2021). The CHI alone treatment consisted of an identical approach, omitting the step where SA was prepared in dH₂O and added to the homogenous CHI solution while stirring. Seed priming was performed by soaking tomato seeds in the two different solutions (SA and CHI+SA) for 12h at 25°C in the dark. After soaking, the seeds were spread on a filter paper to air-dry at room temperature. Once the primed seeds were dried, they were ready to be sown.

Treated and untreated seeds were germinated in presence/absence of an AM fungal inoculum in the soil. A crude inoculum of *Funneliformis mosseae* spores, extraradical mycelium, and colonized roots (MycAgro Lab, Bretenièrre, Francia) was used. Seeds were placed in pots (0.7 l) filled with pumice at the bottom, followed by coconut fibres and a ring with sterilized (at 180°C for 3h) quartz sand and coconut fibres at the top (NMYC). Regarding AMF (MYC) pots, the inoculum was mixed with the sterile sand ring, using 100 ml of inoculum with 300 ml of sterile sand (1:4) for 4 pots. The pots were watered two times a week with tap water and once a week with ½ Hoagland solution (Hoagland and Broyer, 1936), starting one month after sowing. Plants were cultivated in a glasshouse, with average daily temperatures of 24.9±5.35°C, average relative humidity of 42.3–61.8% and maximal photosynthetic photon flux density (PPFD) of 900–1200 µmol photons m⁻² s⁻¹. When necessary, a photoperiod of 12-h-light/12-h-dark was obtained with halogen lamps to guarantee a minimum PPFD of 500-600 µmol photons m⁻² s⁻¹. During the physiological trial, one third of the plants were subjected to water stress (WS), one third to salt stress (SS) and the remaining ones were maintained fully irrigated and used as control (NS). For each treatment

(CTRL, SA, CHI, CHI+SA, MYC, NMYC) and condition (NS, WS, SS), eight biological replicates were used, for a total of 144 plants.

Water deficit time-course

Water deficit was imposed by withholding irrigation on two months old plants belonging to the different experimental theses. Dynamic changes in stomatal conductance (g_s) rates were monitored over time on WS and NS plants using a portable porometer/fluorometer (LI-600, Lincoln, Nebraska USA). Leaf Vapour Pressure Deficit (VPD_{leaf}) values were checked using the same instrument, which also allowed calculation of transpiration rates (E). Daily changes in water loss were also inspected, by measuring the soil relative water content (RWC_{soil} , %) gravimetrically on potted plants exposed to water deprivation according to Pagliarani *et al.* (2022). Additionally, the stem water potential (Ψ_{stem}) was measured on NS and WS plants at the end of the trial (T_f) as described in Pagliarani *et al.* (2022), using a portable pressure chamber (1505D PMS Instrument Company, Albany, OR, USA). The water stress time-course was stopped when RWC values were around 40% and g_s rates were equal or lower than $0.05 \text{ mol H}_2\text{O m}^{-2} \text{ s}^{-1}$. At the beginning (T_0) and end (T_f) of the experiment, plant height, leaf number, stem diameter at the basal level of all plants were measured. The Chlorophyll Content Index (CCI) was also determined on the second and third fully developed leaves from the apex, using a portable chlorophyll meter SPAD 502 (CCM-200; Opti-Sciences). Two leaves *per* plant were analysed, taking three readings per leaf.

Salt stress experiment

After a month and half from the sowing the salt stress was imposed by supplying tomato plants with a solution of NaCl (50 mM), providing 30 ml for each pot. The NaCl concentration was gradually increased every three days first to 100 mM, and then to 150 mM, until reaching the maximum concentration (200 mM), according to Pollastri *et al.* (2018). The maximum concentration was maintained for eleven days, administered every three days. After 20 days of stress, the experiment was stopped and changes in the main eco-physiological parameters were determined on stressed (SS) and non-stressed (NS) plants. Rates of stomatal conductance (g_s), transpiration (E) and assimilation (A_N) were measured in the morning (between 9:00 and 12:00 h a.m.) on each plant, using a portable infrared gas analyzer (LCpro-SD system, ADC BioScientific Ltd, Hertfordshire, UK) equipped with a 6.25 cm^2 -leaf chamber. Three random, fully expanded,

non-senescent leaves exposed to direct sunlight were analyzed working with artificial irradiation ($1200 \mu\text{mol photon m}^{-2} \text{s}^{-1}$) and setting the chamber temperature at 25°C . CO_2 values were those of greenhouse conditions (450 ppm).

Sampling procedure

At the end of the WS and SS experiments, leaves and roots collected from stressed (WS or SS), and not-stressed (NS) plants were immediately frozen in liquid nitrogen and stored at -80°C . Half of the root system was used for the mycorrhizal colonization assessment.

Assessment of mycorrhizal colonization

After two months from seedling inoculation, mycorrhizal colonization was checked on three plants for each priming treatment and condition (NS, WS or SS). Roots were gently washed with tap water, then they were incubated in a KOH (10%) solution for 10 min at 80°C , followed by an overnight staining at room temperature with 0.1% cotton blue in lactic acid. Roots were then gently washed with tap water three times and stored in lactic acid. Twenty randomly selected root fragments (1 cm in length) were mounted on microscope slides (three slides *per* biological replicate). A total of 60 root fragments were observed *per* biological replicate. The level of mycorrhizal colonization was determined using the method described by Trouvelot *et al.* (1986).

Phytohormone quantification

Analysis of plant hormones (SA, JA, ABA, IAA) was carried out by LC/MS as described (De Ollas *et al.* 2021). Briefly, 10 mg of ground dry leaf tissue ($n=3$) was extracted in 1 ml of ultrapure water, after spiking with 25 μl of a solution containing 1 mg l^{-1} of $[\text{}^2\text{H}_2]$ -GA7, $[\text{}^2\text{H}_6]$ -ABA, DHJA, and $[\text{}^{13}\text{C}_6]$ -SA, and 0.1 mg l^{-1} of $[\text{}^2\text{H}_5]$ -IAA in a ball mill (MillMix20, Domel, Železniki, Slovenija). Extracts were centrifuged at 10,000 rpm and 4°C for 10 min and the supernatants recovered and their pH adjusted to 3 with 30% acetic acid. Afterwards, all extracts were partitioned twice against 1 ml of diethylene ether, recovering the organic layer on each occasion. The combined organic layers were evaporated under vacuum (Speed Vac, Jouan, Saint Herblain Cedex, France) and the dry residue resuspended in 500 μl of a 1:9 methanol:water solution by sonication. The resulting solutions were filtered through 0.2- μm polytetrafluoroethylene membrane syringe filters (Albet S.A., Barcelona, Spain). Filtered extracts were separated by reversed phase UPLC

(Acquity SDS, Waters Corp., Milford, MA, USA, or Waters Alliance 2695, Waters Corp) using a C18 column (Luna Omega Polar C18 100 x 2.0 mm, Phenomenex, CA, USA) and an acetonitrile and water gradient (both supplemented with formic acid) at a constant flow rate of 0.3 mL min⁻¹. The mass spectrometer (TQ-D, Micromass Ltd., UK) was operated in negative ionization electrospray mode and the different plant hormones were detected according to their specific transitions, using a multiresidue mass spectrometric method. Hormone quantitation was performed with Masslynx v. 4.0 software (Micromass Ltd., Manchester, UK) after external calibration with standard samples containing known amounts of each plant hormone.

Proline content

Proline content was determined according to Bates *et al.* (1973) and Shabnam *et al.* (2016), with slight modifications. Briefly, leaf tissues (10 mg of dry weight) from three biological replicates were extracted by homogenization in 1 mL of 1% aqueous sulfosalicylic acid (w/v). One mL of plant extract was reacted with 400 µl acid ninhydrin reagent (1.25% ninhydrin in 80% glacial acetic acid) in a water bath at 100°C for 30 min or 60 min. The reaction was stopped by incubating the samples on ice for 10 min. Proline content was determined by reading the sample absorbance at 520 nm using a multi-mode reader (Synergy H1 Hybrid Multi-Mode Microplate Reader, BioTek, USA).

RNA extraction, library preparation and sequencing

Total RNA was extracted from 0.1 g of leaf material according to the C-TAB-based method by Chang *et al.* (1993). Quantity and quality of the extracted RNA samples were determined using a Nanodrop 2000 spectrophotometer (Thermo Fisher Scientific, Waltham, MA, USA). RNA integrity was assessed on an RNA 6000 Nano Labchip using a Bioanalyzer 1000 (Agilent Technologies, Santa Clara, USA) prior to library preparation. Only samples showing a RIN (RNA Integrity Number) value > 8 were processed for sequencing. Library preparation and RNA sequencing were performed at an external service (IGATech s.r.l laboratories, Udine, Italy). In detail, Universal Plus mRNA-Seq kit (Tecan Genomics, Redwood City, CA) was used for library preparation following the manufacturer's instructions and sequenced on an Illumina NovaSeq 6000 apparatus (Illumina, San Diego, CA, USA). Trimming of lower quality bases and adapters was performed using the ERNE software according to Del Fabbro *et al.* (2013).

RNA-seq data analysis

For alignment, reads were mapped onto the reference genome GCF_000188115.5_SL3.1 (Hosmani *et al.*, 2019) using STAR v. 2.7.10 (Dobin *et al.*, 2013), a splice junction mapper designed for RNA-Seq reads, under default parameters. The software htseq-count v. 2.0.2 (Anders *et al.*, 2015) was utilized to count the overlapping of reads with genes. The data were then used to identify differentially expressed genes (DEGs) using the DESeq2 package v1.34.0 (Love *et al.*, 2014). The variance on read count was calculated based on three biological replicates per condition by applying a negative binomial distribution to model the count data, therefore identifying genes showing significant expression changes among the different tested conditions. The DEG identification was performed after normalization of the count data and correction for multiple testing, both accounted by DESeq2, through the Wald test. During DESeq2 analysis, the shrinkage estimation of effect size (LFC estimates) was used, to generate more accurate Log₂ foldchange estimates and considering the variability among replicates. A cut-off of the *p*-adjusted value < 0.05 was used to classify a gene differentially expressed (DEG) in comparison with the reference (*i.e.* untreated (CTRL) in not-stressed condition (NS) and without AM fungal inoculation (NMYC)). Both the identified DEGs and all transcripts of the tomato (*Solanum lycopersicum* L.) transcriptome were annotated through Blast2GO v5.2.5 (Conesa *et al.*, 2005) to obtain an updated functional annotation and to assign the corresponding Gene Ontology (GO) terms. A gene class functional enrichment analysis was then conducted using Blast2GO to reveal the biological processes, pathways, or other functional categories that are enriched among the identified DEGs.

Non-targeted polar metabolite profiling

Polar metabolites were separated using hydrophilic interaction liquid chromatography (HILIC) coupled to hybrid quadrupole-time of flight mass spectrometry (QTOF-MS) according to Andrade *et al.* (2021). HILIC separation was performed on a 2.1×100 mm InfinityLab Poroshell 120 HILIC-Z, 1.9 μm (Agilent Technologies, Inc., Santa Clara, CA, USA) using acetonitrile:water, 95:5 (v/v) (solvent A) and acetonitrile:water, 2:98 (v/v) (solvent B), both supplemented with ammonium formate at 0.063% and 0.126%, respectively, as solvents and at a flow rate of 0.3 ml min⁻¹. During chromatographic runs, column temperature was at 40°C. Tomato leaf samples (10 mg of dry weight) were extracted in triplicate by ultrasonication in 300 μl of 80% aqueous methanol

supplemented with kinetin (1 mg l^{-1}) as internal standard for relative quantitation. After extraction, samples were centrifuged at 10,000 rpm and 4°C for 10 min and the supernatants recovered. Subsequently, supernatants were diluted 1:4 with acetonitrile (LC/MS grade) and filtered through $0.2 \mu\text{m}$ PTFE syringe filters directly into chromatography vials. Mass chromatographic data were acquired in positive and negative ionization modes within the 50–1000 amu mass range. Nitrogen was used both as nebulization and desolvation gas (60 and 800 l h^{-1} and 350°C temperature, respectively). During measurements, capillary and cone voltages were set at 3.5 kV and 30 V for positive electrospray and 2.3 kV and 30 V for negative electrospray, respectively. An additional acquisition function to obtain collision-induced dissociation (CID) information was set by performing a voltage ramp between 6–40 eV. To ensure accurate mass data acquisition, a lockmass reference (Leu-Enkephalin, $[\text{M}+\text{H}]^{+} 556.2771$ and $[\text{M}-\text{H}]^{-} 554.2614$) was regularly infused during runs. Data files from each ionization mode were converted to mzML with msconvert (Chambers *et al.* 2012) and processed independently with XCMS (Smith *et al.*, 2006). Mass chromatographic features were annotated and grouped with CAMERA (Kuhl *et al.*, 2012). Peak areas were normalized to internal standard area and actual sample weight before statistical analyses. Significantly altered mass chromatographic features were subsequently identified as individual compounds by matching m/z and retention time values with those of authentic standards or tentatively annotated by matching experimental mass spectra in public databases (Metlin, Massbank or HMDB).

Statistical analysis

Statistical analyses were performed using the R software (version 4.1.1). For colonization a two-way ANOVA test was conducted to determine the effect of priming treatment (T) and stress condition (S). Regarding physiological, biometric (including CCI), and hormonal data (including proline), a three-way ANOVA test was conducted to determine the effect of priming treatment (T), AM fungal inoculation (I), and stress condition (S), and of their interaction ($\text{T} \times \text{I} \times \text{S}$). A one-way ANOVA was applied to evaluate the combined effect of priming treatment and inoculation on soil relative water content (RWC_{soil}) data during the first four days of experiment. Standard error of all means was calculated. Mean separation was performed using the Tukey HSD test ($p \leq 0.05$). Graphs were generated using the Prism software (La Jolla, CA, USA v.9.5.0), while two-dimensional Partial Least Squares Discriminant Analysis (2D-PLSDAs) was executed using the

MetaboAnalyst 6.0 software. The UpSet plot, heatmaps and GO-enriched plot were elaborated respectively using the upsetplot, pheatmap R and ggplot2/dplyr packages (R version 4.1.1).

Integration between transcriptomic and metabolomic data

To integrate data from RNA-seq and untargeted metabolomics, the DIABLO (N-integration, multi-block sPLS-DA) framework was used (Singh *et al.*, 2019). The input datasets included RNA-seq data (normalized transcript counts of 14'315 genes resulted as DEGs in at least one condition) and metabolomic data (peak areas of 230 significantly-altered mass chromatographic features), both normalized using variance stabilizing transformation and \log_2 transformed. For design the model, the optimal number of features was obtained by performing sparse partial least squares discriminant analysis (sPLS-DA) for each dataset. The final DIABLO model was refined through cross-fold validation repeated 10 times, considering the overall and the balanced classification error rates to identify the optimal number of components. The function *plotArrow()* was used to visualise the level of agreement between the datasets. Correlations between the different types of variables (RNA-seq and metabolomic data) was visualized through the relevance network plots via *network()*, which were inspected and customized by using Cytoscape v3.10.2. In the networks, edges showing weight higher than 0.95 were considered as key edges.

Results

AM fungal root colonization

To assess differences in root mycorrhizal colonization, the mycorrhization frequency (F%) and arbuscules in mycorrhizal root parts (a%) were evaluated (Fig. 1 and Supplementary Table S1). No colonization structures were detected in not-inoculated roots. Regarding inoculated plants, F% tended to be higher following SA treatment with respect to other treatments (Fig. 1A), but no differences were observed among the samples in terms of arbuscule abundance (a%; Fig. 1B). Under water stress (WS), CHI+SA roots displayed significantly higher F% values compared with CTRL (Fig. 1A). Differently, the salt stress (SS) condition associated with SA and CHI+SA treatments led to a lower F% than that of CTRL plants and higher a% values, although not statistically significant (Fig. 1B).

Effect of seed priming and mycorrhizal inoculation on tomato eco-physiological responses to water deficit and salinity

Over the whole duration of the water deficit time-course (15 days), inoculated (MYC) SA and CHI+SA plants depleted water by transpiration more slowly with respect to both inoculated CTRL and all not-inoculated (NMYC) plants, indicating that water extraction dynamics under WS were improved in mycorrhized SA- and CHI+SA-primed plants (Fig. 2A, Supplementary Table S2). This trend was particularly evident during the early times after water withdrawal, especially considering the significant changes occurring in the soil relative water content (RWC_{soil}) of pots containing CHI+SA_MYC plants. In absence of seed priming, the inoculation with AMF alone did not alter the plant water loss dynamics, as RWC_{soil} values in CTRL_MYC plants were close, even slightly lower in the last days of WS imposition than those measured in all NMYC plants (Fig. 2A). Moreover, the analysis of stomatal conductance (g_s) patterns in AMF-inoculated plants (Fig. 2B, Supplementary Table S3) showed that, compared to CTRL_MYC, CHI+SA_MYC and SA_MYC plants were able to maintain higher g_s rates during the first days of water shortage. This finding might therefore reflect a direct effect of seed priming in the regulation of water relations. Notably, this response was observed despite a more pronounced increase in leaf Vapour Pressure Deficit (VPD_{leaf}) occurred in these plants (Fig. 2C). However, patterns of leaf transpiration (E) over stress progression were similar to those of g_s (Fig. 2D, Supplementary Table S3), likely indicating that the glasshouse ambient conditions were overall stable during the experimental trial and did not alter the plant physiological response to water deprivation. The combined effect of seed priming agents and AM fungal inoculation on the monitored parameters was also confirmed at the end of the WS time-course (T_f) by comparing the different plant groups in terms of stem water potential (Ψ_{stem}) (Supplementary Fig. S1) and g_s and E measurements (Fig. 2E and Supplementary Fig. S2). Xylem pressure values were significantly higher in CHI+SA_MYC and SA_MYC plants with respect to all other plant groups subjected to WS. This condition was more accentuated in the comparison between SA_NMYC_WS and SA_MYC_WS plants: the former reached Ψ_{stem} around -0.8 MPa, attesting a water deficit condition for potted tomato plants of this age and size (1 month and half-old and 27 cm height), whereas the latter showed a Ψ_{stem} close to -0.2 MPa, not significantly different from that of all NS plant groups (Supplementary Fig. S1, Supplementary Table S3). Although at T_f all WS plants showed g_s rates up to one fourth lower than those of the corresponding NS controls, stomatal limitation in the inoculated SA_MYC plants was

milder, resulting in significantly higher g_s values (Fig. 2E, Supplementary Table S3). The same did occur at the level of leaf transpiration patterns (Supplementary Fig. S2).

The effect of AM fungal inoculation and seed priming treatments, alone or in combination, was also noticed by analysing the leaf gas exchanges of tomato plants in response to SS at the end of the treatment with NaCl (200 mM). In absence of AM fungal inoculation and stress imposition (NMYC_NS), g_s rates were significantly lower in SA- and CHI+SA-primed plants than in CTRL, likely suggesting an effect of the treatment in the regulation of stomata opening. Such a condition was also observed upon SS. In fact, while g_s significantly dropped in all NMYC plants but at different extend based on the priming treatment, CHI+SA NMYC SS plants only showed a three-fold decrease in g_s in comparison with CHI+SA NMYC NS plants. Furthermore, SA NMYC SS experienced a two-fold, non-significant reduction in g_s with respect to SA NMYC NS plants (Fig. 3A, Supplementary Table S4). These data suggested that the negative effect of SS on stomatal closure was counteracted by chemical priming. In presence of AM symbiosis (MYC) and in absence of stress, the stomata of CTRL plants were significantly more closed than CTRL_NMYC plants. Notably, such limitation was not observed in CHI+SA_MYC and SA_MYC plants compared to the respective NMYC plants. Nonetheless, independently of seed priming, MYC plants were all able to maintain g_s rates upon SS conditions, with values close or even slightly higher than those measured in NS leaves (Fig. 3A). It thus emerged a significant effect of the tripartite interaction among priming treatment (T), inoculation (I) and stress condition (S) on the regulation of stomatal closure in response to salinity. This observation was also confirmed by analysing changes in E , which mirrored the trend described for g_s (Fig. 3B). The effect of the T×I×S interaction also significantly affected the plant photosynthetic capacity in terms of assimilation rates (A_N). In fact, in absence of both AM fungal inoculation and seed priming, net photosynthesis steeply decreased (by more than one half) in CTRL_SS plants with respect to the CTRL_NS condition (Fig. 3C). Moreover, although under SS there was a trend towards increased A_N in AMF-inoculated plants without priming agents compared with uninoculated plants under the same conditions, the changes were not statistically significant. The A_N levels in SA_MYC_SS and CHI+SA_MYC_SS plants were similar to those in CTRL_NMYC_SS, suggesting that the benefits of priming and presence of AMF under SS were not additive, at least for this parameter (Fig. 3C). In parallel, eco-physiological responses, as well as AM fungal colonization, were also checked in plants treated with CHI alone (Supplementary Figs. S3-S4). Although not statistically significant

differences were detected, AM fungal colonization in those plants was lower, either in terms of frequency of mycorrhization in not stressed condition (Supplementary Fig. S3A) or in terms of arbuscule abundance upon WS (Fig. S3B), in comparison with the CHI+SA treatment. Moreover, no significant changes were observed by analysing stomatal conductance rates at the end of the WS and SS imposition in CHI- and CHI+SA-treated plants (Supplementary Fig. S4A, B). Finally, considering the effect of the combination of seed priming and AM fungal inoculation, it emerged that, unlike CHI+SA-treated plants, tomato plants treated with chitosan alone were not able to maintain g_s levels upon SS (Supplementary Fig. S4B). For these reasons and based on the need to avoid adding further complexity to the analysed processes, we did not consider chitosan for further analyses.

Biometric features (Supplementary Table S5) indicated that while significant reductions in shoot height were observed following exposure to both stresses and in presence of AM fungal inoculation, the overall effects of MYC and seed priming on the other morphometric parameters were generally limited. Particularly, SA_MYC plants upon both stress conditions showed the least number of leaves (Supplementary Table S5). Stem diameter was primarily affected by the imposed stress, as well as the leaf chlorophyll content index (CCI). For these two parameters, independently of seed priming or AM fungal inoculation, both WS and SS tomato plants showed significantly lower values compared with NS ones (Supplementary Table S5).

Phytohormones and proline are differently accumulated in tomato leaves based on stress, priming treatment and AM symbiosis

Changes in the content of the main stress-associated phytohormones, namely SA, jasmonic acid (JA), abscisic acid (ABA) and indol-3-acetic acid (IAA), were investigated to elucidate the impact of either seed priming treatments or AM fungal inoculation or their combination on plant responses to the imposed stresses. In NMYC plants, SA concentrations decreased in CTRL and SA leaves upon both stress conditions. SA levels were still high in CHI+SA plants following WS, while they decreased during SS with respect to the corresponding NS samples, suggesting a regulation of the hormone metabolism based on the abiotic stress type (Fig. 4A, Supplementary Table S6). A completely opposite trend emerged from the analysis of MYC_NS leaf samples, which showed a much lower SA accumulation than NMYC_NS plants. Regardless of the priming treatment, SA content in MYC plants was always higher upon both WS and SS in comparison to MYC_NS plants

(Fig. 4A). JA levels in NMYC plants were significantly lower in WS and SS leaves, independently of the priming treatment. This phenomenon did not occur in MYC plants, which during WS maintained JA content similar (CTRL_MYC and CHI+SA_MYC) or even significantly higher (SA_MYC) than the corresponding NS plants. The T×I×S effects were particularly evident in SS plants. A reduction in the hormone levels was in fact noticed upon SS in CTRL_MYC and SA_MYC samples, whereas in CHI+SA_MYC_SS plants JA concentrations were close to those measured in NS and WS conditions (Fig. 4B, Supplementary Table S6).

Upon stress conditions, ABA content increased in NMYC plants, with no significant variations between WS and SS. In AMF-inoculated plants, except for a slight, but not significant, increase in CTRL_MYC_WS leaves, ABA levels were similar among all other samples (Fig. 4C, Supplementary Table S6). The only variable that significantly influenced IAA accumulation patterns was the stress condition, particularly salinity both in MYC and NMYC plants (Fig. 4D, Supplementary Table S6). About the osmoprotectant solute proline, levels of the metabolite increased in NMYC plants based on the stress application, with a steeper accumulation in SS samples (Fig. 4E, Supplementary Table S6). A similar pattern occurred in CTRL plants in presence of AM fungal inoculation. The impact of T×I×S interaction was instead evident in the case of inoculated seed primed plants, but with a different trend based on the seed priming agent. Indeed, in SA_MYC samples proline increased both upon WS and SS with respect to NS conditions, whereas in CHI+SA significantly higher concentrations of the solute were measured exclusively upon SS (Fig. 4E, Supplementary Table S6).

General overview of tomato leaf transcriptomic profiles

To investigate systemic transcriptional changes, a transcriptomic approach *via* RNA-seq was performed. After Illumina NovaSeq 6000 sequencing of 54 libraries, a total of 1,857,600,000 raw reads were generated. The average percentage of mapped reads obtained from all analyzed samples to the reference tomato genome was 88.5% (Supplementary Dataset S1; see supporting datasets available at the Dryad data repository). Total read counts for each sample are reported in the Supplementary Dataset S2. A total of 14,318 genes (out of the 37,604 annotated *Solanum lycopersicum* genes) were differentially regulated across all samples. Complete lists of identified Differentially Expressed Genes (DEGs) in the 17 conditions in comparison to the control (CTRL_NMYC_NS) are reported

in Supplementary Datasets S3-S19 (see supporting datasets available at the Dryad data repository). A significant transcriptomic remodelling occurred following stress exposure, and particularly in plants subjected to SS, with SA_MYC_SS and CHI+SA_MYC_SS showing the highest number of DEGs (7150 and 6890, respectively; Fig. 5A and Supplementary Dataset S20). CHI+SA_MYC_SS was the condition showing the highest number of up-regulated DEGs (3512 genes; Fig. 5A), while in SA_MYC_SS leaves the highest number of down-regulated genes was detected (3696 genes; Fig. 5A). NMYC_NS plants *plus* CHI+SA and SA were interested by a limited leaf transcriptomic reprogramming with respect to CTRL_NMYC_NS (166 and 107 up- and down regulated genes for CHI+SA_NMYC_NS, and 171 up- and 69 down regulated genes for SA_NMYC_NS; Fig. 5A).

From the identification of DEGs in all samples compared to the control condition (CTRL_NMYC_NS), two distinct DEG clusters were detected (Fig. 5B). Cluster I harboured DEGs mainly up-regulated by both water deficit and salinity in MYC plants and mainly down-regulated in NS condition (Fig. 5B). Cluster II included DEGs primarily up-regulated in NS plants, and particularly in absence of AM fungal inoculation (Fig. 5B).

A GO enrichment analysis allowed grouping the identified up and down DEGs into functional classes based on the biological process (Supplementary Figs. S5, S6). In NS condition, a high number of GO terms were identified and particularly in SA_MYC_NS and CHI+SA_MYC_NS (Supplementary Fig. S5A, S6A). It is worth noting that “ABA-8’-hydroxylase activity”, correlated to ABA catabolic processes, was among the enriched terms in CHI+SA_MYC_WS (Supplementary Fig. S6B). This was in agreement with the lower ABA levels quantified in CHI+SA_MYC_WS leaves in comparison to CTRL_MYC_WS plants (Fig. 4C).

The impact of abiotic stress conditions on tomato leaf transcriptome

A core of stress marker genes was differentially regulated upon both stresses, independently of inoculation and seed priming. Particularly, 274 DEGs were up-regulated by WS (Supplementary Dataset S27). Most of them encoded proteins related to stress response (Supplementary Dataset S21), including two genes encoding dehydrins, *i.e.*, *ERD14* and *TAS14*. In detail, *TAS14* was particularly expressed upon WS in NMYC plants (average Log₂FC value of 6.03±0.44) compared to MYC samples (average Log₂FC of 3.96±0.87), consistently with the higher ABA content in NMYC with respect to MYC plants (Fig. 4). Four genes were exclusively up-regulated upon WS,

regardless of AM fungal inoculation and seed priming treatment (Fig. 6A): an abscisic stress-ripening protein 1 (*ASRI*), a serine/threonine-protein kinase SAPK10 (LOC101268620), a class I heat shock protein (LOC101256536), and a JA O-methyltransferase-like (LOC101265985; Supplementary Dataset S20).

Considering SS, irrespectively of the AM fungal inoculation and seed priming treatments, 517 DEGs were found up-regulated. Among them, genes involved in sugar transport and calcium signalling cascades, such as sodium/calcium exchangers, scarecrow-like transcription factors (TFs), calcium binding proteins, and heat shock proteins, were identified (Supplementary Dataset S22). Additionally, four genes, encoding a transcriptional regulatory protein DOT6 (LOC104644498), a protein SUPPRESSOR OF MORE AXILLARY GROWTH (MAX) 2-like (LOC101262825), a probable serine/threonine-protein kinase PBL1 (LOC101264267), and an uncharacterized protein (LOC101245413) were exclusively regulated upon SS (Fig. 6A and Supplementary Dataset S20).

The impact of AM symbiosis on tomato leaf transcriptome

Transcriptome reprogramming events dependent on AM fungal inoculation were also explored. Starting from CTRL_MYC_NS, 2026 genes (1153 up-regulated, 873 down-regulated) were differentially expressed. Among them, genes involved in defence, TFs, signal transduction, proteolysis, and genes encoding transporters and heat shock proteins were detected. A comparison of this dataset with that obtained by Cervantes *et al.* (2016) on the leaf transcriptome of tomato AMF-inoculated plants, led to identify 28 genes (18 up-regulated, 10 down-regulated) showing a similar expression trend (Supplementary Dataset S23). The remaining 45 DEGs in common with the previous dataset showed a different expression profile (Supplementary Dataset S23), likely due to the differences in the genotypes of the partners and in growth conditions. Looking at the 25 most up-regulated genes in CTRL_MYC_NS leaves, the most up-regulated ones (Supplementary Dataset S3) encoded a nudix hydrolase 8-like (LOC101261537, $\text{Log}_2\text{FC}=6.15$), which is known to play a role in plant defence (Fonseca and Dong, 2014), two genes coding for a subtilisin-like protease SBT1.9 (*sbt4d* and LOC101257301), two genes involved in phosphorous homeostasis, one encoded an inorganic pyrophosphatase 1-like (*psi14C*) and one purple acid phosphatase 17-like (LOC101249252). Among the up-regulated genes also a probable carotenoid cleavage dioxygenase 4 (LOC101250535) was detected, while a gene encoding a 9-cis-epoxycarotenoid

dioxygenase (*LeNCEDI*) resulted to be slightly down-regulated together with a carotenoid 9,10(9',10')-cleavage dioxygenase 1-like gene (*LeCCD1B*; Supplementary Dataset S20). These data partially supported what observed in terms of ABA accumulation patterns in these plants (Fig. 4C).

Looking at MYC plants subjected to WS (MYC_WS), 199 DEGs were identified as up-regulated. Forty-five out of these corresponded to pentatricopeptide repeat (PPR)-containing proteins (Supplementary Dataset S24). Furthermore, a gene encoding a 4-coumarate-CoA ligase-like 5 (LOC101251665) was found slightly up-regulated, regardless of priming treatment (Supplementary Dataset S20). Six transcripts were exclusive of MYC_WS plants (Fig. 6A), including genes encoding an inorganic pyrophosphatase 1-like (LOC101246036), a serine/threonine-protein kinase (LOC101262118) and a DPH4 homologue (LOC104645200; Supplementary Dataset S20).

MYC plants upon salt stress (MYC_SS) showed the up-regulation of 464 DEGs (Supplementary Dataset S25). Among these genes, 63 encoded putative PPR-containing proteins and it is worth noting that 38 DEGs were identified as exclusive for this condition (Fig. 6A; Supplementary Dataset S20).

The combined effect of AM symbiosis and priming agents on leaf transcriptome

A relevant combined effect of AM fungal inoculation and seed priming in the reprogramming of leaf transcriptome was observed under both NS and stressed conditions. Conversely, the presence of priming agents alone in NMYC plants only slightly perturbed the leaf transcriptome. Among the most up-regulated genes in the SA_NS treatments, a homologue of *LEA14* was detected (*ER5*, Log₂FC=5.26), already reported to protect pepper against desiccation (Zhang *et al.*, 2023). Considering the CHI+SA_NS treatment, one of the most up-regulated genes encoded a dehydration-responsive element-binding protein 1A-like (LOC104648613, Log₂FC=5.24) involved in oxidative stress responses (Zhang *et al.*, 2021). Looking at mycorrhizal plants, among the first 25 mostly up-regulated genes in SA_MYC_NS, 17 shared an expression profile similar to CTRL_MYC_NS (Supplementary Dataset S26). In CHI+SA_MYC_NS plants a higher number of DEGs was identified (6455 DEGs; 3400 up-regulated, 3055 down-regulated) with respect to CTRL_MYC_NS and CHI+SA_NMYC_NS (Fig. 5A, Supplementary Dataset S19). Transcripts involved in cell-wall metabolism, defence and transport were up-regulated in CHI+SA_MYC_NS

(Supplementary Dataset S19). The most up-regulated gene putatively encoded a O-acyltransferase WSD1-like (LOC101248922, $\text{Log}_2\text{FC}=9.42$) involved in wax metabolism.

Upon WS, the combined effect of SA priming and AM fungal inoculation on leaf transcriptome resulted in the up-regulation of several genes encoding heat shock proteins (36), abscisic acid 8'-hydroxylases (2), proteins putatively related to defence/resistance (23), ethylene-responsive TFs (8), and several PPR-containing proteins (198). Among the 47 genes exclusively up-regulated in SA_MYC_WS (Fig. 6A), the most expressed encoded a gibberellin 2-beta-dioxygenase ($\text{Log}_2\text{FC}=6.16$) and a dehydration-responsive element-binding protein ($\text{Log}_2\text{FC}=2.28$; Supplementary Dataset S27). Similarly, SA_MYC_SS plants showed up-regulation of transcripts putatively related to defence and resistance (23), PPR-containing proteins (218), heat shock proteins (13), and ethylene-responsive TFs (18). The most exclusively up-regulated gene encoded a feruloyl CoA ortho-hydroxylase ($\text{Log}_2\text{FC}=7.44$), linked to the phenylpropanoid pathway (Sun *et al.*, 2015; Supplementary Dataset S28).

In CHI+SA_MYC_WS, in addition to the up-regulation of genes encoding several PPR-containing proteins (190) and heat shock proteins (27), two genes related to ethylene-responsive TFs were strongly up-regulated (LOC101268109, $\text{Log}_2\text{FC}=11.57$ and LOC101246484, $\text{Log}_2\text{FC}=5.46$) together with genes encoding an ABA hydroxylase (LOC101249565) and an abscisic-aldehyde oxidase (AO1). An ABA 8'-hydroxylase was found among the first 20 most up-regulated genes in this sample (LOC101254720, $\text{Log}_2\text{FC}=6.45$; Supplementary Dataset S13). The strong up-regulation of a gene coding for an ascorbate-specific transmembrane electron transporter (LOC101259871, $\text{Log}_2\text{FC}=6.25$) in CHI+SA_MYC_WS, as well as in SA_MYC_WS ($\text{Log}_2\text{FC}=5.69$), was also observed. Compared to the other tested conditions, 38 up- and 45 down-regulated DEGs were exclusively identified in CHI+SA_MYC_WS plants (Fig. 6A; Supplementary Dataset S29). Looking at CHI+SA_MYC_SS, the up-regulation of genes encoding heat shock proteins-related (35), ethylene-responsive TFs (22), and proteins associated to defence response (26) were detected. Among the genes exclusively up-regulated in CHI+SA_MYC_SS leaves, a glutathione-S-transferase gene (LOC101266443; Supplementary Dataset S30).

A look into whole leaf metabolite profiling

Elaboration of metabolite profiling data revealed 14'516 chromatographic mass features (m.f.) differentially expressed in response to the various conditions (Supplementary Datasets S31-S32; see supporting datasets available at the Dryad data repository). Among this m.f. dataset, 372 chromatographic masses were putatively annotated to chemically different compounds (Supplementary Datasets S33 and S34). Supervised bi-dimensional analysis (2D-PLSDs) of these 372 leaf chromatographic masses, which were potentially among the most significant, showed a clear separation among the different stress conditions (NS, WS, SS; Fig. 7A). Furthermore, AM fungal inoculation had an impact under both stressed and NS conditions, as demonstrated by output of 2D-PLSDs (Fig. 7B). Out of the 372 detected chromatographic masses, 230 individual metabolites were annotated (Supplementary Dataset S35), and 64 of those were further selected as they exhibited the highest variability in the accumulation intensity ($\text{Log}_2\text{FC} > 1$ compared to CTRL_NMYC_NS) in at least one condition (Supplementary Dataset S36). Regardless of AM fungal inoculation and seed priming treatments, cluster analysis of the 64 selected metabolites highlighted that WS and SS had a significant impact on their modulation, leading to the identification of three distinct clusters (Fig. 7C). Cluster I included metabolites that were mainly accumulated under both stress conditions; cluster II grouped metabolites more abundant in response to salinity; cluster III was characterized by metabolites showing higher concentrations under NS and WS conditions. During WS, MYC plants accumulated more of those compounds compared to NMYC plants (Fig. 7C).

A closer look to the distribution of single metabolites within each cluster revealed that in cluster I, in absence of AM fungal inoculation, the combined impact of seed priming (SA and CHI+SA) and WS resulted in a higher accumulation of primary metabolites like fructose, ascorbic acid, pyruvic acid, arginine, aspartate, and of secondary metabolites, including fumaric acid, t-aconitic acid, quinic acid, benzoic acid, quercetin hexose, 12-oxo-phytodienoic acid (12-OPDA), 1-caffeoylquinic acid. Metabolites included in cluster II were mainly influenced by SS conditions and among the most abundant there were prodelfinidin A1, glutathione (GSH), amino adipic acid, pyroglutamic acid, L-aspartate semialdehyde, caffeoyl-glucaric acid and proline. Notably, proline displayed accumulation patterns consistent with those obtained through the target analytical approach (Fig. 7C). The interaction between AM fungal inoculation and WS positively influenced most of the metabolites grouped in cluster III, which comprised coumarin, mevalonic acid,

physalin L, jasmonil isoleucine, tyrosine, tangeretin, p-benzoquinones, quercetin deoxyhexose pentose, 3-hydroxyproline, quercetin hexose, feruloyl flavonoid glycoside, beta-tomatidine (Fig. 7C).

Integration between transcriptomic and metabolomic data reveals overlapping and distinct metabolic response to WS and SS in tomato plants in presence of seed priming agents and AM fungal inoculation

The DIABLO framework allowed the identification of potential interactions between transcripts and metabolites in the different tested conditions. The arrow plot from multiblock sPLS-DA revealed distinct clusters for NS, WS, and SS conditions, with the exception of CHI+SA_MYC_NS (Fig. 8). Short arrows under NS conditions indicated a close correlation between RNA-seq and metabolomic data for both MYC and NMYC, while relatively longer arrows were observed for the CTRL_NMYC_SS, SA_NMYC_SS, and CHI+SA_NMYC_SS conditions, suggesting less consistency between the two datasets (Fig. 8). Overall, the obtained output showed that, upon SS, the combination of AM fungal inoculation and chemical seed priming treatments functioned distinctly than all other conditions in regulating the plant metabolic response. Conversely, under WS the main distinction was noticed between untreated (CTRL) and treated (CHI+SA and SA) plants, independently of AM fungal inoculation (Fig. 8). Moreover, a network analysis conducted on the same datasets highlighted that overlapping and specific metabolic routes were regulated in AMF-inoculated plants treated with priming agents depending on WS or SS (Supplementary Figs. S7, S8). Particularly, the activation of transcripts and corresponding secondary metabolites involved in defense processes (such as tomatine, tomatidine and flavonols) and in lipid signaling (*e.g.* diacylglycerol and phosphatidylglycerol) represented a response common to the two abiotic stresses (Supplementary Figs. S7, S8). Nonetheless, signals mainly dependent on either WS or SS were also observed: this was the case of the activation of genes/metabolites with a strong antioxidant role (*e.g.* tangeretin, kaempferol, physalin) or involved in cell homeostasis regulation and signal transduction, including hormone-signaling, upon WS (Supplementary Fig. S7), and of specific sugar signaling routes at the basis of osmotic response upon SS (such as genes encoding sugar transporters, sugar biosynthetic enzymes, photosynthesis-related proteins, in association with osmoprotectant solutes, like proline; Supplementary Fig. S8).

Discussion

Priming of plants with chemical or biological agents has received increasing attention in the last years (Gohari *et al.*, 2024; Savvides *et al.*, 2016). It was already shown that the combined use of different agents, potentially acting synergistically, is a promising strategy for stimulating stress tolerance in plants (Balestrini *et al.*, 2018; González Guzmán *et al.*, 2022; Gohari *et al.*, 2024). In this framework, we investigated the effectiveness of the potential beneficial effects of chemical seed treatments (*i.e.* SA or CHI+SA) alone or in combination with post-germination AM fungal inoculation. The evaluation of seed priming on the ability of root colonization by AMF highlighted that the mycorrhization varied based on the interaction between stress conditions and treatments. Arbuscule abundance was not significantly affected by the tested variables, suggesting that the seed priming treatments did not deeply affect the AM fungal ability to colonize the root, at least considering these tomato and fungal genotypes.

Chemical seed priming and AM fungal inoculation modify the tomato plant response to water deficit and salinity

Overall, our results showed that the interplay between seed priming and AM fungal inoculation positively modulated plant stress responses to the imposed stresses. During WS, primed AMF-inoculated plants were able to maintain higher stomatal conductance and transpiration rates in comparison with plants subjected to either seed priming or inoculation alone. These plants also exhibited a lower water loss over the experimental trial, in agreement with previous studies demonstrating that SA and CHI improved water stress tolerance in crop species (Hidangmayum *et al.*, 2023a, b; Das *et al.*, 2023). The impact of AM symbiosis on plant performance in conditions of water limitation was extensively studied (Balestrini *et al.*, 2017), showing an improvement of tolerance for different plant species, including tomato (Chitarra *et al.*, 2016; Ruiz-Lozano *et al.*, 2016; Volpe *et al.*, 2018; Rivero *et al.*, 2018). The novel aspect here is that the synergistic effect of seed priming and subsequent AM fungal colonization further boosts the plant's ability to endure water deprivation. The collected data further suggested that the combination of CHI+SA and AM fungal inoculation was the most effective one in providing a long-term protective effect against water deficit (as attested by measurement of g_s and Ψ_{stem} values at the end of the stress imposition), whereas chemical priming alone could ultimately lead to detrimental effect (at least for some physiological parameters, as it was the case of Ψ_{stem} in SA_NMYC_WS plants). Although the

effect of exogenous application of chitosan and salicylic acid on stomata closure and photosynthetic processes upon water stress is well documented ([Hidangmayum et al., 2019](#); [Hidangmayum et al., 2023a,b](#); Janda *et al.*, 2014; Gao *et al.*, 2023; Silva *et al.*, 2023), available literature investigates short-term (over days) effects of these compounds in either seedlings or leaves of treated plants. Moreover, most information deals with foliar application rather than seed priming treatment, the latter representing a more recent research field (Gohari *et al.*, 2024). Our findings provide novel insights into this subject, showing that long-lasting (2 months after sowing the primed seeds) priming defence responses could also be established, resulting in improved physiological adaptation to water shortage conditions.

Notably, physiological adjustments also occurred when tomato plants were exposed to salinity (SS). AM symbiosis alone improved the plant capacity to counteract SS effects in terms of stomatal limitation, as CTRL_MYC plants did not experience the same steep drop in g_s and E of CTRL_NMYC. This is in agreement with the key role of AMF in the regulation of tolerance mechanisms to salt and osmotic stress reported in several plant species (Porcel *et al.*, 2012; Pollastri *et al.* 2018; Chandra *et al.*, 2023). Unlike to WS, during SS exposure, the most important effect of the combined treatment resulted in a balance of the plant physiological responses, being gas exchange rates similar between NS and SS conditions. Already in absence of stress application, stomata opening of CTR_MYC and SA and CHI+SA_MYC plants was significantly reduced with respect to NMYC plants, in turn resulting in a slowdown of photosynthesis. Additionally, seed primed NMYC and MYC plants were smaller than the untreated NMYC ones, indicating that the improved adaptation to stress encompassed phenotypic modifications able to alter the physiological performances (He *et al.*, 2022). Our results also pointed to an AMF-mediated balancing effect in the growth-defence trade-off mechanisms, which persisted with both seed priming treatments.

Overlapping and distinct transcriptomic and metabolic signatures are induced in stressed plants following treatments with seed priming agents and AM fungal inoculation

It is long-time known that plant tolerance to water deficit and salt stress involves profound changes in hormone metabolism and related signalling routes (Yu *et al.*, 2020; Salvi *et al.*, 2021). Here, besides a stress-dependent impact on the accumulation profiles of the main phytohormones, a significant effect of AM symbiosis, alone or in combination with both SA and CHI+SA treatments,

revealed reversed response with respect to untreated NMYC plants. Both SA and JA levels showed an increasing trend in SA and CHI+SA_MYC plants upon both WS and SS. It was already reported that seed priming with SA increased endogenous hormone content, including SA and ABA in cotton (Fang *et al.*, 2018). Additionally, proline accumulation was more accentuated in NMYC plants following both stress conditions, reflecting the well-known role of this solute in protecting cells from oxidative damages caused by osmotic stresses (Yoshida *et al.*, 1997; Forlani *et al.*, 2019). Proline concentration patterns agreed with expression changes of genes involved in proline metabolism. Particularly, *P5CS* (LOC101244293), a gene encoding the delta-1-pyrroline-5-carboxylate synthetase, catalyzing the first step of proline biosynthesis (Yoshida *et al.*, 1997), was up-regulated in NMYC plants. CHI+SA combined with AM fungal inoculation led to significantly higher levels of the metabolite already in NS conditions, suggesting the establishment of specific basal defence mechanisms shaping the plant stress perception. This result, coupled with the physiological modifications, could support a prompter ability of these plants in limiting cell water loss during stress through osmotic adjustment (Bartlett *et al.*, 2014; Forlani *et al.*, 2019).

At the transcriptomic level, regardless of inoculation or priming treatment, water deprivation caused a reprogramming of genes typically associated with stress responses, such as the dehydrin-encoding genes *ED14* and *TAS14*, and several heat shock proteins-related genes, as previously described (Iovieno *et al.*, 2016). ERD14 was reported in Arabidopsis to help plants coping with oxidative stress under dehydration (Nguyen *et al.*, 2020), while *TAS14* was already associated with WS response in tomato (Chitarra *et al.*, 2016) and it is considered one of the main tomato drought markers, transcriptionally activated by ABA signaling (Muñoz-Mayor *et al.*, 2012). Among the genes exclusively up-regulated upon WS, two genes encoding an abscisic stress-ripening protein 1 (*ASR1*) and a jasmonate o-methyltransferase-like were found. Tomato plants overexpressing *ASR1* showed enhanced survival rates under WS (Golan *et al.*, 2014), while up-regulation of a jasmonate o-methyltransferase in rice allowed the plants to produce MeJA in drought conditions (Kim *et al.*, 2009). The average expression value of *ASR1* in MYC plants subjected to WS was lower than in the NMYC ones, suggesting a role of AMF in mediating the ABA production, as previously reported (Begum *et al.*, 2020).

Salinity also caused a profound leaf transcriptomic remodelling, independently of priming treatments. The significant up-regulation of genes related to ion transport and stress response likely played a role in maintaining cell homeostasis and preventing sodium toxicity. Among the

transcripts exclusively regulated under SS, a gene encoding a suppressor of max2-like protein (*SMXL2*) was identified. Genes from the SMXL family, which are recognized as downstream components of the strigolactone (SL) signalling pathway, were proven to control salt stress responses in soybean and chickpea (Fu *et al.*, 2022; Basso *et al.*, 2024).

Overall, under the same growth conditions (*i.e.*, not-stressed, water deficit or high salinity), and irrespective of the seed priming treatment, AMF-inoculated tomato plants showed a distinct transcriptomic profile compared to not-inoculated. In NS conditions, the systemic foliar response to AM fungal inoculation was comparable to that reported by Cervantes *et al.* (2016), revealing the expression of genes already reported as key signature of AM fungal colonization (Liu *et al.*, 2007; López-Ráez *et al.*, 2010; Takeda *et al.*, 2011). In stress conditions, MYC plants showed up-regulation of a large set of PPR-containing protein genes, involved in RNA processing and in various biological processes, including response to ABA signal under stress conditions (Yuan and Liu, 2012). In *Populus alba*, transcription of *PPR* genes was induced following AM fungal inoculation in plants both grown on heavy metal-contaminated soils and upon optimal conditions (Cicatelli *et al.*, 2014); in *Citrus sinensis*, one of the most highly expressed genes in mycorrhizal plants encoded a PPR protein (Gao *et al.*, 2016). Our results further confirm a role for these proteins in the regulation of response to abiotic stresses in mycorrhizal plants.

In CHI+SA_MYC_WS plants, the transcriptome analysis highlighted a significant up-regulation of genes involved in ABA metabolism, such as an abscisic acid hydroxylase and an abscisic-aldehyde oxidase, which are considered integral to the ABA signalling pathway (Merilo *et al.*, 2015). A reduction in the expression level of an abscisic stress-ripening protein 1 (*ASR1*) gene in CHI+SA_MYC_WS was detected. This gene was previously found to be up-regulated in tomato leaves following WS treatment (Maskin *et al.*, 2001), so it could be argued that the decreased expression in presence of AM fungal inoculation and priming agents might be correlated with a less stressed status of the plants. In parallel, upon SS, CHI+SA_MYC plants increased the expression of a glutathione-S-transferase-encoding gene, involved in cell detoxification processes (Xu *et al.*, 2015). Together, these findings might indicate that the interaction between AM fungal inoculation and the CHI+SA-seed treatment not only triggers general stress response mechanisms, but also activates specific defence pathways. These results also imply that effects of the seed priming treatment were still effective after two months from the sowing.

In agreement with the observed transcriptomic changes, analysis of whole leaf metabolic signatures outlined an extensive remodelling of primary and secondary metabolite profiles primarily controlled by the stress condition or the AM fungal inoculation. Hierarchical clustering of metabolomic data attested that soluble sugars (like fructose and sucrose), osmoprotectant solutes (including proline), primary and secondary metabolites with a known antioxidant and defence role (such as glutathione, ascorbic acid and flavonols, flavanols and several precursors of the phenylpropanoid biosynthetic pathway) were the metabolic classes most differentially accumulated based on the three tested variables (stress, AM symbiosis and seed priming treatments). The increased accumulation of antioxidant metabolites, such as ascorbic acid (Zhou *et al.*, 2016), was supported at the molecular level by the positive modulation of specific genes, such as an ascorbate-specific transmembrane electron transporter and a glutathione-S-transferase, which were induced in the same samples. Secondary metabolites were mainly affected by presence of AMF. In leaves, metabolomic data demonstrated an enhanced content of the alkaloid beta-tomatidine in plants inoculated with AMF, irrespective of seed priming, especially under WS conditions. It has been observed that the concentration of tomatidine in tomato leaves substantially increases under water scarcity and when NaCl is added, suggesting that this compound is involved in chemical defense mechanisms (Bailly, 2021). The systemic accumulation of alkaloid compounds in tomato after AM fungal inoculation has also been documented, and it has been proposed that this class of metabolites may elicit plant metabolic responses to herbivore attack (Rivero *et al.*, 2021). This suggests that AMF may mediate the systemic accumulation of tomatidine in plants subjected to WS and modulate the related stress responses. Furthermore, several flavonoids and related compounds, including tangeretin, feruloyl flavonoid glycoside, and quercetin, were also accumulated in AMF-inoculated plants under water deficit. Flavonoids are well-known antioxidant compounds that play a protective role against a plethora of abiotic stresses, such as heat stress, salinity and drought (Hichri *et al.*, 2011; Martinez *et al.* 2016). High concentration of flavonoids in MYC WS plants has already been documented (Xie *et al.*, 2023), including tomato (Aseel *et al.*, 2019). Both AMF presence and water deficit affected the flavonoid content in tomato, suggesting that AMF may regulate the biosynthesis of these antioxidant molecules under stress conditions. A similar pattern was observed for coumarin, another antioxidant compound exerting a role in the activation of tomato tolerance to environmental stresses (Bourgau *et al.*, 2006). Notably, recent studies have reported that beneficial

microorganisms such as AMF can enhance coumarin production in tomato leaves (Iula *et al.*, 2021). Our transcriptomic data indicated an up-regulation of a 4-coumarate-CoA ligase-like 5 related gene in AMF-inoculated plants (both treated and untreated with priming agents) under water stress. This gene family includes multiple members and encodes several enzymes involved in the biosynthesis of hydroxycinnamate CoA esters, a critical step in diverting general phenylpropanoid metabolism toward coumarin biosynthesis (Hamberger and Halbrock, 2004). Particularly, the 4-coumarate-CoA ligase-like 5 gene was up-regulated in roots of soybean colonized by AMF, suggesting an involvement of AMF in fine tuning the expression of this gene (Marquez *et al.*, 2018). Here, a link between the accumulation of coumarin metabolites and the induction of this gene in AMF-inoculated plants was observed under water deprivation, likely indicating that an AMF-mediated systemic response occurred in tomato leaves.

About the effect of priming agents on leaf metabolomic profiles, some secondary metabolites including a flavonoid-related one, *i.e.*, quercetin-hexose, and an oxylipin, *i.e.*, 12-OPDA, were more abundant in SA and CHI+SA-primed plants, particularly under water deficit. The observed accumulation of quercetin-hexose, also observed in WS AMF-inoculated plants, further supports the hypothesis that priming agents may enhance the accumulation of antioxidant compounds (Król *et al.*, 2015). Concerning 12-OPDA, in Syrian bean capper plants (*Zygophyllum fabago*) pretreated with SA and exposed to different levels of lead, an increased accumulation of 12-OPDA was observed in the leaves, accompanied by an inhibition of the accumulation of the JA derivative Jasmonoyl-Isoleucine (JA-Ile; López-Orenes *et al.*, 2020). Plants with elevated levels of 12-OPDA were demonstrated to display an improved drought resistance associated with a reduction in stomatal aperture; moreover, it has been reported that drought interrupts the conversion of 12-OPDA to JA (Savchenko *et al.*, 2014). Similarly, in our study, tomato plants subjected to WS and primed with SA and CHI+SA showed an increased amount of 12-OPDA, whereas JA accumulated less in stressed than not-stressed plants. Interestingly, it has also been suggested that 12-OPDA acts in combination with ABA to elicit stomatal closure in response to drought (Savchenko *et al.*, 2014). Given the higher ABA levels in primed plants under WS compared to their untreated counterparts, it could be hypothesized that stomatal closure was driven by a priming elicitation of an interplay between ABA and 12-OPDA, as previously suggested (Savchenko *et al.*, 2014).

Collectively, our achievements suggest that seed treatments and AM fungal inoculation, alone or in combination, stimulate the tomato antioxidant and osmoprotective machinery, leading to an overall improvement of the plant physiological performance upon different abiotic stress conditions. It was furthermore shown that, besides overlapping defence responses (*e.g.* lipid signalling pathways), tomato plants subjected to chemical seed priming and AM fungal inoculation were able to activate molecular and metabolic responses specific of the type of stress. In fact, while defence signals were primarily represented by differential expression of genes linked to antioxidant secondary metabolites upon WS, the transcriptional reprogramming of defence processes is shifted towards osmotic stress responses upon SS. Elicitation of multiple defence-associated metabolic routes is therefore at the basis of priming, confirming the success of the adopted approaches in enhancing plant stress tolerance.

Accepted Manuscript

Acknowledgements

Mass spectrometric measurements were performed at Universitat Jaume I Servei Central d'Instrumentació Científica (SCIC) with the help of Dr. Cristian Vicent.

Author contribution

VA, MGG, VF and RB conceived and designed the research. LG, CP, EC, FS, SDR, EZ, WC, AI, AS, FV conducted all measurements and wet lab experiments. LG, CP, FS and WC conducted data analyses. LG, CP, FS and RB wrote the first draft of the manuscript. All authors read and approved the manuscript.

Conflict of interest statement

Authors have no conflict of interest declared.

Funding

This work and Financial support was provided through projects OPTIMUS PRIME (Italian Ministry of University and Research, grant DD 16302 12/11/2021, and Spanish Ministry of Science and Innovation, grant PCI2021-121920 funded by MCIN/AEI/10.13039/501100011033) and by the “European Union NextGenerationEU/PRTR”, EXTREMO grant PID2020-118126RB-I00 funded by MCIN/AEI/10.13039/501100011033, UJI-B2019-24 and UJI-B2022-23 funded by Universitat Jaume I. LG and EC's fellowships were funded by the PRIMA project OPTIMUS PRIME (Italian Ministry of University and Research DD 16302, 12/11/2021 and Spanish Ministry of Science and Innovation PCI2021-121920, respectively).

Data Availability

The RNA-Seq raw data underlying this article are available in the NCBI Sequence Read Archive (SRA) at <https://www.ncbi.nlm.nih.gov/sra> and can be accessed with the BioProject accession number PRJNA1108677.

The statistics and elaboration of RNAseq and untargeted metabolomics data (named as Supplementary Datasets from S1 to S36) that support the findings of this study are openly available in Dryad at the following link <https://doi.org/10.5061/dryad.dbrv15fb9>.

References

- Alagna F, Balestrini R, Chitarra W, Marsico AD, Nerva L.** 2020. Getting ready with the priming: innovative weapons against biotic and abiotic crop enemies in a global changing scenario. In: Hossain MA, Liu F, Burritt DJ, Fujita M, Huang B, eds, Priming-mediated stress and cross-stress tolerance in crop plants. New York, US: Academic Press, 35-56.
- Anders S, Pyl PT, Huber W.** 2015. HTSeq-a Python framework to work with high-throughput sequencing data. *Bioinformatics* **31**, 166-169.
- Andrade A, Boero A, Escalante M, Llanes A, Arbona V, Gómez-Cádenas A, Alemano S.** 2021. Comparative hormonal and metabolic profile analysis based on mass spectrometry provides information on the regulation of water-deficit stress response of sunflower (*Helianthus annuus* L.) inbred lines with different water-deficit stress sensitivity. *Plant Physiology and Biochemistry* **168**, 432-446.
- Aseel DG, Rashad YM, Hammad SM.** 2019. Arbuscular mycorrhizal fungi trigger transcriptional expression of flavonoid and chlorogenic acid biosynthetic pathways genes in tomato against Tomato Mosaic Virus. *Scientific Reports* **9**, 9692.
- Augé RM.** 2004. Arbuscular mycorrhizae and soil/plant water relations. *Can J Soil Sc.* **84**: 373-381.
- Bailly C.** 2021. The steroidal alkaloids α -tomatine and tomatidine: Panorama of their mode of action and pharmacological properties. *Steroids* **176**, 108933.
- Balestrini R, Brunetti C, Cammareri M, et al.** 2021. Strategies to modulate specialized metabolism in Mediterranean crops: From molecular aspects to field. *International Journal of Molecular Sciences* **22**, 2887.
- Balestrini R, Rosso LC, Veronico P, Melillo MT, De Luca F, Fanelli E, Colagiero M, Salvioli di Fossalunga A, Pentimone, I.** 2019. Transcriptomic responses to water deficit and nematode infection in mycorrhizal tomato roots. *Frontiers in Microbiology* **10**, 47441.
- Balestrini R, Chitarra W, Antoniou C, Ruocco M, Fotopoulos V.** 2018. Improvement of plant performance under water deficit with the employment of biological and chemical priming agents. *The Journal of Agricultural Science* **156**, 680-688.
- Balestrini R, Lumini E.** 2018. Focus on mycorrhizal symbioses. *Applied Soil Ecology* **123**, 299-304.
- Balestrini R, Chitarra W, Fotopoulos V, Ruocco M.** 2017. Potential role of beneficial soil microorganisms in plant tolerance to abiotic stress. In: Lukac M, Gamboni M, Grenni P, eds. Soil biological communities and ecosystem resilience. Sustainability in plant and crop protection series. Cham, Switzerland: Springer, 191-207.
- Balusamy SR, Rahimi S, Sukweenadhi J, Sunderraj S, Shanmugam R, Thangavelu L, Perumalsamy H.** 2022. Chitosan, chitosan nanoparticles and modified chitosan biomaterials, a potential tool to combat salinity stress in plants. *Carbohydrate Polymers* **284**, 119189.
- Bartlett MK, Zhang Y, Kreidler N, Sun S, Ardy R, Cao K, Sack L.** 2014. Global analysis of plasticity in turgor loss point, a key drought tolerance trait. *Ecology Letters* **17**, 1580-1590.

- Basso MF, Contaldi F, Lo Celso F, Baratto CM, Grossi-de-Sa MF, Barone G, Ferrante A, Martinelli F** (2024) Identification and expression profile of the SMAX/SMXL family genes in chickpea and lentil provide important players of biotechnological interest involved in plant branching. *Planta* **259**,1.
- Bastías DA, Balestrini R, Pollmann S, Gundel PE.** 2022. Environmental interference of plant-microbe interactions. *Plant, Cell and Environment* **45**, 3387-3398.
- Bates LS, Waldren RP, Teare ID.** 1973. Rapid determination of free proline for water stress studies. *Plant Soil* **39**, 205-207.
- Begum N, Ahanger MA, Zhang L.** 2020. AMF inoculation and phosphorus supplementation alleviates drought induced growth and photosynthetic decline in *Nicotiana tabacum* by up-regulating antioxidant metabolism and osmolyte accumulation. *Environmental and Experimental Botany* **176**, 104088.
- Bourgaud F, Hehn A, Lariat R, Doerper S, Gontier E, Kellner S, Matern U.** 2006. Biosynthesis of coumarins in plants: a major pathway still to be unravelled for cytochrome P450 enzymes. *Phytochemistry Reviews* **5**, 293-308.
- Cervantes-Gómez RG, Bueno-Ibarra MA, Cruz-Mendivil A, Calderón-Vázquez CL, Ramírez-Douriet CM, Maldonado-Mendoza IE, Villalobos-López MA, Valdez-Ortíz A, López-Meyer M.** 2016. Arbuscular mycorrhizal symbiosis-induced expression changes in *Solanum lycopersicum* leaves revealed by RNA-seq analysis. *Plant Molecular Biology Reporter* **34**, 89-102.
- Chakma R, Biswas A, Saekong P, Ullah H, Datta A.** 2021. Foliar application and seed priming of salicylic acid affect growth, fruit yield, and quality of grape tomato under drought stress. *Scientia Horticulturae* **280**, 109904.
- Chambers MC, Maclean B, Burke R, et al.** 2012. A cross-platform toolkit for mass spectrometry and proteomics. *Nature Biotechnology* **30**, 918-920.
- Chandra P, Yadav S, Kumar A, Sheoran P, Mann A.** 2023. Arbuscular mycorrhizal fungi–plant interaction for salinity and drought stress alleviation. In: Kumar A, Dhansu P, Mann A, eds. *Salinity and drought tolerance in plants: physiological perspectives*. Singapore, MY: Springer, 365-386.
- Chang S, Puryear J, Cairney J.** 1993. A simple and efficient method for isolating RNA from pine trees. *Plant Molecular Biology Reporter* **11**, 113-116.
- Chitarra W, Pagliarani C, Maserti B, Lumini E, Siciliano I, Cascone P, Schubert A, Gambino G, Balestrini R, Guerrieri E.** 2016. Insights on the impact of arbuscular mycorrhizal symbiosis on tomato tolerance to water stress. *Plant Physiology* **171**, 1009-1023.
- Cicatelli A, Todeschini V, Lingua G, Biondi S, Torrigiani P, Castiglione S.** 2014. Epigenetic control of heavy metal stress response in mycorrhizal versus non-mycorrhizal poplar plants. *Environmental Science and Pollution Research* **21**, 1723-1737.
- Conesa A, Götz S, García-Gómez JM, Terol J, Talón M, Robles M.** 2005. Blast2GO: a universal tool for annotation, visualization and analysis in functional genomics research. *Bioinformatics* **21**, 3674-3676.
- Cortina C, Culiáñez-Macià FA.** 2005. Tomato abiotic stress enhanced tolerance by trehalose biosynthesis. *Plant Science* **169**, 75-82.

- Cutler SR, Rodriguez PL, Finkelstein RR, Abrams SR.** 2010. Abscisic acid: emergence of a core signaling network. *Annual Review of Plant Biology* **61**, 651-679.
- Das D, Bisht K, Chauhan A, Gautam S, Jaiswal JP, Salvi P, Lohani P.** 2023. Morpho-physiological and biochemical responses in wheat foliar sprayed with zinc-chitosan-salicylic acid nanoparticles during drought stress. *Plant Nano Biology* **4**, 100034.
- De Ollas C, González-Guzmán M, Pitarch Z, Matus JT, Candela H, Rambla JL, Granell A, Gómez-Cadenas A, Arbona V.** 2021. Identification of ABA-mediated genetic and metabolic responses to soil flooding in tomato (*Solanum lycopersicum* L. Mill). *Frontiers in Plant Science* **12**, 613059.
- Del Fabbro C, Scalabrin S, Morgante M, Giorgi FM.** 2013. An extensive evaluation of read trimming effects on Illumina NGS data analysis. *PLoS One* **8**, e85024.
- Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras TR.** 2013. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21.
- Duc NH, Csintalan Z, Posta K.** 2018. Arbuscular mycorrhizal fungi mitigate negative effects of combined drought and heat stress on tomato plants. *Plant Physiology and Biochemistry* **132**, 297-307.
- Duc NH, Szentpéteri V, Mayer Z, Posta K.** 2023. Distinct impact of arbuscular mycorrhizal isolates on tomato plant tolerance to drought combined with chronic and acute heat stress. *Plant Physiology and Biochemistry* **201**, 107892.
- Fang S, Gao K, Hu W, Snider JL, Wang S, Chen B, Zhou Z.** 2018. Chemical priming of seed alters cotton floral bud differentiation by inducing changes in hormones, metabolites and gene expression. *Plant Physiology and Biochemistry* **130**, 633-640.
- Fang Y, Xiong L.** 2015. General mechanisms of drought response and their application in drought resistance improvement in plants. *Cellular and Molecular Life Sciences* **72**, 673-689.
- Fathi A, Tari DB.** 2016. Effect of drought stress and its mechanism in plants. *International Journal of Life Sciences* **10**, 1-6.
- Fonseca JP, Dong X.** 2014. Functional characterization of a Nudix hydrolase *AtNUDX8* upon pathogen attack indicates a positive role in plant immune responses. *PLoS One* **9**, e114119.
- Forlani G, Trovato M, Funck D, Signorelli S.** 2019. Regulation of Proline Accumulation and its Molecular and Physiological Functions in Stress Defence. In: Hossain M, Kumar V, Burritt D, Fujita M, Mäkelä P, eds. *Osmoprotectant-Mediated Abiotic Stress Tolerance in Plants*. Cham, Switzerland: Springer, 73-97.
- Francini A, Sebastiani L.** 2019. Abiotic stress effects on performance of horticultural crops. *Horticulturae* **5**, 67.
- Fróna D, Szenderák J, Harangi-Rákos M.** 2021. Economic effects of climate change on global agricultural production. *Nature Conservation* **44**, 117-139.
- Fu X, Wang J, Shangguan T, Wu R, Li S, Chen G, Xu S.** 2022. SMXLs regulate seed germination under salinity and drought stress in soybean. *Plant Growth Regulation* **96**, 397-408.

- Galmés J, Ochogavía JM, Gago J, Roldán EJ, Cifre J, Conesa MÀ.** 2013. Leaf responses to drought stress in Mediterranean accessions of *Solanum lycopersicum*: anatomical adaptations in relation to gas exchange parameters. *Plant, Cell and Environment* **36**, 920-935.
- Galviz-Fajardo YC, Bortolin GS, Deuner S, Amarante LD, Reolon F, Moraes DMD.** 2020. Seed priming with salicylic acid potentiates water restriction-induced effects in tomato seed germination and early seedling growth. *Journal of Seed Science* **42**, e202042031.
- Gao X, Zhao S, Xu QL, Xiao JX.** 2016. Transcriptome responses of grafted *Citrus sinensis* plants to inoculation with the arbuscular mycorrhizal fungus *Glomus versiforme*. *Trees* **30**, 1073-1082.
- Gao Q, Liu Y, Liu Y, Dai C, Zhang Y, Zhou F, Zhu Y.** 2023. Salicylic acid modulates the osmotic system and photosynthesis rate to enhance the drought tolerance of *Toona ciliata*. *Plants* **12**, 4187.
- Giannakoula A, Ilias I.** 2013. The effect of water stress and salinity on growth and physiology of tomato (*Lycopersicon esculentum* Mil.). *Archives of Biological Sciences* **65**, 611-620.
- Gohari G, Alavi Z, Esfandiari E, Panahirad S, Hajihoseinlou S, Fotopoulos V.** 2020. Interaction between hydrogen peroxide and sodium nitroprusside following chemical priming of *Ocimum basilicum* L. against salt stress. *Physiologia Plantarum* **168**, 361-373.
- Gohari G, Jiang M, Manganaris GA, Zhou J, Fotopoulos V.** 2024. Next generation chemical priming: with a little help from our nanocarrier friends. *Trends in Plant Science* **29**, 150-166.
- Golan I, Dominguez PG, Konrad Z, Shkolnik-Inbar D, Carrari F, Bar-Zvi D.** 2014. Tomato abscisic acid stress ripening (ASR) gene family revisited. *PLoS One* **9**, e107117.
- González Guzmán M, Cellini F, Fotopoulos V, Balestrini R, Arbona V.** 2022. New approaches to improve crop tolerance to biotic and abiotic stresses. *Physiologia Plantarum* **174**, e13547.
- Hamberger B, Hahlbrock K.** 2004. The 4-coumarate: CoA ligase gene family in *Arabidopsis thaliana* comprises one rare, sinapate-activating and three commonly occurring isoenzymes. *Proceedings of the National Academy of Sciences* **101**, 2209-2214.
- Hassanpour M, Jafari H, Sharifi S, Rezaie J, Lighvan ZM, Mahdavinia GR, Gohari G, Akbari A.** 2021. Salicylic acid-loaded chitosan nanoparticles (SA/CTS NPs) for breast cancer targeting: Synthesis, characterization and controlled release kinetics. *Journal of Molecular Structure* **1245**, 131040.
- Hichri I, Barrieu F, Bogs J, Kappel C, Delrot S, Lauvergeat V.** 2011. Recent advances in the transcriptional regulation of the flavonoid biosynthetic pathway. *Journal of Experimental Botany* **62**, 2465-2483.
- Hidangmayum A, Dwivedi P, Katiyar D, Hemantaranjan A.** 2019. Application of chitosan on plant responses with special reference to abiotic stress. *Physiology and Molecular Biology of Plants* **25**, 313-326.
- Hidangmayum A, Dwivedi P, Kumar P, Upadhyay SK.** 2023a. Seed priming and foliar application of chitosan ameliorate drought stress responses in mungbean genotypes through modulation of morpho-physiological attributes and increased antioxidative defense mechanism. *Journal of Plant Growth Regulation* **42**, 6137-6154.
- Hidangmayum A, Dwivedi P.** 2023b. Effect of chitosan seed priming on mungbean seedlings subjected to different levels of water potential. *Acta Physiologia Plantarum* **45**, 6.

- Hilker M, Schmölling T.** 2019. Stress priming, memory, and signalling in plants. *Plant, Cell and Environment* **42**, 753-761.
- Hoagland DR, Broyer TC.** 1936. General nature of the process of salt accumulation by roots with description of experimental methods. *Plant Physiology* **11**, 471-507.
- Horton RM, Mankin JS, Lesk C, Coffel E, Raymond C.** 2016. A Review of recent advances in research on extreme heat events. *Current Climate Change Reports* **2**, 242-259.
- Hosmani PS, Flores-Gonzalez M, van de Geest H, et al.** 2019. An improved de novo assembly and annotation of the tomato reference genome using single-molecule sequencing, Hi-C proximity ligation and optical maps. *BioRxiv* doi: <https://doi.org/10.1101/767764> [Preprint].
- Hu H, Xiong L.** 2014. Genetic engineering and breeding of drought-resistant crops. *Annual Review of Plant Biology* **65**, 715-741.
- Iovieno P, Punzo P, Guida G, et al.** 2016. Transcriptomic changes drive physiological responses to progressive drought stress and rehydration in tomato. *Frontiers in Plant Science* **7**, 371.
- Iriti M, Varoni EM.** 2015. Chitosan-induced antiviral activity and innate immunity in plants. *Environmental Science and Pollution Research* **22**, 2935-2944.
- Iula G, Miras-Moreno B, Lucini L, Trevisan M.** 2021. The mycorrhiza- and Trichoderma-mediated elicitation of secondary metabolism and modulation of phytohormone profile in tomato plants. *Horticulturae* **7**, 394.
- Kim EH, Kim YS, Park SH, Koo YJ, Choi YD, Chung YY, Lee IJ, Kim JK.** 2009. Methyl jasmonate reduces grain yield by mediating stress signals to alter spikelet development in rice. *Plant Physiology* **149**, 1751-1760.
- Król P, Igielski R, Pollmann S, Kępczyńska E.** 2015. Priming of seeds with methyl jasmonate induced resistance to hemi-biotroph *Fusarium oxysporum* f. sp. *lycopersici* in tomato via 12-oxo-phytodienoic acid, salicylic acid, and flavonol accumulation. *Journal of Plant Physiology* **179**, 122-132.
- Kuhl C, Tautenhahn R, Böttcher C, Larson TR, Neumann S.** 2012. CAMERA: an integrated strategy for compound spectra extraction and annotation of liquid chromatography/mass spectrometry data sets. *Analytical Chemistry* **84**, 283-289.
- Kulak M, Jorrín-Novo JV, Romero-Rodriguez MC, Yildirim ED, Gul F, Karaman S.** 2021. Seed priming with salicylic acid on plant growth and essential oil composition in basil (*Ocimum basilicum* L.) plants grown under water stress conditions. *Industrial Crops and Products* **161**, 113235.
- Kumar A, Verma JP.** 2018. Does plant-microbe interaction confer stress tolerance in plants: A review? *Microbiological Research* **207**, 41-52.
- Janda T, Gondor OK, Yordanova R, Szalai G, Pál M.** 2014. Salicylic acid and photosynthesis: signalling and effects. *Acta Physiologiae Plantarum* **36**, 2537-2546.
- Lamers J, van der Meer T, Testerink C.** 2020. How plants sense and respond to stressful environments. *Plant Physiology* **182**, 1624-1635.
- Le Gall H, Philippe F, Domon JM, Gillet F, Pelloux J, Rayon C.** 2015. Cell wall metabolism in response to abiotic stress. *Plants* **4**, 112.

- Lesk C, Rowhani P, Ramankutty N.** 2016. Influence of extreme weather disasters on global crop production. *Nature* **529**, 84-87.
- Leventis G, Tsiknia M, Feka M., Ladikou EV, Papadakis IE, Chatzipavlidis I, Papadopoulou K, Ehaliotis C.** 2021. Arbuscular mycorrhizal fungi enhance growth of tomato under normal and drought conditions, via different water regulation mechanisms. *Rhizosphere* **19**, 100394.
- Li Y, Ye W, Wang M, Yan X.** 2009. Climate change and drought: a risk assessment of crop-yield impacts. *Climate Research* **39**, 31-46.
- Liang SM, Zhang F, Zou YN, Kuća K, Wu QS.** 2021. Metabolomics analysis reveals drought responses of trifoliolate orange by arbuscular mycorrhizal fungi with a focus on terpenoid profile. *Frontiers Plant Science* **12**, 740524.
- Liu J, Maldonado- Mendoza I, Lopez- Meyer M, Cheung F, Town CD, Harrison MJ.** 2007. Arbuscular mycorrhizal symbiosis is accompanied by local and systemic alterations in gene expression and an increase in disease resistance in the shoots. *The Plant Journal* **50**, 529-544.
- Liu MY, Li QS, Ding WY, et al.** 2023. Arbuscular mycorrhizal fungi inoculation impacts expression of aquaporins and salt overly sensitive genes and enhances tolerance of salt stress in tomato. *Chemical and Biological Technologies in Agriculture* **10**, 5.
- López-Orenes A, Alba JM, Kant MR, Calderón AA, Ferrer MA.** 2020. OPDA and ABA accumulation in Pb-stressed *Zygophyllum fabago* can be primed by salicylic acid and coincides with organ-specific differences in accumulation of phenolics. *Plant Physiology and Biochemistry* **154**, 612-621.
- López-Ráez JA, Verhage A, Fernández I, García JM, Azcón-Aguilar C, Flors V, Pozo MJ.** 2010. Hormonal and transcriptional profiles highlight common and differential host responses to arbuscular mycorrhizal fungi and the regulation of the oxylipin pathway. *Journal of Experimental Botany* **61**, 2589-2601.
- Love MI, Huber W, Anders S.** 2014. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biology* **15**, 550.
- Ma QJ, Sun MH, Lu J, Kang H, You CX, Hao YJ.** 2019. An apple sucrose transporter MdSUT2.2 is a phosphorylation target for protein kinase MdCIPK22 in response to drought. *Plant Biotechnology Journal* **17**, 625-637.
- Marquez N, Giachero ML, Gallou A, Debat HJ, Cranenbrouck S, Di Rienzo JA, Declerck S.** 2018. Transcriptional changes in mycorrhizal and nonmycorrhizal soybean plants upon infection with the fungal pathogen *Macrophomina phaseolina*. *Molecular Plant-Microbe Interactions* **31**, 842-855.
- Martinez V, Mestre TC, Rubio F, Girones-Vilaplana A, Moreno DA, Mittler R, Rivero RM.** 2016. Accumulation of flavonols over hydroxycinnamic acids favors oxidative damage protection under abiotic stress. *Frontiers in Plant Science* **7**, 838.
- Maskin L, Gudesblat GE, Moreno JE, Carrari FO, Frankel N, Sambade A, Rossi M, Iusem ND.** 2001. Differential expression of the members of the *Asr* gene family in tomato (*Lycopersicon esculentum*). *Plant Science* **161**, 739-746.

- Mauch-Mani B, Baccelli I, Luna E, Flors V.** 2017. Defense priming: an adaptive part of induced resistance. *Annual Review of Plant Biology* **68**, 485-512.
- Merilo E, Jalakas P, Laanemets K, Mohammadi O, Hõrak H, Kollist H, Brosché M.** 2015. Abscisic Acid Transport and Homeostasis in the Context of Stomatal Regulation. *Molecular Plant* **8**, 1321-1333.
- Moradi Tarnabi Z, Iranbakhsh A, Mehregan I, Ahmadvand R.** 2020. Impact of arbuscular mycorrhizal fungi (AMF) on gene expression of some cell wall and membrane elements of wheat (*Triticum aestivum* L.) under water deficit using transcriptome analysis. *Physiology and Molecular Biology of Plants* **26**, 143-162.
- Munns R, Gilliham M.** 2015. Salinity tolerance of crops - what is the cost? *New Phytologist* **208**, 668-673.
- Muñoz-Mayor A, Pineda B, Garcia-Abellán JO, et al.** 2012. Overexpression of dehydrin *tas14* gene improves the osmotic stress imposed by drought and salinity in tomato. *Journal of Plant Physiology* **169**, 459-468.
- Nerva L, Balestrini R, Chitarra W.** 2023. From plant nursery to field: persistence of mycorrhizal symbiosis balancing effects on growth-defence tradeoffs mediated by rootstock. *Agronomy* **13**, 229.
- Nerva L, Giudice G, Quiroga G, Belfiore N, Lovat L, Perria R, Volpe MG, Moffa L, Sandrini M, Gaiotti F.** 2022. Mycorrhizal symbiosis balances rootstock-mediated growth-defence tradeoffs. *Biology and Fertility of Soils* **58**, 17-34.
- Nguyen PN, Tossounian MA, Kovacs DS, et al.** 2020. Dehydrin ERD14 activates glutathione transferase Phi9 in *Arabidopsis thaliana* under osmotic stress. *Biochimica et Biophysica Acta* **1864**, 129506.
- Pagliarani C, Moine A, Chitarra W, Nerva L, Catoni M, Tavazza R, Matić S, Vallino M, Secchi F, Noris E.** 2022. The C4 protein of tomato yellow leaf curl Sardinia virus primes drought tolerance in tomato through morphological adjustments. *Horticulture Research* **9**, uhac164.
- Pascale A, Proietti S, Pantelides IS, Stringlis IA.** 2020. Modulation of the root microbiome by plant molecules: the basis for targeted disease suppression and plant growth promotion. *Frontiers in Plant Science* **10**, 1741.
- Pellegrino E, Bedini S, Avio L, Bonari E, Giovannetti M.** 2011. Field inoculation effectiveness of native and exotic arbuscular mycorrhizal fungi in a Mediterranean agricultural soil. *Soil Biology and Biochemistry* **43**, 367-376.
- Pollastri S, Savvides A, Pesando M, et al.** 2018. Impact of two arbuscular mycorrhizal fungi on *Arundo donax* L. response to salt stress. *Planta* **247**, 573-585.
- Porcel R, Aroca R, Ruíz-Lozano JM.** 2012. Salinity stress alleviation using arbuscular mycorrhizal fungi. A review. *Agronomy for Sustainable Development* **32**, 181-200.

- Porcel R, Ruiz-Lozano JM.** 2004. Arbuscular mycorrhizal influence on leaf water potential, solute accumulation and oxidative stress in soybean plants subjected to drought stress. *Journal of Experimental Botany* **55**, 1743-1750.
- Qadir M, Quill rou E, Nangia V, Murtaza G, Singh M, Thomas RJ, Drechsel P, Noble AD.** 2014. Economics of salt- induced land degradation and restoration. *Natural Resources Forum* **38**, 282-295.
- Recchia GH, Konzen ER, Cassieri F, Caldas DGG, Tsai SM.** 2018. Arbuscular mycorrhizal symbiosis leads to differential regulation of drought-responsive genes in tissue-specific root cells of common bean. *Frontiers in Microbiology* **9**, 1339.
- Rivas-San Vicente M, Plasencia J.** 2011. Salicylic acid beyond defence: its role in plant growth and development. *Journal of Experimental Botany* **62**, 3321-3338.
- Rivero J,  lvarez D, Flors V, Azc n-Aguilar C, Pozo MJ.** 2018. Root metabolic plasticity underlies functional diversity in mycorrhiza-enhanced stress tolerance in tomato. *New Phytologist* **220**, 1322-1336.
- Rivero J, Lidoy J, Llopis-Gim nez  , Herrero S, Flors V, Pozo MJ.** 2021. Mycorrhizal symbiosis primes the accumulation of antiherbivore compounds and enhances herbivore mortality in tomato. *Journal of Experimental Botany* **72**, 5038-5050.
- Ro ca M, Mihalache G, Stoleru V.** 2023. Tomato responses to salinity stress: From morphological traits to genetic changes. *Frontiers in Plant Science* **14**, 1118383.
- Rosenzweig C, Elliott J, Deryng D, et al.** 2014. Assessing agricultural risks of climate change in the 21st century in a global gridded crop model intercomparison. *Proceedings of the National Academy of Sciences of the USA* **111**, 3268-3273.
- Ruiz-Lozano JM, Aroca R, Zamarre o  M, Molina S, Andreo-Jim nez B, Porcel R, Garc a-Mina JM, Ruyter-Spira C, L pez-R ez JA.** 2016. Arbuscular mycorrhizal symbiosis induces strigolactone biosynthesis under drought and improves drought tolerance in lettuce and tomato. *Plant, Cell and Environment* **39**, 441-452.
- Salvi P, Manna M, Kaur H, Thakur T, Gandass N, Bhatt D, Muthamilarasan M.** 2021. Phytohormone signaling and crosstalk in regulating drought stress response in plants. *Plant Cell Reports* **40**, 1305-1329.
- S nchez-Rodr guez E, Moreno DA, Ferreres F, Rubio-Wilhelmi Mdel M, Ruiz JM.** 2011. Differential responses of five cherry tomato varieties to water stress: Changes on phenolic metabolites and related enzymes. *Phytochemistry* **72**, 723-729.
- S nchez-Romera B, Calvo-Polanco M, Ruiz-Lozano JM, Zamarre o  M, Arbona V, Garc a-Mina JM, G mez-Cadenas A, Aroca R.** 2018. Involvement of the def-1 mutation in the response of tomato plants to arbuscular mycorrhizal symbiosis under well-watered and drought conditions. *Plant and Cell Physiology* **59**, 248-261.

Santos MS, Nogueira MA, Hungria M. 2019. Microbial inoculants: reviewing the past, discussing the present and previewing an outstanding future for the use of beneficial bacteria in agriculture. *AMB Express* **9**, 205.

Sardans J, Urbina I, Grau O, Asensio D, Ogaya R, Peñuelas J. 2020. Long-term drought decreases ecosystem C and nutrient storage in a Mediterranean holm oak forest. *Environmental and Experimental Botany* **177**, 104135.

Savchenko T, Kolla VA, Wang CQ, et al. 2014. Functional convergence of oxylipin and abscisic acid pathways controls stomatal closure in response to drought. *Plant Physiology* **164**, 1151-1160.

Savvides A, Ali S, Tester M, Fotopoulos V. 2016. Chemical priming of plants against multiple abiotic stresses: mission possible? *Trends Plant Science* **21**, 329-340.

Schachtman DP, Goodger JQ. 2008. Chemical root to shoot signaling under drought. *Trends Plant Science* **13**, 281-287.

Shabnam N, Tripathi I, Sharmila P, Pardha-Saradhi P. 2016. A rapid, ideal, and eco-friendlier protocol for quantifying proline. *Protoplasma* **253**, 1577-1582.

Silva J M, da Silva Júnior GB, Bonifácio A, Dutra AF, de Mello Prado R, de Alcântara Neto F, Zuffo AM, Melo RS, de Sousa Pereira TL, de Sousa RS. 2023. Exogenous salicylic acid alleviates water stress in watermelon plants. *Annals of Applied Biology* **182**, 121–130.

Singh A, Shannon CP, Gautier B, Rohart F, Vacher M, Tebbutt SJ, Lê Cao KA. 2019. DIABLO: an integrative approach for identifying key molecular drivers from multi-omics assays. *Bioinformatics* **35**, 3055-3062.

Smith CA, Want EJ, O'Maille G, Abagyan R, Siuzdak G. 2006. XCMS: processing mass spectrometry data for metabolite profiling using nonlinear peak alignment, matching, and identification. *Analytical Chemistry* **78**, 779-787.

Smith SE, Read DJ. 2008. *Mycorrhizal Symbiosis*, 3rd edn. London, UK: Academic Press.

Spatafora JW, Chang Y, Benny GL, et al. 2016. A phylum-level phylogenetic classification of zygomycete fungi based on genome-scale data. *Mycologia* **108**, 1028-1046.

Sun X, Zhou D, Kandavelu P, Zhang H, Yuan Q, Wang BC, Rose J, Yan Y. 2015. Structural insights into substrate specificity of feruloyl-coa 6'-hydroxylase from *Arabidopsis thaliana*. *Scientific Reports* **5**, 1035.

Takeda N, Haage K, Sato S, Tabata S, Parniske M. 2011. Activation of a *Lotus japonicus* subtilase gene during arbuscular mycorrhiza is dependent on the common symbiosis genes and two cis-active promoter regions. *Molecular Plant-Microbe Interactions* **24**, 662-670.

Tripathi DK, Bhat JA, Antoniou C, Kandhol N, Singh VP, Fernie AR, Fotopoulos V. 2024. Redox regulation by priming agents towards a sustainable agriculture. *Plant and Cell Physiology* **65**, 1087-1102.

Trouvelot A, Kough JL, Gianinazzi-Pearson V. 1986. Mesure du taux de mycorrhization VA d'un système racinaire. Recherche de méthodes d'estimation ayant une signification fonctionnelle. In Gianinazzi-Pearson V, Gianinazzi S, eds. *Mycorrhizae: physiology and genetics*. Paris, France: INRA, 217-221.

Tugizimana F, Mhlongo M, Piater L, Dubery I. 2018. Metabolomics in plant priming research: The way forward? *International Journal of Molecular Sciences* **19**, 1759.

Ujvári G, Grassi A, Cristani C, Pagliarani I, Avio L, Blandino M, Capo L, Giovannetti M, Turrini A, Agnolucci M. 2023. Native mycorrhizal communities in maize roots as affected by plant genotype, starter fertilization and a seed-applied biostimulant. *Plant and Soil* **492**, 519-539.

Volpe V, Chitarra W, Cascone P, et al. 2018. The association with two different arbuscular mycorrhizal fungi differently affects water stress tolerance in tomato. *Frontiers in Plant Science* **9**, 1480.

Wang Y, Sun C, Ye Z, Li C, Huang S, Lin T. 2024. The genomic route to tomato breeding: Past, present, and future. *Plant Physiology* **30**, kiae248.

Xie W, Hao Z, Zhou J, Fu W, Guo L, Zhang X, Chen B. 2023. Integrated transcriptomics and metabolomics reveal specific phenolic and flavonoid accumulation in licorice (*Glycyrrhiza uralensis* Fisch.) induced by arbuscular mycorrhiza symbiosis under drought stress. *Plant Physiology and Biochemistry* **205**, 108173.

Xu J, Xing XJ, Tian YS, Peng RH, Xue Y, Zhao W, Yao QH. 2015. Transgenic arabidopsis plants expressing tomato glutathione s-transferase showed enhanced resistance to salt and drought stress. *PLoS One* **10**, e0136960.

Yoshida Y, Kiyosue T, Nakashima K, Yamaguchi-Shinozaki K, Shinozaki K. 1997. Regulation of levels of proline as an osmolyte in plants under water stress. *Plant and Cell Physiology* **38**, 1095-102.

Yu Z, Duan X, Luo L, Dai S, Ding Z, Xia G. 2020. How plant hormones mediate salt stress responses. *Trends in Plant Science* **25**, 1117-1130.

Yuan H, Liu D. 2012. Functional disruption of the pentatricopeptide protein SLG1 affects mitochondrial RNA editing, plant development, and responses to abiotic stresses in *Arabidopsis*. *The Plant Journal* **70**, 432-444.

Zhang W, Xia K, Feng Z, Qin Y, Zhou Y, Feng G, Zhu H, Yao Q. 2024. Tomato plant growth promotion and drought tolerance conferred by three arbuscular mycorrhizal fungi is mediated by lipid metabolism. *Plant Physiology and Biochemistry* **208**, 108478.

Zhang X, Han C, Gao H, Cao Y. 2019. Comparative transcriptome analysis of the garden asparagus (*Asparagus officinalis* L.) reveals the molecular mechanism for growth with arbuscular mycorrhizal fungi under salinity stress. *Plant Physiology and Biochemistry* **141**, 20-29.

Zhang Z, Chang X, Luo S, Wang Y, Xuan S, Zhao J, Chen X. 2023. Transcriptome analysis of two pepper genotypes infected with pepper mild mottle virus. *Frontiers in Genetics* **14**, 1164730.

Zhang Z, Liu J, Huber DJ, Qu H, Yun Z, Li T, Jiang Y. 2021. Transcriptome, degradome and physiological analysis provide new insights into the mechanism of inhibition of litchi fruit senescence by melatonin. *Plant Science* **308**, 110926.

Zhou X, Gu Z, Xu H, Chen L, Tao G, Yu Y, Li K. 2016. The effects of exogenous ascorbic acid on the mechanism of physiological and biochemical responses to nitrate uptake in two rice cultivars (*Oryza sativa* L.) under aluminum stress. *Journal of Plant Growth Regulation* **35**, 1013-1024.

Accepted Manuscript

Figure legends

Fig. 1

Arbuscular mycorrhizal fungal (AMF) colonization assessment in tomato potted plants (*Solanum lycopersicum* cv Moneymaker) inoculated with a commercial inoculum (MycAgro, *Funneliformis mosseae*). Top panels show representative images of AMF-inoculated roots. (A) F%, Frequency of mycorrhization in root system; bar (AI) = 120 μm ; (B) a%, arbuscule abundance in the colonized root system; bar (BI) = 60 μm , bar (BII) = 20 μm . All results are reported as mean \pm SE. Lower case letters above bars denote significant differences as assessed by Tukey HSD test ($p < 0.05$) and are reported only when the priming treatment (T) \times stress condition (S) interaction is significant. ns, *: not significant or significant at $p \leq 0.05$, respectively. CTRL: untreated, SA: treated with salicylic acid, CHI+SA: treated with chitosan + salicylic acid, NMYC: not-inoculated, MYC: inoculated, NS: not-stressed condition, WS: water stress condition, SS: salt stress condition. Analysis of variance on the single variables is reported in Supplementary Table S1.

Fig. 2

Effects of priming treatment and arbuscular mycorrhizal fungal (AMF) inoculation on the physiological performances of tomato (*Solanum lycopersicum* cv Moneymaker) potted plants during a water stress time-course. (A) Changes in soil relative water content (RWC_{soil} %) monitored over the whole duration of the water stress time-course. The inset beside panel A) magnifies differences in RWC_{soil} values among the diverse groups of plants during the first four days after water withdrawal (T1-T5). (B-D) Dynamic changes in the rates of (B) stomatal conductance (g_s , $\text{mol H}_2\text{O m}^{-2} \text{s}^{-1}$), (C) leaf vapour pressure deficit (VPD_{leaf} , KPa) and (D) leaf transpiration (E , $\text{mmol H}_2\text{O m}^{-2} \text{s}^{-1}$) monitored over the whole duration of the water stress time-course. (E) Rates of g_s measured at the end (Tf) of the experiment. All results are reported as mean \pm SE. In panel A), asterisks denote significant differences determined by one-way ANOVA on RWC_{soil} data (*, **, and *** significant at $p \leq 0.05$, $p \leq 0.01$, and $p \leq 0.001$, respectively, with corresponding values reported in the panel inset), while different lower case letters in the panel A) inset highlight significant differences among plant groups within each day of trial, as assessed by Tukey HSD test ($p < 0.05$); in (E) statistically significant differences were evaluated considering the priming treatment (T) \times inoculation (I) \times stress condition (S) interaction, and lower case letters are indicated only when the T \times I \times S interaction is significant, as assessed by Tukey HSD test ($p < 0.05$). ns, *, **, and ***: not significant or significant at $p \leq 0.05$, $p \leq 0.01$, and $p \leq 0.001$, respectively. CTRL: untreated, SA: treated with salicylic acid, CHI+SA: treated with chitosan + salicylic acid, NMYC: not-inoculated, MYC: inoculated, NS: not-stressed

condition, WS: water stress condition. Analysis of variance on the single variables is reported in Supplementary Tables S2-S3.

Fig. 3

Effects of priming treatment and arbuscular mycorrhizal fungal (AMF) inoculation on the physiological performances of tomato (*Solanum lycopersicum* cv Moneymaker) under salt stress at the end of the experiment. (A) stomatal conductance (g_s , mol H₂O m⁻² s⁻¹), (B) leaf transpiration (E , mmol H₂O m⁻² s⁻¹) and (C) assimilation rates (A_N , μmol CO₂ m⁻² s⁻¹). All results are reported as mean ± SE. Lower case letters above bars denote significant differences as assessed by Tukey HSD test ($p < 0.05$) and are reported only when the priming treatment (T) × inoculation (I) × stress condition (S) interaction is significant. ns, *, **, and ***: not significant or significant at $p \leq 0.05$, $p \leq 0.01$, and $p \leq 0.001$, respectively. CTRL: untreated, SA: treated with salicylic acid, CHI+SA: treated with chitosan + salicylic acid, NMYC: not-inoculated, MYC: inoculated, NS: not-stressed condition, SS: salt stress condition. Analysis of variance on the single variables is reported in Supplementary Table S4.

Fig. 4

Content of phytohormones and proline in leaves of tomato (*Solanum lycopersicum* cv Moneymaker) potted plants. Concentrations of (A) salicylic acid - SA, (B) jasmonic acid - JA, (C) abscisic acid - ABA, (D) indole-3-acetic acid - IAA, and (E) proline. For phytohormones, data are expressed as ng g⁻¹ of dry weight (DW), for proline, data are expressed as μmol g⁻¹ DW. All results are the mean ± SE. Lower case letters above bars denote significant differences as assessed by Tukey HSD test ($p < 0.05$) and are reported only when the priming treatment (T) × inoculation (I) × stress condition (S) interaction is significant. ns, *, **, and ***: not significant or significant at $p \leq 0.05$, $p \leq 0.01$, and $p \leq 0.001$, respectively. CTRL: untreated, SA: treated with salicylic acid, CHI+SA: treated with chitosan + salicylic acid, NMYC: not-inoculated, MYC: inoculated, NS: not-stressed condition, WS: water stress condition, SS: salt stress condition. Analysis of variance on the single variables is reported in Supplementary Table S6.

Fig. 5

Overview of Differentially Expressed Genes (DEGs) compared to control (CTRL_NMYC_NS) between the seventeen different conditions. (A) Total number of up- and down-regulated DEGs generated using the *ggplot2* R package (R version 4.1.1). (B) Heatmap and clustering (Pearson

distance) of the normalized expression levels of the 14,316 DEGs among the different conditions, generated using the *pheatmap* R package (R version 4.1.1). Columns indicate different conditions, whereas rows indicate genes and their expression profiles. Color intensity indicates different levels of expression (\log_2 Fold Change).

Fig. 6

UpSet plot to summarize up (A) and down (B) Differentially Expressed Genes (DEGs) overlapping in each condition. Black dots in each panel matrix represent unique and overlapping DEGs. Connected circles indicate an intersection of DEGs among conditions. Bottom left horizontal bars show the number of total detected DEGs per condition, while the top bar graph in each panel summarizes the number of DEGs for each unique or overlapping combination generated using the *upsetplot* R package (R version 4.1.1).

Fig. 7

Overview of metabolomic changes in tomato (*Solanum lycopersicum* cv Moneymaker) leaves under the different tested conditions: inoculated (MYC), not-inoculated (NMYC), seed-treated with chemical priming (SA, CHI+SA), untreated (CTRL), under not-stressed (NS), water stress (WS) or salt stress (SS). (A) Two-dimensional Partial Least Squares Discriminant Analysis (2D-PLSDAs) of the 372 most significant chromatographic masses in NS and stressed (WS, SS) conditions, regardless of priming treatment and inoculation. (B) 2D-PLSDAs of the 372 most significant chromatographic masses in NMYC and MYC plants, regardless of priming treatment and stress condition. 2D-PLSDAs plots were generated with the MetaboAnalyst software, considering the normalised dataset as \log_{10} . (C) Heatmap and clustering (Euclidean distance) of 64 differentially accumulated metabolites selected among those having the highest variability in accumulation intensity ($\log_2FC > 1$ compared to CTRL_NMYC_NS) in at least one condition. The heatmap was generated using the *pheatmap* R package (R version 4.1.1). Data in cells are \log_2 -transformed mean values of three biological replicates.

Fig. 8

Arrow plot from multiblock sPLS-DA performed on the metabolomic and RNA-seq data of all tested conditions. Start of the arrow indicates the centroid between all data sets for each condition, while tips of the arrows indicate their location in each block.

Fig. 9

Overview of long-lasting physiological, metabolic and molecular responses occurring in AMF-inoculated (MYC) tomato plants (*Solanum lycopersicum* cv Moneymaker) either subjected to chemical seed priming (*i.e.* SA_MYC and CHI+SA_MYC) or not (CTRL_MYC). Green and yellow arrows respectively indicate decrease and increase in the measured ecophysiological parameters/gene expression/metabolite content, while the = symbol refers to absence of changes. ABA: abscisic acid, JA: jasmonic acid, CTRL: untreated, SA: treated with salicylic acid, CHI+SA: treated with chitosan + salicylic acid, MYC: inoculated, WS: water stress, SS: salt stress.

Accepted Manuscript




Fig. 1 Arbuscular mycorrhizal fungal (AMF) colonization assessment in tomato potted plants (*Solanum lycopersicum* cv MoneyMaker) inoculated with a commercial inoculum (MycAgro, *Funneliformis mosseae*). Top panels show representative images of AMF-inoculated roots. (A) F%, Frequency of mycorrhization in root system; bar (AI) = 120 μ m; (B) a%, arbuscule abundance in the colonized root system; bar (BI) = 60 μ m, bar (BII) = 20 μ m. All results are reported as mean \pm SE. Lower case letters above bars denote significant differences as assessed by Tukey HSD test ($p < 0.05$) and are reported only when the priming treatment (T) \times stress condition (S) interaction is significant. ns, *: not significant or significant at $p \leq 0.05$, respectively. CTRL: untreated, SA: treated with salicylic acid, CHI+SA: treated with chitosan + salicylic acid, NMYC: not-inoculated, MYC: inoculated, NS: not-stressed condition, WS: water stress condition, SS: salt stress condition. Analysis of variance on the single variables is reported in Supplementary Table S1.

Accepted Manuscript




Fig. 2 Effects of priming treatment and arbuscular mycorrhizal fungal (AMF) inoculation on the physiological performances of tomato (*Solanum lycopersicum* cv MoneyMaker) potted plants during a water stress time-course. (A) Changes in soil relative water content (RWC_{soil} %) monitored over the whole duration of the water stress time-course. The inset beside panel A) magnifies differences in RWC_{soil} values among the diverse groups of plants during the first four days after water withdrawal (T1-T5); (B-D) Dynamic changes in the rates of (B) stomatal conductance (g_s, mol H₂O m⁻² s⁻¹), (C) leaf vapour pressure deficit (VPD_{leaf}, kPa) and (D) leaf transpiration (E, mmol H₂O m⁻² s⁻¹) monitored over the whole duration of the water stress time-course; (E) Rates of g_s measured at the end (T_f) of the experiment. All results are reported as mean ± SE. In panel A), asterisks denote significant differences determined by one-way ANOVA on RWC_{soil} data (*, **, and *** significant at p ≤ 0.05, p ≤ 0.01, and p ≤ 0.001, respectively, with corresponding values reported in the panel inset), while different lower case letters in the panel A) inset highlight significant differences among plant groups within each day of trial, as assessed by Tukey HSD test (p < 0.05); in (E) statistically significant differences were evaluated considering the priming treatment (T) × inoculation (I) × stress condition (S) interaction, and lower case letters are indicated only when the T × I × S interaction is significant, as assessed by Tukey HSD test (p < 0.05). ns, *, **, and ***: not significant or significant at p ≤ 0.05, p ≤ 0.01, and p ≤ 0.001, respectively. CTRL: untreated, SA: treated with salicylic acid, CHI+SA: treated with chitosan + salicylic acid, NMYC: not-inoculated, MYC: inoculated, NS: not-stressed condition, WS: water stress condition. Analysis of variance on the single variables is reported in Supplementary Tables S2-S3.




Fig. 3 Effects of priming treatment and arbuscular mycorrhizal fungal (AMF) inoculation on the physiological performances of tomato (*Solanum lycopersicum* cv MoneyMaker) under salt stress at the end of the experiment. (A) stomatal conductance (g_s , mol H₂O m⁻² s⁻¹), (B) leaf transpiration (E , mmol H₂O m⁻² s⁻¹) and (C) assimilation rates (A_N , μmol CO₂ m⁻² s⁻¹). All results are reported as mean ± SE. Lower case letters above bars denote significant differences as assessed by Tukey HSD test ($p < 0.05$) and are reported only when the priming treatment (T) × inoculation (I) × stress condition (S) interaction is significant. ns, *, ***, not significant or significant at $p \leq 0.05$, $p \leq 0.001$, respectively. CTRL: untreated, SA: treated with salicylic acid, CHI+SA: treated with chitosan + salicylic acid, NMYC: not-inoculated, MYC: inoculated, NS: not-stressed condition, SS: salt stress condition. Analysis of variance on the single variables is reported in Supplementary Table S4.





Fig. 4. Content of phytohormones and proline in leaves of tomato (*Solanum lycopersicum* cv Moneymaker) potted plants. Concentrations of (A) salicylic acid - SA, (B) jasmonic acid - JA, (C) abscisic acid - ABA, (D) indole-3-acetic acid - IAA, and (E) proline. For phytohormones, data are expressed as ng g⁻¹ of dry weight (DW), for proline, data are expressed as μmol g⁻¹ DW. All results are the mean ± SE. Lower case letters above bars denote significant differences as assessed by Tukey HSD test ($p < 0.05$) and are reported only when the priming treatment (T) × inoculation (I) × stress condition (S) interaction is significant. ns, *, **, and ***: not significant or significant at $p \leq 0.05$, $p \leq 0.01$, and $p \leq 0.001$, respectively. CTRL: untreated, SA: treated with salicylic acid, CHI+SA: treated with chitosan + salicylic acid, NMYC: not-inoculated, MYC: inoculated, NS: not-stressed condition, WS: water stress condition, SS: salt stress condition. Analysis of variance on the single variables is reported in Supplementary Table S6.

A



B




Fig. 5. Overview of Differentially Expressed Genes (DEGs) compared to control (CTRL_NMYC_NS) between the seventeen different conditions. (A) Total number of up- and down-regulated DEGs generated using the *ggplot2* R package (R version 4.1.1). (B) Heatmap and clustering (Pearson distance) of the normalized expression levels of the 14,316 DEGs among the different conditions, generated using the *heatmap* R package (R version 4.1.1). Columns indicate different conditions, whereas rows indicate genes and their expression profiles. Color intensity indicates different levels of expression (Log_2 Fold Change).




Fig. 6. UpSet plot to summarize up (A) and down (B) Differentially Expressed Genes (DEGs) overlapping in each condition. Black dots in each panel matrix represent unique and overlapping DEGs. Connected circles indicate an intersection of DEGs among conditions. Bottom left horizontal bars show the number of total detected DEGs per condition, while the top bar graph in each panel summarizes the number of DEGs for each unique or overlapping combination generated using the *upsetplot* R package (R version 4.1.1).




Fig. 7. Overview of metabolomic changes in tomato (*Solanum lycopersicum* cv Moneymaker) leaves under the different tested conditions: inoculated (MYC), not-inoculated (NMYC), treated with chemical priming (SA, CHI+SA), untreated (CTRL), under not-stressed (NS), water stress (WS) and salt stress (SS). (A) Two-dimensional Partial Least Squares Discriminant Analysis (2D-PLSDAs) of the 372 most significant chromatographic masses in NS and stressed (WS, SS) conditions, regardless of priming treatment and inoculation. (B) 2D-PLSDAs of the 372 most significant chromatographic masses in NMYC and MYC plants, regardless of priming treatment and stress condition. 2D-PLSDAs plots were generated with the MetaboAnalyst software, considering the normalised dataset as Log_{10} . (C) Heatmap and clustering (Euclidean distance) of 64 differentially accumulated metabolites selected among those having the highest variability in accumulation intensity ($\text{Log}_2\text{FC} > 1$ compared to CTRL_NMYC_NS) in at least one condition. The heatmap was generated using the *heatmap* R package (R version 4.1.1). Data in cells are Log_2 -transformed mean values of three biological replicates.




Fig. 8. Arrow plot from multiblock sPLS-DA performed on the metabolomic and RNA-seq data of all tested conditions. Start of the arrow indicates the centroid between all data sets for each condition, while tips of the arrows indicate their location in each block.

Accepte




Fig. 9 Overview of long-lasting physiological, metabolic and molecular responses occurring in AMF-inoculated (MYC) tomato plants (*Solanum lycopersicum* cv Moneymaker) either subjected to chemical seed priming (i.e. SA_MYC and CHI+SA_MYC) or not (CTRL_MYC). Green and yellow arrows respectively indicate decrease and increase in the measured ecophysiological parameters/gene expression/metabolite content, while the = symbol refers to absence of changes. ABA: abscisic acid, JA: jasmonic acid, CTRL: untreated, SA: treated with salicylic acid, CHI+SA: treated with chitosan + salicylic acid, MYC: inoculated, WS: water stress, SS: salt stress.

Accepted