



Legionella and legionellosis in touristic-recreational facilities: Influence of climate factors and geostatistical analysis in Southern Italy (2001–2017)



Osvalda De Giglio^a, Fabrizio Fasano^a, Giusy Diella^a, Marco Lopuzzo^a, Christian Napoli^b, Francesca Apollonio^a, Silvia Brigida^c, Carla Calia^a, Carmen Campanale^a, Angelo Marzella^a, Chrysovalentinos Pousis^a, Serafina Rutigliano^a, Francesco Triggiano^a, Giuseppina Caggiano^a, Maria Teresa Montagna^{a,*}

^a Department of Biomedical Science and Human Oncology, University of Bari "Aldo Moro", Piazza G. Cesare 11, 70124, Bari, Italy

^b Department of Medical Surgical Sciences and Translational Medicine, "Sapienza" University of Rome, Via di Grottarossa 1035/1039, 00189, Rome, Italy

^c Water Research Institute-Italian National Research Council, Bari, Italy

ARTICLE INFO

Keywords:

Legionella

Geostatistical analysis

Climate

Touristic facilities

ABSTRACT

Legionella is the causative agent of Legionnaires' disease, a flu-like illness normally acquired following inhalation or aspiration of contaminated water aerosols. Our recent studies revealed that climatic parameters can increase the number of reported cases of community-acquired Legionnaires' disease. Here, we evaluated the presence of *Legionella* in water networks and the distribution of Legionnaires' disease cases associated with touristic-recreational facilities in the Apulia region (southern Italy) during the period 2001–2017 using geostatistical and climatic analyses. Geostatistical analysis data revealed that the area with the highest concentration of *Legionella* in water systems also had the greatest number of cases of Legionnaires' disease associated with touristic-recreational facilities. Climatic analysis showed that higher daily temperature excursion (difference between maximum and minimum temperature) on the day of sampling was more often associated with *Legionella*-positive samples than *Legionella*-negative samples. In addition, our data highlighted an increased risk of Legionnaires' disease with increases in precipitation and average temperature and with decreases in daily temperature excursion (difference between maximum and minimum temperature over the course of 24 h in the days of incubation period of disease) and minimum temperature.

Healthcare professionals should be aware of this phenomenon and be particularly vigilant for cases of community-acquired pneumonia during such climatic conditions and among the tourist population. The innovative geo-statistical approach used in this study could be applied in other contexts when evaluating the effects of climatic conditions on the incidence of *Legionella* infections.

1. Introduction

Legionella is a facultative intracellular Gram-negative microorganism responsible of infectious disease commonly called legionellosis. The different clinical forms of legionellosis range from severe pneumonia known as Legionnaires' disease (LD), to a flu-like illness, Pontiac fever. Legionellosis is usually acquired by inhalation or aspiration of aerosols originating from water sources containing

Legionella. This microorganism is ubiquitous in natural and artificial water systems (Costa et al., 2010; Graells et al., 2018; Haupt et al., 2012; Montagna et al., 2012, 2016; 2017a, 2017b; 2018; De Giglio et al., 2019) sediments and soils as individual cells, but is most commonly found colonising amoeba or within biofilms (Fields, 1996). *Legionella* grows at temperatures between 25 °C and 45 °C (Garrity et al., 2005), but can survive at temperatures ranging from 5.7 °C to 63 °C, especially if the water is stagnant (Borella et al., 2005; Napoli et al.,

Abbreviations: (LD), Legionnaire's disease; (sg), serogroup; (IQR), interquartile range

* Corresponding author.

E-mail addresses: osvalda.degiglio@uniba.it (O. De Giglio), fabrizio.fasano1979@libero.it (F. Fasano), giusy.diella@uniba.it (G. Diella), marcolopuzzo@gmail.com (M. Lopuzzo), christian.napoli@uniroma1.it (C. Napoli), francesca.apo@libero.it (F. Apollonio), silvia.brigida@ba.irs.cnr.it (S. Brigida), carla.calia@uniba.it (C. Calia), campanale.carmen@libero.it (C. Campanale), marzella.angelo@libero.it (A. Marzella), vpousis@gmail.com (C. Pousis), rutigliano.s@gmail.com (S. Rutigliano), francescotrigg@hotmail.it (F. Triggiano), giuseppina.caggiano@uniba.it (G. Caggiano), mariateresa.montagna@uniba.it (M.T. Montagna).

<https://doi.org/10.1016/j.envres.2019.108721>

Received 21 May 2019; Received in revised form 2 September 2019; Accepted 3 September 2019

Available online 06 September 2019

0013-9351/ © 2019 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license

(<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

2010). Several studies on the ecology of the *Legionella* genus have highlighted the limitations of detection system reliant on culture-based methods and the importance of taking the life cycle of the pathogen into account. For example, significant numbers of *Legionella* isolates have been recovered from cold waters sources (Wullings and van der Kooij, 2006; Wadowsky et al., 1985), including Antarctic lakes ($\sim 0^{\circ}\text{C}$) (Carvalho et al., 2008) using cultivation-independent techniques. For this reason, it is important to investigate all environmental parameters influencing the presence of *Legionella* (Graells et al., 2018).

The susceptibility of different individuals to the same source of infection depends on individual host factors and on concomitant diseases (Montagna et al., 2006; Napoli et al., 2010). To date, 61 species and more than 70 serogroups (sg) of *Legionella* have been reported (Euzéby, 2019, <http://www.bacterio.net/legionella.html>). In particular, the species *Legionella pneumophila*, which is most frequently associated with human disease, includes 16 serogroups. However, the number of *Legionella* species is constantly changing due to recent isolated species (Palmer et al., 2016; Relich et al., 2018). Although previous studies have shown that *Legionella pneumophila* sg 1 strains are the most cause of disease in humans, an increasing number of cases are being attributed to other species and serogroups (ECDC, 2018; Napoli et al., 2010; Palmer et al., 2016; Relich et al., 2018).

The surveillance of legionellosis in Europe is coordinated by the European Centre for Disease Prevention and Control (ECDC), and in Italy by the Italian National Institute of Health. Since 2010, the ECDC has coordinated European Legionnaires' Disease Surveillance Network (ELDSNet) an international surveillance of travel-associated LD. According to national data (ISS, 2018), in 2017, a total of 2014 legionellosis cases were reported in Italy (33.2 cases per 1 million inhabitants), corresponding to a slight increase with the previous year (28.2/1,000,000). After community-associated cases ($n = 1,580$, 78.5%), the largest number of cases was associated with travel ($n = 239$, 11.9%; Italian citizen diagnosed in Italy), followed by healthcare facility-associated cases ($n = 124$, 6.2%) and other cases ($n = 71$, 3.5%). If foreign tourists who likely acquired the infection in Italy but were diagnosed elsewhere were also considered, the number of travels-associated legionellosis cases in 2017 increased to 436 (ISS, 2018; European Centre for Disease Prevention and Control (ECDC), 2018). Overall, the case fatality rate in Italy ranges from 10.1% for community-associated cases to 51.1% for nosocomial cases (ISS, 2018).

In 2015, all of the national recommendations for the prevention and control of legionellosis, including those for touristic-recreational facilities, were incorporated into a single updated document (Linee, 2015; Regione 2015). These instructions list risk assessment evaluation as one of the most effective prevention measures for managing *Legionella* contamination of water systems. This is particularly relevant in touristic-recreational facilities because they often have complex water systems with a large number of outlets (e.g., showers) (Ricketts et al., 2010). In the absence of regular flushing, such as in unoccupied rooms or during annual closure, water may stagnate in pipes, favouring the growth of *Legionella*. In addition, it can be difficult to maintain adequate temperature control of hot and cold water within long pipe systems (Phin et al., 2014).

In 1998, the government of Apulia region (southern Italy) established a regional epidemiological observatory (OER-Apulia) at the Department of Biomedical Science and Human Oncology, University of Bari Aldo Moro. The observatory is tasked with monitoring and performing surveillance of sanitary events, including all clinical forms of legionellosis. Since 2000, ongoing environmental and clinical monitoring of *Legionella* has been conducted throughout all of the regional territories (Montagna et al., 2006, 2012; Napoli et al., 2007, 2009a, 2009b), with all environmental and clinical isolates being stored in a stock archive for further studies (Iatta et al., 2013a, 2013b; Montagna et al., 2016). Data are regularly entered into an informatic and integrated platform that allows for the delivery of real-time information concerning notified cases and the distribution and degree of

contamination in public and private water systems.

One of the main activities of the Apulia region is tourism thanks to its geographical and climatic characteristics, well as its popularity as a pilgrimage destination (Castellani Pastoris et al., 2002). The official tourism observatory of the Apulia region, Puglia Promozione, publishes data regarding tourism in Apulia compared with other regions in Italy in collaboration with the Italian National Institute of Statistics. Apulia is among the top 10 Italian tourism regions in terms of the number of overnight stays (Portale dell'Agenzia Regionale del Turismo PUGLIA-PROMOZIONE, Report data 2017, <https://www.agenziapugliapromozione.it/portal/guest/osservatorio-del-turismo>). Over the last 17 years, the Apulia region recorded 519 cases of LD, amongst which 230 cases (44.3%) were contracted from touristic-recreational facilities (Portale dell'Osservatorio Epidemiologico regionale, <https://www.sanita.puglia.it/osservatorio-epidemiologico-regionale>). Currently, Italian and regional guidelines (Linee, 2015; Regione Puglia 2015) on the control and prevention of legionellosis at touristic-recreational facilities are considered reference documents only and are not enforced. As such, very little is known about the implementation rates of these guidelines (Montagna et al., 2018).

Several studies have reported that climate change leading to increased water temperature, drought, higher rainfall, rising sea levels and flooding could play an important role in the transmission of waterborne diseases, including legionellosis (Beauté et al., 2016; Brandsema et al., 2014; Conza et al., 2014; Garcia-Vidal et al., 2013; Karagiannis et al., 2009; Walker, 2018). Therefore, combining classical statistical analysis methods with georeferencing analyses may be of particular use in understanding the sources of contamination (Laganà et al., 2015).

The aim of the current study was to evaluate the presence of *Legionella* in water networks and the distribution of legionellosis cases associated with touristic-recreational facilities in the Apulia region of southern Italy during the period 2001–2017, using geostatistical and climatic analyses.

2. Materials and methods

2.1. Geographical area of study

Apulia, in south-eastern Italy, is a region between the Adriatic and Ionian seas. With a land surface area of 19,347 km², the region has about 800 km of coastline and 4 million inhabitants. It is characterised by a typical Mediterranean climate, with mild, dry winters and hot summers. Rainfall is the largest source of groundwater recharge, but is highly irregular throughout the year. Therefore, water is supplied by the Apulian Aqueduct, which draws from a large network ($\sim 20,000\text{ km}^2$) of interconnected natural and artificial water reservoirs (Sinni-Pertusillo, Sele, Calore, Locone, Ofanto, Fortore) (Fig. 1) and has a complex system of drainage wells, with different purposes (drinking, irrigation and industrial) (De Giglio et al., 2016). Based on data from the Italian National Agency for the Environmental Protection, Apulia has some of the highest average temperatures across all regions of Italy; moreover, although it is not among the regions with the highest rainfall (ISPRA 2014, http://www.isprambiente.gov.it/files/pubblicazioni/statoambiente/copy_of_SA_55_14_Valori_climatici_normali.pdf), it is characterised by areas (Gargano), where rainfall is consistently higher than in the rest of the region (Istituto Superiore per la Protezione e la Ricerca Ambientale - ISPRA, Regione Puglia 2014). Based on homogeneous climatic data, it is possible to distinguish five distinct areas of the region (Gargano, Tavoliere, Murgia, Salento, Arco Jonico) (ISPRA 2014, http://www.isprambiente.gov.it/files/pubblicazioni/statoambiente/copy_of_SA_55_14_Valori_climatici_normali.pdf; De Giglio et al., 2016).



Fig. 1. Integrated water supply and transport system for civil use in the Apulia region [Source: Apulian aqueduct S.p.A.].

2.2. Touristic-recreational facilities

After considering a proportionate and representative territorial distribution of all Apulian tourist-recreational facilities ($n = 6332$), 495 facilities (7.8% of all Apulian tourist-recreational facilities) were enrolled in our study, with 6638 water samples analysed in total. The characteristics of each facility, including type (e.g., hotel, B&B), classification (number of hotel stars: 1–5), period of activity (i.e., annual or seasonal) and tourist arrivals (grouped by municipality, month and year), were acquired from two regional official websites ([Portale ufficiale del turismo della Regione Puglia](https://www.viaggiareinpuglia.it/hp/it), <https://www.viaggiareinpuglia.it/hp/it>; [Portale dell'Open Data Puglia](http://www.dataset.puglia.it/), <http://www.dataset.puglia.it/>).

2.3. Microbiological investigation

Sampling and processing procedures were performed according to the Italian guidelines (Linee, 2000, 2015). Water samples (1 L) were collected from showers (42.4%), taps (36.1%), boilers (26.6%) and other sampling points (16.6%) in sterile containers containing sodium thiosulphate pentahydrate (0.01%, w/v) to neutralise chloride present in the water, and were transported at environmental temperature in the dark. Samples (1 L) were filtered through 0.2- μm isopore polycarbonate membranes (Millipore Corporation, Bedford, MA, USA), with the membrane then resuspended in 10 ml of the same water sample and vortexed. A 5 ml aliquot of the suspension was then incubated at 50 °C for 30 min. Aliquots (100 μl) of both the heat-treated and untreated samples were seeded onto GVPc (glycine vancomycin polymyxin cycloheximide) agar plates (BioMérieux, Marcy l'Etoile, France) and incubated at 36 ± 1 °C for 10 days in a modified atmosphere (2.5% CO_2). It was not considered necessary to carry out acid treatment because the analysed samples were not abundant in contaminating flora. Putative *Legionella* colonies were subcultured on buffered charcoal yeast extract (BCYE) agar (BioMérieux, Marcy l'Etoile, France) and BCYE agar without cysteine (BioMérieux, Marcy l'Etoile, France). Colonies that grew only in the presence of cysteine were identified as *Legionella* and

confirmed using a latex agglutination test with polyvalent (Oxoid Spa, Milan, Italy) and monovalent antisera (Biogenetics Srl, Tokyo, Japan) (ISO 11731, 1998) as per the manufacturer's instruction. As recommended by the Italian guidelines (Linee, 2015), water samples containing < 100 colony-forming units per litre (cfu/L) were considered negative for *Legionella*. Facilities with at least one positive sample were considered positive for *Legionella*.

2.4. Data collection

2.4.1. Cases of travel-associated legionellosis

All legionellosis cases notified to OER-Apulia in the years 2001–2017 were included in this study. A case of legionellosis was considered travel-associated if the patient had slept away from home for one or more nights in the 10 days prior to the onset of symptoms (Linee, 2015; Regione Puglia, 2015; 2018). Legionellosis cases notified to OER-Apulia also included foreign tourists (coming from another location or country) who had stayed in the Apulia region (ELDSNET surveillance system).

2.4.2. Climatic characteristics

Climatic data were obtained from meteorological stations present in the region ([Portale della Protezione Civile Puglia](http://www.protezionecivile.puglia.it/), <http://www.protezionecivile.puglia.it/>).

For environmental surveillance, we considered the air temperature (maximum, minimum, average) and thermal excursion (difference between maximum and minimum daily temperature) on the day of sampling. The association between *Legionella* load in water samples and air temperature was studied using only the water samples that were positive for *Legionella*. In addition, a case-control study was carried out to verify any statistically significant differences between *Legionella*-positive and -negative water samples.

In relation to clinical surveillance, to evaluate the influence of climatic parameters on LD cases spread, maximum, minimum and average temperatures, thermal excursion (difference between maximum and

minimum temperature), daily precipitation and relative humidity (RH) were investigated over the days corresponding to the incubation period of LD (2–10 days prior to the onset of symptom) and within the month that symptoms began (e.g., if a case occurred in April, we considered the climatic parameters from April 1st to April 30th).

All climatic data were recovered from meteorological stations within a maximum distance of 30 km from the sampling point (for environmental analysis) or from the facility designated as the probable source of infection (for clinical surveillance).

2.5. Geo-statistical analysis

All of the water sampling locations and touristic-recreational facilities were georeferenced using geographic coordinates provided by the regional authorities for “hot spot” analysis. Generally, this analysis is used to pinpoint locations of statistically significant high- and low-value clusters of a phenomenon, evaluating each feature (e.g., *Legionella* load in water samples) in relation to the features of its neighbour and against all features in the dataset (Getis and Ord, 1992). A feature with a high value may be a statistically significant hot spot if it is also surrounded by other features with high values, as opposed to simply being a data outlier. When the observed local mean differed from the global mean and this difference was too large to be the result of random chance, a statistically significant z-score is generated and a hot spot cluster is detected.

The Getis-Ord algorithm (Getis and Ord, 1992, 1996; Ord and Getis, 1995) was used to identify a hot spot. This tool, also called the G_i^* statistic, calculates spatial dependencies between an observed datum and its neighbours within a threshold distance that can be pre-defined or set by the user. The G_i^* statistic indicates whether spatial entities with high (or very low) presence values tend to cluster within a certain distance in certain areas.

To determine spatial relationships in the hot spot analyses for punctual geometry, the most commonly method used in literature (Sánchez-Martín et al., 2019) is Fixed distance band where each feature is analysed within the context of neighbouring features. Neighbouring features within the specified critical distance (Distance Band or Threshold Distance) receive a weight of 1 and exert influence on computations for the target feature. Neighbouring features outside the critical distance receive a weight of zero and have no influence on a target feature's computations.

To calculate distances between neighbours, the Euclidean distance (which uses the straight-line distance between points A and B, i.e., the shortest possible distance) is the most commonly method used in literature (Sánchez-Martín et al., 2019). Because there was no precedent for our analyses of *Legionella* and legionellosis, we applied Fixed distance band with Euclidean distance, ensuring that each feature had at least one neighbour.

Maps of *Legionella* distribution were generated using an interpolation technique, known as inverse distance weighting (IDW), based on G_i^* statistic scores from the hot spot analysis. The principle of the IDW method is to consign more weight to nearby points compared with distant points. IDW weights the contribution of each input point by a normalised inverse of the distance from the control point to the interpolated point (Chang, 2010). The IDW interpolation method assumes that each input point has local influence that decreases with distance (Fisher et al., 1987). QGIS version 3.2 (Download QGIS version 3.2, <https://qgis.org/it/site/forusers/download.html>) was used for georeferencing of the points, using the Getis-Ord algorithm and to generate the maps through IDW.

2.6. Statistical analysis

R software version 3.5.1 was used for statistical analysis, with $p < 0.05$ considered statistically significant.

Fisher's exact test or a χ^2 test with Yates correction were used to

investigate the association between air temperature and *Legionella* load in water networks, and between climatic parameters (temperature, humidity, daily precipitation) and legionellosis cases.

The Shapiro-Wilk normality test (SWt), the Kolmogorov-Smirnov test (KSt) and a quantile-quantile (QQ) plot were used to evaluate normal or non-normal distributions of the *Legionella* load and climatic data. When SWt and KSt vales have a p -value $p < 0.05$, the distribution was assumed to be non-normal and non-parametric tests were used.

Kendall's Tau-b test was used to investigate the correlation between *Legionella* load and air temperature on the day of water sampling. In cases where the correlation coefficient was > 0 or < 0 , a positive or negative correlation was assumed, respectively.

A regression study was carried out to explain the relationship between the trend in *Legionella* load (dependent) and air temperature (independent) by estimation of the curve with respect to the main models (Linear, Logarithmic, Inverse, Quadratic, Cubic, Compound, Power, S).

The Wilcoxon rank sum test with continuity correction was used to verify statistically significant differences in the case-control study or to verify statistically significant differences among different areas of the study.

Kruskal-Wallis and Pairwise comparisons using a Wilcoxon rank sum test were used to compare climatic parameters of different areas within the Apulia region.

2.6.1. Multivariate analysis of risk factors for LD

A Poisson regression model was used to perform the multivariate analysis of climatic risk factors associated with LD. Climatic data were obtained from the main weather stations in the five homogeneous climatic areas of Apulia (Gargano, Tavoliere, Murgia, Salento and Arco Jonico) from 2001 to 2017, with all data stratified by month. These data were then correlated with cases (presented as no. of cases/10,000 tourist arrivals). The climatic characteristics for each month in which a clinical LD case was recorded were divided into two parts. In the first part the climatic data for the incubation period of the disease were entered and the number of cases that occurred per 10,000 tourist arrivals was estimated using the following formula:

[(No of arrivals in the month/No of days in the month) x No of days in the incubation period].

In the second part the climatic parameters for the month, excluding the days corresponding to the incubation period, along with the arrivals during the incubation period, were considered. In the case of months in which no cases occurred, the climatic parameters of the same month were considered.

The final model included only those variables with a p -value of < 0.05 in the preliminary model with all the variables. To quantify the effects of meteorological variables, we computed the influences ($e^{\beta} - 1$), which virtually corresponded to the relative risks (Conza et al., 2013). R software version 3.5.1 was used for this analysis.

3. Results

3.1. Hot spot analysis of water samples and climatic parameters

During the period 2001–2017, *Legionella* was detected at 339 (68.5%) of the tested facilities and in 2459 (37%) of the water samples. *Legionella pneumophila* sg 1 was the most frequently isolated species and sg (72%). Hot spot analysis of the *Legionella* load (cfu/L) is shown in Fig. 2. Hot spots with a 99% confidence interval (red point) were considered the most relevant. The area in red contained the greatest number of relevant hot spots based on *Legionella* load.

Among the *Legionella*-positive samples, the mean bacterial load was 10.8×10^4 cfu/L (SD, 37.6×10^4 cfu/L) with a median of 3.6×10^3 (IQR, 10.1×10^4 cfu/L). Fig. 3 shows the QQ plot of the *Legionella* load (cfu/L).

SWt and KSt analyses produced a W statistic = 0.20, $p < 0.0001$




Fig. 2. Hot spot analysis of the *Legionella* load in water samples from touristic-recreational facilities in the Apulia region (2001–2017).




Fig. 3. QQ plot of the *Legionella* load (cfu/L) from 2001 to 2017.




Fig. 4. QQ plot of the air temperature (°C) versus with all water sampling points.

and $D = 0.39$, $p < 0.0001$, respectively, indicating that the distribution was not normal.

Kendall's Tau-b (T_b) test showed that the correlation coefficients were very low, meaning that it was not possible to confirm a correlation between the *Legionella* load (cfu/L) and the following parameters:

- maximum temperature (°C) (T_b ; $p = 0.018$, correlation coefficient = 0.035);
- minimum temperature (°C) (T_b ; $p = 0.008$, correlation coefficient = 0.039);

- average temperature (°C) (T_b ; $p = 0.012$, correlation coefficient = 0.037);
- daily thermal excursion (°C) (T_b ; $p = 0.045$, correlation coefficient = 0.030).

Table 1
Case control study between the temperature parameters and *Legionella*-positive and -negative water samples.

Air temperature (°C)	POSITIVE water samples (N = 2459) (median ± IQR)	NEGATIVE water samples (N = 4179) (median ± IQR)	Wilcoxon rank sum test with continuity correction (W)	p-value
Maximum	23.0 ± 9.7	22.1 ± 10.8	3,674,900	0.2533
Minimum	15.7 ± 9.3	16.2 ± 10.2	3,529,100	0.1662
Average	19.3 ± 9.9	19.1 ± 9.9	3,613,200	0.9423
Daily thermal excursion	6.8 ± 3.2	6.6 ± 3.4	3,881,600	0.0023

IQR = interquartile range.

Legend

Hot spot analysis

Gi_Bin

- Cold Spot - 99% Confidence
- Cold Spot - 95% Confidence
- Cold Spot - 90% Confidence
- Not Significant
- Hot Spot - 90% Confidence
- Hot Spot - 95% Confidence
- Hot Spot - 99% Confidence

IDW on Hot spot analysis

GZ Score - Hot spot Analysis

- -2.27 - -1.78
- -1.79 - -1.18
- -1.19 - -0.70
- -0.69 - -0.38
- -0.37 - 0.14
- 0.15 - 0.74
- 0.75 - 1.32
- 1.33 - 1.84
- 1.85 - 2.21
- 2.22 - 2.85




Fig. 5. Hot spot analyses of LD cases in tourist recreational facilities in the Apulia region (2001–2017).

Table 2
Cases of LD linked to tourist recreational facilities and tourist turnout in the Apulia region by month (for years 2001–2017).

Month	Cases of LD/tourist arrivals	%
Jan	1/1,330,533	0.00008
Feb	3/1,537,958	0.00020
Mar	5/1,978,748	0.00025
Apr	19/3,377,922	0.00056
May	30/4,004,798	0.00075
Jun	43/6,054,586	0.00071
Jul	19/8,047,139	0.00024
Aug	25/10,075,702	0.00025
Sep	44/5,438,084	0.00081
Oct	30/3,145,781	0.00095
Nov	10/1,800,913	0.00056
Dec	1/1,788,470	0.00006
Total	230/48,580,636	0.00047

Table 3
Cases of LD linked to tourist recreational facilities and tourist turnout in the Apulia region by season (for years 2001–2017).

Season	Cases of LD/tourist arrivals	%
Autumn	84/10,384,778	0.00081
Winter	5/4,656,962	0.00011
Spring	54/9,361,468	0.00058
Summer	87/24,177,427	0.00036
Total	230/48,580,636	0.00047

SWt and KSt analyses produced statistically significant (W statistic = 0.98251, $p < 0.0001$ and $D = 0.39$, $p < 0.0001$, respectively) results for the association between air temperature data and all water sampling points, indicating that the distribution was not normal. The QQ plot for the temperature data is presented in Fig. 4.




Fig. 6. QQ plot of the (A) temperature (°C), (B) daily precipitation (mm) and (C) daily relative humidity (%) versus LD cases.

Table 1 shows the data from the case-control study between *Legionella*-positive and negative water samples versus air temperature.

In the case-control study, only the parameter “daily thermal excursion” showed a statistically significant difference ($W = 3,881,600$, $p = 0.002$) between the samples, with the daily thermal excursion shown to be higher in *Legionella*-positive water samples compared with *Legionella*-negative samples.

Analysis of daily temperature excursion in both the hot spot area (n. observations: 830 for positive samples and 715 for negative samples; median value = 7.3 °C vs. 6.6 °C; $W = 156,840$, $p < 0.0001$) and the other areas (n. observations: 1629 for positive samples and 3464 for negative samples; median value = 6.7 °C vs. 6.6 °C; $W = 2,193,500$, $p = 0.03142$) revealed that the daily air temperature excursion on the days on which positive samples were collected was higher than that for the days on which negative samples were collected in the two area group. Moreover, a significant statistical difference was detected for the positive water samples in the two area groups ($W = 463,240$; $p = 0.0005$).

3.2. Hot spot analysis of legionellosis cases and climatic parameters

During the period 2001–2017, 230 cases of LD were associated with

tourist recreational facilities in the Apulia region. Other clinical forms of legionellosis were not notified. Among these 230 cases (87.9% caused by *Legionella pneumophila* sg 1), the mean age was 64.5 years \pm 11.1, with a higher percentage of males (67%) than females. Fig. 5 shows the hot spots for LD cases. Hot spots with a 99% confidence interval (red points) were considered the most relevant. The area shown in red indicates the area in which the greatest number of relevant hot spots of LD cases were located.

Comparison of the data presented in Fig. 2 and Fig. 5 revealed that the areas with the highest concentration of *Legionella* in the water samples also had the highest number of cases of LD associated with staying in a tourist facility.

Analysis of the arrivals registered at the tourist recreational facilities versus the number of LD cases that occurred between 2001 and 2017 revealed an incidence of one LD case per 211,220 tourist arrivals.

Moreover, it was possible to associate the number of travel-associated LD cases with the number of tourist arrivals per month and season (Tables 2 and 3).

The number of LD cases/tourist arrivals was greater in autumn (1 case/123,628), than in summer ($X^2 = 28.705$, $df = 1$, $p < 0.0001$), or winter (Fisher's exact test, $p < 0.0001$). September and October were identified as having the greatest number of LD cases/tourist arrivals;

Table 4

Poisson regression model of meteorological factors associated with risk of LD (by month from 2001 to 2017 in the five principal climatologic areas of Apulia region).

	β	$(e\beta - 1) = \text{RR} (\%)$	p-value
Preliminary model			
Intercept	1,797,044		< 0.0001***
Tavoliere Area	-0.9692	-62.10	< 0.0001***
Gargano Area (Hot spot Area)	0.6309	87.90	< 0.0001***
Murgia Area	-0.8347	-56.60	< 0.0001***
Salento Area	0.1297	13.85	< 0.0001***
Arco Jonico Area	-32.098	-95.50	< 0.0001***
Year	-0.0918	-8.80	< 0.0001***
Month	0.3133	36.80	< 0.0001***
Maximum temperature (°C)	0.3042	35.60	0.80
Minimum temperature (°C)	-47.390	-99.10	< 0.0001***
Average temperature (°C)	0.4484	56.60	< 0.0001***
Daily thermal excursion (°C)	-23.096	-90.10	0.049*
Daily precipitation (mm)	0.08901	9.30	< 0.0001***
RH	0.0253	2.56	0.65
Final model			
Intercept	1,793,297		< 0.0001***
Tavoliere Area	-0.9663	-62.00	< 0.0001***
Gargano Area (Hot spot Area)	0.6303	87.80	< 0.0001***
Murgia Area	-0.8327	-56.50	< 0.0001***
Salento Area	0.1299	13.87	< 0.0001***
Arco Jonico Area	-32.195	-96.00	< 0.0001***
Year	-0.0916	-62.90	< 0.0001***
Month	0.3133	36.80	< 0.0001***
Minimum temperature (°C)	-44.471	-98.80	< 0.0001***
Average temperature (°C)	0.4497	56.80	< 0.0001***
Daily thermal excursion (°C)	-20.121	-86.60	< 0.0001***
Daily precipitation (mm)	0.0882	9.20	< 0.0001***

*, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$.

however results of the two-to-two comparison between September/October and the other months only showed statistically significant difference against:

- January (Fisher's exact test: September, $p = 0.005$; October, $p = 0.0005$)
- March (Fisher's exact test: September, $p = 0.01$; October, $p = 0.002$)
- July (Pearson's X^2 test with Yates' continuity correction: September, $X^2 = 24.985$, $df = 1$, $p = 0.0005$; October, $X^2 = 21.596$, $df = 1$, $p = 0.003$)
- August (Pearson's X^2 test with Yates' continuity correction: September, $X^2 = 23.745$, $df = 1$, $p < 0.0001$; October, $X^2 = 27.015$, $df = 1$, $p = 0.0002$)
- December (Fisher's exact test: September, $p = 0.006$; October, $p = 0.019$)

The QQ plot is shown in Fig. 6. Neither the temperature (Fig. 6 A) (SWt-W statistic = 0.97756, $p = 0.002$; KSt-D statistic = 0.99512, $p < 0.0001$) nor the precipitation (Fig. 6 B) (SWt-W statistic = 0.45785, $p < 0.0001$, KSt-D statistic = 0.5, $p < 0.0001$) showed a normal distribution. Analysis of the RH (Fig. 6 C) produced conflicting results between the different tests (SWt-W statistic = 0.98201, $p = 0.05894$, KSt-D statistic = 1, $p < 0.0001$).

A Poisson regression model was used to conduct multivariate analysis of risk factors for LD in accordance to the model used by Conza et al. (2013). The results of these analyses are reported in Table 4.

The results of multivariate analysis (Table 4) showed that daily precipitation (mm) and average temperature (°C) directly influenced the RR of LD (9.2% increase in the RR of LD for every mm of precipitation and 56.8% increase for every 1 °C of temperature). Moreover, the results of multivariate analysis identified an inversely proportional relationship between minimum temperature (°C) and daily thermal excursion and the RR of LD (98.8% decrease in the RR of LD for every 1 °C of temperature and a 86.6% decrease for every 1 °C change in

thermal excursion).

In the multivariate study, the higher RR of LD is confirmed in the hot spot area (Gargano).

Analysis of the climatic parameters of different areas of Apulia (Portale della Protezione Civile Puglia, available at <http://www.protezionecivile.puglia.it/>) (Fig. 7A–D), showed that the hot spot area for LD (Gargano) had very low minimal temperatures and thermal excursion, but had the highest daily precipitation (Table 5A–D).

Based on 2001–2017 Regional Civil Protection data on daily precipitation, autumn was the wettest out of the four seasons (Kruskal-Wallis $X^2 = 286.48$, $df = 11$, $p < 0.001$; pairwise comparisons using a Wilcoxon rank sum test: $p < 0.001$ compared to with all the other seasons).

The median daily precipitation for autumn was 2.4 mm (IQR = 2.1) compared with 2 mm (IQR = 1.8) in winter, 1.7 mm in spring (IQR = 1.6) and 0.5 mm in summer (IQR = 1.2). Interestingly, the number of cases of LD relative to tourist arrival numbers was lower in the summer months of July and August compared with the number calculated for autumn months of September and October (Table 2). In autumn, there is a 6 °C–7 °C decrease in the median minimum temperature, with a simultaneous reduction of the daily thermal excursion of almost 3 °C. Most importantly though, this is coupled with an increase in median daily precipitation, from 0.2 mm in August to 2.1 mm in September, with similar levels maintained in October and November.

Taking into account only the climatic parameters showing statistically significant differences, we then examined areas with hot spots of LD cases (99 observations) versus other areas of the Apulia region (111 observations) over the days corresponding to the incubation period of LD. A statistically significant difference was identified among the climatic factors of the different areas (Table 6).

In the areas with LD cases hot spots, only the average temperature data was in contrast to the findings of the risk factor analysis shown in Table 4. However, given the percentage of RR shown in Table 4, the other climatic factors appeared to largely compensate for this discrepancy.

4. Discussion

Previous studies have investigated the impacts of climate change, including higher water temperature, drought, higher rainfall, rising sea levels and flooding, on the transmission of infectious disease (Hicks et al., 2007; Fisman et al., 2005; Karagianis et al., 2009; Ricketts et al., 2009; Walker, 2018). Recently investigations into the effects of climatic parameters on the notification rates of community-acquired LD in four European countries, including Italy (Beauté et al., 2016), confirmed that the growth of *Legionella* in natural environments is enhanced by warmer temperatures (De Giglio et al., 2019).

Our results showed that only temperature excursion affected the proliferation of *Legionella*.

In comparison, the role of temperature appears to be more complex because water flowing in artificial water networks is affected by the external temperature. An increase in seasonal temperatures can cause proliferation of *Legionella* in the cold water supply of a building and, consequently, a greater risk of disease. Therefore, designers and engineers must guarantee an artificial network system able to protect water from external thermal variations so as to avoid microbial growth in the built environment (Walker, 2018).

Conversely, our results also showed the highest risk LD was associated with decreases in minimum temperature. As reported previously (Beauté et al., 2016; Wullings and van der Kooij, 2006; Wadowsky et al., 1985; Carvalho et al., 2008), it may be that *Legionella* or its protozoan hosts have a lower optimum growth temperature and range in some regions or in certain climates.

The effects of the studied environmental factors on the protozoan hosts of *Legionella* may also play a role in the incidence of LD, which should be investigated further. In addition, *Legionella* is not only found


in natural environments but are also present in manmade water systems where the temperature can be artificially modified (Beauté et al., 2016).

Our data indicated that decreases in temperature excursion and increases in daily precipitation can affect the risk of LD. While the effect of thermal excursion has been poorly documented, many authors have reported that greater daily precipitation increases the number of cases

of LD (Simmering et al., 2017; Beauté et al., 2016; Halsby et al., 2014; Nai-Tzu Chen et al., 2014; Brandsema et al., 2014; Conza et al., 2013; Dunn et al., 2013; Garcia-Vidal et al., 2013; Karagiannis et al., 2009; Ricketts et al., 2009; Hicks et al., 2007).

Several studies (Brandsema et al., 2014) have reported that during milder climatic periods, the risk of LD increases as a result of increased

(A) Minimum temperature (°C) in the climatic area of Apulia region (month by month, 2001-2017 years)



Kruskal-Wallis chi-squared = 60.524, df = 5, p<0.001

Table 5 A. Minimum temperature (°C), pairwise comparisons using Wilcoxon rank sum test (p)

Area	Tavoliere	Arco Jonico	Murgia	Gargano
Tavoliere				
Arco Jonico	0.0002			
Murgia	0.05	<0.0001		
Gargano	0.28	<0.0001	0.37	
Salento	0.28	0.004	0.003	0.036

(B) Average temperature (°C) in the climatic areas of Apulia region (month by month, 2001-2017 years)





Fig. 7. Box plot of climatic parameters for five climatic areas of Apulia by month (2001-2017): (A) minimum and (B) average temperature, (C) daily thermal excursion and (D) daily precipitation.

Table 5 B. Average temperature (°C), pairwise comparisons using Wilcoxon rank sum test (*p*)

Area	Tavoliere	Arco Jonico	Murgia	Gargano
Tavoliere				
Arco Jonico	0.085			
Murgia	0.018	<0.0001		
Gargano	0.003	<0.0001	0.59	
Salento	0.52	0.26	0.002	0.00036

(C) Daily thermal excursion (°C) in the climatic area of Apulia region (month by month, 2001-2017 years)




Kruskal-Wallis chi-squared = 456.25, df = 5, *p* < 0.001

Table 5 C. Daily thermal excursion (°C), pairwise comparisons using Wilcoxon rank sum test (*p*)

Area	Tavoliere	Arco Jonico	Murgia	Gargano
Tavoliere				
Arco Jonico	<0.0001			
Murgia	0.0006	<0.0001		
Gargano	<0.0001	0.001	<0.0001	
Salento	0.06	<0.0001	0.029	<0.0001

(D) Daily precipitation (mm) in the climatic area of Apulia region (month by month, 2001-2017 years)

Fig. 7. (continued)



Kruskal-Wallis chi-squared = 61.18, df = 5, $p < 0.001$

Table 5 D. Daily precipitation (mm), pairwise comparisons using Wilcoxon rank sum test (p)

Area	Tavoliere	Arco Jonico	Murgia	Gargano
Tavoliere				
Arco Jonico	0.91			
Murgia	0.006	0.04		
Gargano	<0.0001	0.001	<0.0001	
Salento	0.47	0.54	0.17	<0.0001

Fig. 7. (continued)

precipitation and, consequently, increased RH. In agreement with other reports (Beauté et al., 2016; Conza et al., 2014; Chen et al., 2014), our study did not show a significant correlation with RH. This may be because to influence the survival of *Legionella*, the RH levels during the incubation period of the disease must reach at least 80% (Berendt, 1980) which would be uncommon in the Apulia region. Some authors (Hicks et al., 2007) hypothesised that heavy rain can favour the proliferation of other microorganisms or increase organic sediments in the water network which, in turn, aid the growth of *Legionella*. In fact, *L. pneumophila* has been found in rainwater on roads and in roof collection systems, as well as in floodwater (Brandsema et al., 2014; Sakamoto et al., 2009; Ahmed et al., 2014; Schets et al., 2010). Thus, it is likely that when the outside temperature exceeds 25 °C, the bacteria can be dispersed in the air from wet roads or from vehicles driving through puddles and generating contaminated aerosols. Similar to a study of travel-related LD in Glasgow, Scotland (Bhopal and Fallon, 1991), our data showed a greater number of cases in autumn (September, October), when precipitation was increased compared with the rest of the year.

The geostatistical analysis conducted in the current study revealed that the area with the highest *Legionella* load in water systems coincided with the area presenting the highest number of LD cases associated with staying in touristic-recreational facilities.

Studies conducted to date have applied geostatistical methods to public health issues such as the presence of heavy metals, soil insecticides, chemical contaminants, pollens, insects, plants and protozoa in different environments (Fortes Bde et al., 2004; Goovaerts, 2011; Li et al., 2006; Hou et al., 2017; Núñez-Alonso et al., 2019). Only Laganà et al. (2015) used geostatistical analysis to evaluate the distribution of *Legionella* in a healthcare facility. We applied this method throughout a geographical area, analysing tourist facilities and assessing the effects of climate on the incidence of LD.

The hot spot for LD cases in the Apulia region may be caused by the greater number of religious tourists (Castellana Pastoris et al., 2002) with known risk factors (such as older age, male sex, chronic lung disease and various conditions associated with immunodeficiency) to this area (Ginevra et al., 2009). Travelers are particularly at risk because they may stay in accommodation contaminated with *Legionella*, and the response to infection may not be supported by appropriate and timely diagnostic tests and medical treatment (Mouchtouri and Rudge, 2015). Legionellosis is a significant public health problem, especially if we consider that the number of cases notified each year represents only a fraction of the real number of cases as a result of several limitations: i) surveillance systems are not always adequately applied in some countries (Montagna et al., 2018), and ii) although *Legionella* is widely distributed in the environment, growing it in a laboratory setting is

Table 6
Climatic factors in areas with hot spots of LD cases versus other areas of the Apulia region (2001–2017 years).

Climatic factor	Area with hot spot of LD cases [median (A) ± IQR]	Other areas of the Apulia region [median (B) ± IQR]	Wilcoxon rank sum test with continuity correction (W)	p-value	RR (%) (from Table 4, final model) (C)	Higher RR (%) in the area of Hot spot vs other area of Apulia Region ^a
Minimum temperature (°C)	12.0 ± 6.4	16.8 ± 6.5	14,772	0.005	-98.80	+474.3
Average temperature (°C)	16.5 ± 8.0	22.5 ± 6.3	7256.5	< 0.0001	56.80	-340.8
Daily thermal excursion (°C)	10.4 ± 3.2	11.6 ± 5.5	6149.5	0.005	-86.60	+103.92
Daily precipitation (mm)	0.4 ± 3.6	0.0 ± 0.6	3883.5	0.003	9.20	+3.68

IQR = interquartile range.

^a calculated on the basis of the following formula: [(A-B) x C].

challenging because of its intracellular lifestyle and dependency on the host (Graells et al., 2018). For this reason, culture-independent methods are necessary to detect species other than *Legionella pneumophila* sg 1, which is still considered the primary causative agent of LD (Palmer et al., 2016; Kocazeybek et al., 2016; Relich et al., 2018). Currently, > 90% of cases are diagnosed using a urinary antigen that mainly detects *Legionella pneumophila* sg 1 strains. In the last decade, several seroprevalence studies have shown a potential ongoing exposure to other serogroups and *Legionella* species (Lee et al., 2008; Rudbeck et al., 2008). Consequently, there has recently been a substantial effort made to characterise the virulence determinants of *Legionella pneumophila* sg 1 strains, including secretory effectors or surface factors (Mercante and Winchell, 2015). Conflicting data has also been reported for the Apulia region. For example, a previous surveillance study of health and community facilities (Napoli et al., 2010) showed that clinical LD cases caused by *Legionella pneumophila* sg 1 strains did not coincide with its distribution in aquatic environments, which were often contaminated with other *Legionella* species or serogroups (De Giglio et al., 2019). In comparison, the current study, which concerns only the cases of LD associated with staying in tourist-recreational facilities, showed that *Legionella pneumophila* sg 1 was the most frequent species and serogroup both in environmental samples and in clinical samples. This discrepancy confirms the need to improve laboratory diagnostics techniques to provide an epidemiological framework that is more responsive to changes in real-time.

5. Conclusion

Our study highlights the influences of climatic parameters on the increased risk of LD during periods of higher rainfall. Public health physicians should be aware of this phenomenon and consider more appropriate diagnostic and therapeutic protocols in cases of travel-associated pneumonia. Furthermore, the innovative geo-statistical approach used in this study should be further implemented to evaluate the distribution of *Legionella* and legionellosis cases in other areas, taking climatic influences into account. Such measures would more precisely define where and when it is necessary to intervene with preventative measures to protect the health of tourists.

Conflict of interest

The authors declare they have no actual or potential competing financial interests.

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Acknowledgement

We thank Kate Fox, DPhil, and Tamsin Sheen, Phd, from Edanz Group (www.edanzediting.com/ac) for editing a draft of this manuscript.

References

- Ahmed, W., Brandes, H., Gyawali, P., Sidhu, J.P., Toze, S., 2014. Opportunistic pathogens in roof-captured rainwater samples, determined using quantitative PCR. *Water Res.* 53, 361–369.
- Beauté, J., Sandini, S., Uldum, S.A., Rota, M.C., Brandsema, P., Giesecke, J., 2016. Short-term effects of atmospheric pressure, temperature, and rainfall on notification rate of community-acquired Legionnaires' disease in four European countries. *Epidemiol. Infect.* 144, 3483–3493.
- Berendt, R.F., 1980. Survival of *Legionella pneumophila* in aerosols: effect of relative humidity. *J. Infect. Dis.* 141, 689.
- Bhopal, R.S., Fallon, R.J., 1991. Seasonal variation of Legionnaires' disease in Scotland. *J. Infect.* 22, 153–160.

- Borella, P., Montagna, M.T., Stampi, S., Stancanelli, G., Romano-Spica, V., Triassi, M., Marchesi, I., Bargellini, A., Tatò, D., Napoli, C., Zanetti, F., Leoni, E., Moro, M., Scaltriti, S., Ribera Alcalá, G., Santarpia, R., Boccia, S., 2005. *Legionella* contamination in hot water of Italian hotels. *Appl. Environ. Microbiol.* 71, 5805–5813.
- Brandsema, P.S., Euser, S.M., Karagiannis, I., Den Boer, J.W., Van Der Hoek, W., 2014. Summer increase of Legionnaires' disease 2010 in The Netherlands associated with weather conditions and implications for source finding. *Epidemiol. Infect.* 142 (11), 2360–2371.
- Carvalho, F.R.S., Nastasi, F.R., Gamba, R.C., Foronda, A.S., Pellizari, V.H., 2008. Occurrence and diversity of Legionellaceae in polar Lakes of the Antarctic Peninsula. *Curr. Microbiol.* 57, 294–300.
- Castellani Pastoris, M., Ricci, M.L., Rota, M.C., 2002. A community cluster of legionnaires' disease among pilgrims to San Giovanni Rotondo, Puglia, Italy. *Weekly releases (1997–2007)*. 6 (27).
- Chang, K., 2010. *Introduction to Geographic Information Systems*, vol. 5. Thomas D. Timp, New York, pp. 327–340.
- Chen, N.T., Chen, M.J., Guo, C.Y., Chen, K.T., Su, H.J., 2014. Precipitation increases the occurrence of sporadic legionnaires' disease in Taiwan. *PLoS One* 9 (12), e114337.
- Conza, L., Casati, S., Limoni, C., Gaia, V., 2013. Meteorological factors and risk of community-acquired Legionnaires' disease in Switzerland: an epidemiological study. *BMJ Open* 3 (3), e002428.
- Conza, L., Casati Pagani, S., Gaia, V., 2014. Influence of climate and geography on the occurrence of *Legionella* and amoebae in composting facilities. *BMC Res. Notes* 7, 831.
- Costa, J., da Costa, M.S., Veríssimo, A., 2010. Colonization of a therapeutic spa with *Legionella* spp: a public health issue. *Res. Microbiol.* 161 (1), 18–25.
- De Giglio, O., Barbuti, G., Trerotoli, P., Brigida, S., Calabrese, A., Di Vittorio, G., Lovero, G., Caggiano, G., Uricchio, V.F., Montagna, M.T., 2016. Microbiological and hydro-geological assessment of groundwater in Southern Italy. *Environ. Monit. Assess.* 188, 638.
- De Giglio, O., Napoli, C., Apollonio, F., Brigida, S., Marzella, A., Diella, G., Calia, C., Scrascia, M., Pacifico, C., Pazzani, C., Uricchio, V.F., Montagna, M.T., 2019. Occurrence of *Legionella* in groundwater used for sprinkler irrigation in Southern Italy. *Environ. Res.* 170, 215–221.
- Download QGIS version 3.2 Available online: <https://qgis.org/it/site/forusers/download.html>, Accessed date: 16 March 2019.
- Dunn, C.E., Rowlingson, B., Bhopal, R.S., Diggle, P., 2013. Meteorological conditions and incidence of Legionnaires' disease in Glasgow, Scotland: application of statistical modelling. *Epidemiol. Infect.* 141 (4), 687–696.
- European Centre for Disease Prevention and Control (ECDC), 2018. Legionnaires' Disease. ECDC. Annual Epidemiological Report for 2016. ECDC, Stockholm.
- Euzéby, J.P., 2019. List of Prokaryotic Names with Standing in Nomenclature - Genus *Legionella*. <http://www.bacterio.cict.fr/l/legionella.html>, Accessed date: 16 March 2019.
- Fields, B.S., 1996. The molecular ecology of legionellae. *Trends Microbiol.* 4, 286–290.
- Fisher, N.L., Lewis, T., Embleton, B.J.J., 1987. *Statistical Analysis of Spherical Data*. Cambridge University Press, Cambridge.
- Fisman, D.N., Lim, S., Wellenius, G.A., Johnson, C., Britz, P., Gaskins, M., Maher, J., Mittleman, M.A., Spain, C.V., Haas, C.N., Newbern, C., 2005. It's not the heat, it's the humidity: wet weather increases legionellosis risk in the greater Philadelphia metropolitan area. *J. Infect. Dis.* 192 (12), 2066–2073.
- Fortes Bde, P., Ortiz Valencia, L.I., Ribeiro Sdo, V., Medronho Rde, A., 2004. Geostatistical modeling of *Ascaris lumbricoides* infection. *Cad Saude Publica* 20 (3), 727–734.
- García-Vidal, C., Labori, M., Viasus, D., Simonetti, A., Garcia-Somoza, D., Dorca, J., Gudiol, F., Carratala, J., 2013. Rainfall is a risk factor for sporadic cases of *Legionella pneumophila* pneumonia. *PLoS One* 8 (4), e61036. <https://doi.org/10.1371/journal.pone.0061036>.
- Garrity, G.M., Bell, J.A., Tilburn, T., 2005. Order VI Legionellales family 1. Legionellaceae. In: Brenner, D.J., Krieg, N.R., Staley, J.T. (Eds.), *Bergey's Manual of Systematic Bacteriology*, second ed. vol. 2. Part B. Springer-Verlag, New York, NY, pp. 210–237.
- Getis, A., Ord, J.K., 1992. The analysis of spatial association by use of distance statistics. *Geogr. Anal.* 24, 189–206.
- Getis, A., Ord, J.K., 1996. Local spatial statistics: an overview. In: Longley, P., Batty, M. (Eds.), *Spatial Analysis: Modeling in a GIS Environment*. Geo Information International, Cambridge, pp. 269–294.
- Ginevra, C., Ducloux, A., Vanhems, P., Campès, C., Forey, F., Lina, G., Che, D., Etienne, J., Jarraud, S., 2009. Host-related risk factors and clinical features of community-acquired legionnaires disease due to the Paris and Lorraine endemic strains, 1998–2007. *France. Clin. Infect. Dis.* 49 (2), 184–191.
- Goovaerts, P., 2011. A coherent geostatistical approach for combining choropleth map and field data in the spatial interpolation of soil properties. *Eur. J. Soil Sci.* 62 (3), 371–380.
- Graells, T., Ishak, H., Larsson, M., Guy, L., 2018. The all-intracellular order Legionellales is unexpectedly diverse, globally distributed and lowly abundant. *FEMS Microbiol. Ecol.* 94 (12), fty185.
- Halsby, K.D., Joseph, C.A., Lee, J.V., Wilkinson, P., 2014. The relationship between meteorological variables and sporadic cases of Legionnaires' disease in residents of England and Wales. *Epidemiol. Infect.* 142 (11), 2352–2359.
- Haupt, T.E., Heffernan, R.T., Kazmierczak, J.J., Nehls-Lowe, H., Rheineck, B., Powell, C., Leonhardt, K.K., Chitnis, A.S., Davis, J.P., 2012. An outbreak of Legionnaires disease associated with a decorative water wall fountain in a hospital. *Infect. Control. Hosp. Epidemiol.* 33, 185–191.
- Hicks, L.A., Rose Jr., C.E., Fields, B.S., Drees, M.L., Engel, J.P., Jenkins, P.R., Rouse, B.S., Blythe, D., Khalifah, A.P., Feikin, D.R., Whitney, C.G., 2007. Increased rainfall is associated with increased risk for legionellosis. *Epidemiol. Infect.* 135 (5), 811–817.
- Hou, D., O'Connor, D., Nathanail, P., Tian, L., Ma, Y., 2017. Integrated GIS and multi-variate statistical analysis for regional scale assessment of heavy metal soil contamination: a critical review. *Environ. Pollut.* 231 (Pt 1), 1188–1200.
- Iatta, R., Cuna, T., De Giglio, O., Montagna, M.T., Napoli, C., 2013a. Sequence based typing of *Legionella pneumophila* sg 1 isolated in nosocomial acquired infections in Apulia, Southern Italy. *J. Prev. Med. Hyg.* 54, 167–169.
- Iatta, R., Cuna, T., Napoli, C., De Giglio, O., Montagna, M.T., 2013b. Environmental surveillance and molecular investigation of *Legionella* spp. in Apulia, in the years 2008–2011. *Ann. Ig.* 25, 435–441.
- ISO 11731, 1998. Water Quality – Detection and Enumeration of *Legionella*.
- ISPRa, Regione Puglia, 2014. Ministero delle Infrastrutture e dei Trasporti. The Groundwater and the Seawater Intrusion in Apulia: from Research to the Emergency in the Safeguard of the Water Resource, vol. 92.
- ISPRa, 2014. Valori Climatici Normali di Temperatura e precipitazione in Italia, vol. 55 Available online: http://www.isprambiente.gov.it/files/pubblicazioni/statoambiente/copy_of_SA_55_14_Valori_climatici_normali.pdf.
- Istituto Superiore di Sanità (ISS), 2018. Rapporto Annuale Sulla Legionellosi in Italia Nel 2017, vol. 31. Notiziario ISS, Rome, Italy, pp. 7–12.
- Karagiannis, I., Brandsema, P., van der Sande, M., 2009. Warm, wet weather associated with increased Legionnaires' disease incidence in The Netherlands. *Epidemiol. Infect.* 137, 181–187.
- Kocazeybek, B.S., Yuksel, P., Keskin, D., Sheikh, S., Habib, Z., Yavuzer, S.S., Caliskan, R., Altun, Y.M., Kuskucu, M., Cengiz, M., Dinc, H.O., Karakullukcu, A., Ergin, S., Saribas, S., Yilmaz, N., Tokman, H.B., 2016. Travel-associated infections caused by unusual serogroups of *Legionella pneumophila* identified using *Legionella* BIOCHIP slides in Turkey and Iraq. *Trav. Med. Infect. Dis.* 14 (3), 248–254.
- Laganà, P., Moscato, U., Poscia, A., La Milia, D.I., Boccia, S., Avventuroso, E., Delia, S., 2015. Geostatistics - a tool applied to the distribution of *Legionella pneumophila* in a hospital water system. *Ann. Agric. Environ. Med.* 22 (4), 655–660.
- Lee, H.K., Woo, M.K., Ju, Y.I., Baek, S.J., Song, H.J., Choi, J.S., Kweon, S.S., Jeon, D.Y., Kang, Y.H., 2008. Prevalence of antibodies in response to *Legionella* species, analysis of a healthy population from Jeollanam-do Province, Korea. *J. Microbiol.* 46, 160–164.
- Li, B.G., Cao, J., Liu, W.X., Shen, W.R., Wang, X.J., Tao, S., 2006. Geostatistical analysis and kriging of hexachlorocyclohexane residues in topsoil from Tianjin, China. *Environ. Pollut.* 142 (3), 567–575.
- Linee, 2000. Linee Guida per la Prevenzione ed il Controllo della Legionellosi. Available online: http://www.salute.gov.it/imgs/C_17_normativa_1601_allegato.pdf, Accessed date: 16 March 2019.
- Linee Guida per la Prevenzione ed il Controllo della Legionellosi. Available online: http://www.salute.gov.it/imgs/C_17_pubblicazioni_2362_allegato.pdf, Accessed date: 16 March 2019.
- Mercante, J.W., Winchell, J.M., 2015. Current and emerging *Legionella* diagnostics for laboratory and outbreak investigations. *Clin. Microbiol. Rev.* 28 (1), 95–133.
- Montagna, M.T., Napoli, C., Tatò, D., Spilotros, G., Barbuti, G., Barbuti, S., 2006. Clinical-environmental surveillance of legionellosis: an experience in Southern Italy. *Eur. J. Epidemiol.* 21, 325–331.
- Montagna, M.T., Cuna, T., De Giglio, O., Bollino, A., Iatta, R., Napoli, C., 2012. Spa facilities and legionellosis: legislative structure and state of the art in Apulia (Italy). *Ig Sanita Pubbl* 68 (5), 719–731.
- Montagna, M.T., Cristina, M.L., De Giglio, O., Spagnolo, A.M., Napoli, C., Cannova, L., Deriu, M.G., Delia, S.A., Giuliano, A., Guida, M., Laganà, P., Liguori, G., Mura, I., Pennino, F., Rossini, A., Tardivo, S., Torre, I., Torregrossa, M.V., Villafrate, M.R., Albertini, R., Pasquarella, C., 2016. Serological and molecular identification of *Legionella* isolated from water and surrounding air samples in Italian healthcare facilities. *Environ. Res.* 146, 47–50.
- Montagna, M.T., De Giglio, O., Cristina, M.L., Albertini, R., Pasquarella, C., GISIO, AIA and SIMPIOS WorkingGroups, 2017. *Legionella* indoor air contamination in health care environments. In: In: Capolongo, S., Settimo, G., Gola, M. (Eds.), *Policentrico di Milano, Milan, Italy. Indoor Air Quality in Healthcare Facilities*, vol. 6 SpringerBriefs in Public Health.
- Montagna, M.T., De Giglio, O., Cristina, M.L., Napoli, C., Pacifico, C., Agodi, A., Baldovin, T., Casini, B., Coniglio, M.A., D'Errico, M.M., Delia, S.A., Deriu, M.G., Guida, M., Laganà, P., Liguori, G., Moro, M., Mura, I., Pennino, F., Privitera, G., Romano Spica, V., Sembini, S., Spagnolo, A.M., Tardivo, S., Torre, I., Valeriani, F., Albertini, R., Pasquarella, C., 2017b. Evaluation of *Legionella* air contamination in healthcare facilities by different sampling methods: an Italian multicenter study. *Int. J. Environ. Res. Public Health* 14, 670.
- Montagna, M.T., De Giglio, O., Napoli, C., Diella, G., Rutigliano, S., Agodi, A., Auxilia, F., Baldovin, T., Bisetto, F., Arnoldo, L., Brusaferrò, S., Busetti, M., Calagreti, G., Casini, B., Cristina, M.L., Di Luzio, R., Fiorio, M., Formoso, M., Liguori, G., Martini, E., Molino, A., Mondello, P., Mura, I., Novati, R., Orsi, G.B., Patroni, A., Poli, A., Privitera, G., Ripabelli, G., Rocchetti, A., Rose, F., Sarti, M., Savini, S., Silvestri, A., Sodano, L., Spagnolo, A.M., Tardivo, S., Teti, V., Torregrossa, M.V., Torri, E., Veronesi, L., Zarrilli, R., Pacifico, C., Goglio, A., Moro, M., Pasquarella, C., 2018. Control and prevention measures for legionellosis in hospitals: a cross-sectional survey in Italy. *Environ. Res.* 166, 55–60.
- Mouchtouri, V.A., Rudge, J.W., 2015. Legionnaires' disease in hotels and passenger ships: a systematic review of evidence, sources, and contributing factors. *J. Travel Med.* 22 (5), 325–337.
- Napoli, C., TATO, D., Iatta, R., Montagna, M.T., 2007. [Assessment of occupational risk of *Legionella* spp. infection among dental health-care personnel]. *Ig. Sanita. Pubbl* 63, 683–689 [Article in Italian].
- Napoli, C., Iatta, R., Fasano, F., Marsico, T., Montagna, M.T., 2009a. Variable bacterial load of *Legionella* spp. in a hospital water system. *Sci. Total Environ.* 408, 242–244.
- Napoli, C., Iatta, R., Fasano, F., Marsico, T., Montagna, M.T., 2009b. [Legionella e

- legionellosi in Puglia: sorveglianza negli anni 2000-2008]. *Ann. Ig* 21, 181–188 [Article in Italian].
- Napoli, C., Fasano, F., Iatta, R., Barbuti, G., Cuna, T., Montagna, M.T., 2010. *Legionella* spp. and legionellosis in Southeastern Italy: disease epidemiology and environmental surveillance in community and health care facilities. *BMC Public Health* 10, 660.
- Núñez-Alonso, D., Pérez-Arribas, L.V., Manzoor, S., Cáceres, J.O., 2019. Statistical tools for air pollution assessment: multivariate and spatial analysis studies in the Madrid region. *J. Anal. Methods Chem.* 2019, 9753927.
- Ord, J.K., Getis, A., 1995. Local spatial autocorrelation statistics: distributional issues and an application. *Geogr. Anal.* 27, 286–306.
- Palmer, A., Painter, J., Hassler, H., Richards, V.P., Bruce, T., Morrison, S., Brown, E., Kozak-Muiznieks, N.A., Lucas, C., McNealy, T.L., 2016. *Legionella clemsonensis* sp. nov.: a green fluorescing *Legionella* strain from a patient with pneumonia. *Microbiol. Immunol.* 60 (10), 694–701.
- Phin, N., Parry-Ford, F., Harrison, T., Stagg, H.R., Zhang, N., Kumar, K., Lortholary, O., Zumla, A., Abubakar, I., 2014. Epidemiology and clinical management of Legionnaires' disease. *Lancet Infect. Dis.* 14 (10), 1011–1021.
- Portale dell'Open Data Puglia Available online: <http://www.dataset.puglia.it/>, Accessed date: 16 March 2019.
- Portale dell'Agenzia REgionale del Turismo PUGLIAPROMOZIONE. Report Data 2017, Available online: <https://www.agenziapugliapromozione.it/portal/documents/10180/2920246/Dati%202017>.
- Portale della Protezione Civile Puglia. Available online: <http://www.protezionecivile.puglia.it/>. (Accessed 16 March 2019).
- Portale dell'Osservatorio Epidemiologico regionale. Available online: <https://www.sanita.puglia.it/osservatorio-epidemiologico-regionale>. (Accessed 16 March 2019).
- Portale ufficiale del turismo della Regione Puglia. Available online: <https://www.viaggiareinpuglia.it/hp/it> (Accessed 16 March 2019).
- Regione Puglia, 2015. Indirizzi operativi per la prevenzione e il controllo della legionellosi nelle strutture turistico ricettive ed ad uso collettivo della Regione Puglia.
- Regione Puglia, 2018. Indirizzi operativi per la sorveglianza clinica e ambientale della legionellosi nelle strutture sanitarie ed assistenziali della Regione Puglia.
- Relich, R.F., Schmitt, B.H., Raposo, H., Barker, L., Blosser, S.J., May, M., 2018. *Legionella indianapolisensis* sp. nov., isolated from a patient with pulmonary abscess. *Int. J. Infect. Dis.* 69, 26–28.
- Ricketts, K.D., Yadav, R., Rota, M.C., Joseph, C.A., European Working Group for Legionella Infections, 2010. Characteristics of reoffending accommodation sites in Europe with clusters of Legionnaires disease, 2003-2007. *Euro Surveill.* 15 (40), 19680.
- Ricketts, K.D., Charlett, A., Gelb, D., Lane, C., Lee, J.V., Joseph, C.A., 2009. Weather patterns and Legionnaires' disease: a meteorological study. *Epidemiol. Infect.* 137 (7), 1003–1012 Erratum in: *Epidemiol. Infect.* 2011, 139(9), 1443.
- Rudbeck, M., Molbak, K., Uldum, S., 2008. High prevalence of antibodies to *Legionella* spp. in Danish blood donors. A study in areas with high and average incidence of Legionnaires' disease. *Epidemiol. Infect.* 136 (2), 257–262.
- Sakamoto, R., Ohno, A., Nakahara, T., Satomura, K., Iwanaga, S., Kouyama, Y., Kura, F., Noami, M., Kusaka, K., Funato, T., Takeda, M., Matsubayashi, K., Okumiya, K., Kato, N., Yamaguchi, K., 2009. Is driving a car a risk for Legionnaires' disease? *Epidemiol. Infect.* 137 (11), 1615–1622.
- Sánchez-Martín, J.-M., Rengifo-Gallego, J.-I., Blas-Morato, R., 2019. Hot spot analysis versus cluster and outlier analysis: an enquiry into the grouping of Rural accommodation in Extremadura (Spain). *ISPRS Int. J. Geo-Inf.* 8, 176 2019.
- Schets, F.M., Italiaander, R., van den Berg, H.H., de Roda Husman, A.M., 2010. Rainwater harvesting: quality assessment and utilization in The Netherlands. *J. Water Health* 8, 224–235.
- Simmering, J.E., Polgreen, L.A., Hornick, D.B., Sewell, D.K., Polgreen, P.M., 2017. Weather-dependent risk for Legionnaires' disease, United States. *Emerg. Infect. Dis.* 23 (11), 1843–1851. www.cdc.gov/eid.
- Wadowsky, R.M., Wolford, R., McNamara, A.M., Yee, R.B., 1985. Effect of temperature, pH, and oxygen level on the multiplication of naturally occurring *Legionella pneumophila* in potable water. *Appl. Environ. Microbiol.* 49, 1197–1205.
- Walker, J.T., 2018. The influence of climate change on waterborne disease and *Legionella*: a review. *Perspect. Public Heal.* 138 (5), 282–286.
- Wullings, B.A., van der Kooij, D., 2006. Occurrence and genetic diversity of uncultured *Legionella* spp. in drinking water treated at temperatures below 15 degrees C. *Appl. Environ. Microbiol.* 72, 157–166.