

Religious stigmata as malingering artifact

Report of a case and review of the literature

Domenico Bonamonte, MD, PhD^a, Michelangelo Vestita, MD^{b,*}, Angela Filoni, MD^a,
Giuseppe Giudice, MD^b, Gianni Angelini, MD^a

Abstract

Rationale: Artifacts or simulated diseases are self-inflicted conditions caused by various means and for different purposes. Disease simulation can be motivated, among other things, by illegal purposes, to escape from civil duties or prison sentences, for example, or to exploit specific situations in order to receive a range of benefits. In such cases, the simulator is fully aware of his or her actions and intentions.

Diagnoses, Interventions and Outcomes: We report the case of a 42-year-old woman who, for 3 consecutive years, showed religious stigmata on the forehead and on the dorsal surface of hands and feet immediately before Easter. Lesions showed an acute onset, manifested as erosions and ulcerations, and healed a few days after Easter. Stigmata were immediately made public and every year faithful and curious people went in procession to the house of the patient, offering different kinds of gifts. After intervention of the police authority in the early days of the third episode, the patient and her family repented of their actions. Cutaneous lesions healed in few days and similar events did not occur during the following 2 years.

Lessons: Regardless of personal beliefs, the possibility that stigmata could be self-inflicted for illegal or profit purposes should always be considered.

Abbreviation: SISL = self-inflicted skin lesion.

Keywords: behavioral health, malingering, psychocutaneous, religious, stigmata, ulcers of the skin

1. Introduction

The issue about self-inflicted skin lesions (SISL)^[1] has been critically discussed in a recent position paper of the European Society for Dermatology and Psychiatry. SISL are divided into 2 main groups: the first group is represented by syndromes associated with a denied or hidden pathological behavior, with (malingering) or without (factitious disorders) external incentives; the second group is represented by syndromes associated with a non-denied and nonhidden pathological behavior (compulsive and impulsive disorders). Our case of religious stigmata belongs to the first group of SISL, associated with external incentives and profit purposes.

2. Case presentation

A 42-year-old woman, originally from a small town in Calabria (South of Italy), showed, for the third consecutive year, cutaneous lesions during Easter holidays. Such lesions appeared every year on the forehead and on the dorsal surface of hands and feet (Figs. 1–3). Lesions suddenly become manifest on Holy Monday and healed spontaneously within a few weeks. These alleged stigmata were made public, causing a steady flow of devotees and curious onlookers to the patient's house, offering gifts of various kinds. The year before the local ecclesiastical and police authorities had become interested in the singular event, but legal or religious actions were not undertaken as the stigmata had rapidly disappeared some days after Easter.

The patient came to our attention on the third day of clinical manifestations of the third year, accompanied by her general practitioner. At physical examination, a dozen mildly exudative erosions were present on the forehead, along the hairline, measuring 0.5 cm in diameter, regularly spaced 1 cm from each other. On the dorsal surface of the hands, halfway between carpus and metacarpus, an irregularly round erythematous-ulcerated lesion, swollen and moderately bleeding, was noticeable. The left-hand lesion was larger and more infiltrated. Similar erythematous-ulcerated elements (measuring 1.5 × 1 cm) involved symmetrically the central dorsum of the feet. No lesions were present on wrists, palms, soles, or elsewhere.

During consultation the patient was silent; she gave no specifics on time and mode of onset of lesions, and did not complain of any subjective symptoms. She also showed a wary and absent attitude, occasionally speaking in a detached manner about her religious faith. The psychiatric consultation, which had been refused at the time of the previous episodes, described a hysterical personality with attention-seeking behavior and excessive need

Editor: Dan Lipsker.

The authors have no conflicts of interest to disclose.

^a Section of Dermatology, Department of Biomedical Science and Human Oncology, ^b Unit of Plastic Surgery, Department of Emergencies and Organ Transplantation, University of Bari, Bari, Italy.

* Correspondence: Michelangelo Vestita, 11, Piazza Giulio Cesare, Bari 70124, Italy (e-mail: michelangelovestita@gmail.com).

Copyright © 2016 the Author(s). Published by Wolters Kluwer Health, Inc. All rights reserved.

This is an open access article distributed under the Creative Commons Attribution-NoDerivatives License 4.0, which allows for redistribution, commercial and non-commercial, as long as it is passed along unchanged and in whole, with credit to the author.

Medicine (2016) 95:49(e5354)

Received: 4 May 2016 / Received in final form: 15 October 2016 / Accepted: 18 October 2016

<http://dx.doi.org/10.1097/MD.0000000000005354>



Figure 1. Irregularly round erythematous-ulcerated lesion on hands dorsum.



Figure 2. Similar elements involving symmetrically feet dorsum.

for approval of her family. Moreover, her general practitioner reported us that the intervention of the police, the day before, had stopped the people pilgrimage to the house and had changed the attitude of the patient and her relatives; they showed fear and discretion, diminished the episode, and argued that most likely it would not have occurred again. The patient refused hospitalization but promised she would return for follow-up, but actually



Figure 3. Multiple erosions following the hairline. Notice the consistent diameter and spacing.

Table 1

Time-related milestones of the presented case.

Past episode—Easter, 2 y before
Lesion onset
Pilgrimage
Spontaneous healing in few weeks
Past episode—Easter, 1 y before
Lesion onset
Pilgrimage
Spontaneous healing in few weeks
Described episode—Easter
Lesion onset—day 1
Pilgrimage start—day 1
Police intervention—day 2
Our consultation—day 3
Psychiatric consultation—day 3
Almost complete spontaneous healing—day 6
Complete healing—day 22
No recurrence—1-y follow-up
No recurrence—2-y follow-up

never showed up. Her general practitioner later reported us that lesions were already healing 3 days after our consultation, with complete healing after 3 weeks. Stigmata no longer recurred during Easter holidays on the following 2 years. The time-related milestones concerning the case are summarized in Table 1.

3. Discussion

Different causes and purposes underlie SISL associated with denied or hidden pathological behaviors. Most of the time disease simulations are due to psychiatric problems, such as psychosis and personality disorders. The intrinsic motive of these lesions is a request for family and society attention, as a reaction to difficulties or adverse situations with involuntary cutaneous somatization, masochism with physical pain, and guilt or spiritual responses. As showed by the literature,^[1–9] cases of unconscious simulators, mainly women with chronic cutaneous diseases, belong to this group. Such patients always deny self-induced illness and seem unconcerned about the disease.

However, disease simulation can be motivated by illegal purposes, for example, to escape from specific types of duties (military duty, draft), to avoid detention, and to acquire benefits in an occupational environment (sickness leave extension, recognition as an occupational disease, higher class of disability). In these cases, simulators are aware of their actions and purposes.^[1,2,6,10–13] In our long-term practice, we have directly observed 46 patients affected by SISL with illegal purposes in occupational environments. These patients, overwhelmingly male, are affected by morphologically bizarre lesions, more so than those provoked by unaware pathomimia patients. Besides hands and forearms, unusual areas are involved and the most peculiar and unlikely causative agents are implicated.^[6,13] The certainty of conscious simulation, suspected by a positive history for job uneasiness, by lesions' particular morphology, and by repeated diagnostic tests' negativity, came from patients' admission after repeated friendly and confidential consultations.

Herein we report a rare case of religious stigmata. Affected sites, morphology, acute onset, and duration led to suspicion of an artificially induced disease. As a matter of fact, in our case hands and feet dorsum were affected without the involvement of palms and soles. Moreover, in religious texts, the originally described sites of upper limb involvement are the wrist and not

the hands. Stigmata were also immediately made public by the patient and her family in this case. Our hypothesis was indirectly confirmed by quick resolution of the dermatosis and no recurrence of the lesions during the following 2 years. Moreover, the patient's absent attitude, as though she was living in her inner world, could also underlie a hysterical personality that could be easily exploited by her relatives. This was confirmed by the psychiatric consultation. Skin lesions were probably caused by metal objects, by a mechanism of scraping (on the forehead), or overheating (most likely on the dorsal surface of hands and feet); however, the use of caustic chemicals cannot be excluded.

Differential diagnosis with dermatologic conditions, when singularly considering a lesion among those affecting hands and feet, included contact dermatitis, mycosis, and infective and noninfective granulomatosis; contact dermatitis to a hair accessory such as a frontlet or headband (perhaps with spaced metallic inserts) could have been responsible for the hairline manifestations. However, when assessing the clinical presentation as a whole, together with patient history, we believe that a differential diagnosis is not needed, as there is no single dermatologic condition that could justify the morphology, localization, symmetry, and evolution of such lesions.

To our knowledge, the topic of malingering religious stigmatization has rarely been treated in the medical literature. Kluger and Cribier thoroughly discussed the subject in light of modern medicine,^[14,15] and a few other cases similar to our own have been reported. In particular, a case of a 42-year-old man with a 5-month history of palmar ulcers at different healing stages has been described.^[16] During consultation the patient theatrically explained how he had been blessed by God with Christ's stigmata. However, the patient's partner revealed that such lesions were autoinflicted with a knife. Another case of a 23-year-old male patient with stigmata on the hands has been reported in a small town of Sicily. He showed some erosion on the dorsal surface of hands, and small bullous lesions arranged to create the word "pax" on the left palm. Previously, other odd events, promoted as divine, had occurred in the house of the young man (weeping of an effigy depicting the Madonna, weeping and bleeding of a Jesus Christ's picture, lugubrious sounds at night). Accordingly, a prayer hall in the garage of the house had been set up, complete of an altar and iconography of the Stations of the Cross. A flock of people brought gifts and pledges. The patient later turned out to be hysterical, with an easily swayed personality, and easily manipulated by relatives and acquaintances for profit aim. The intervention of the law authority stopped the lucrative affair.^[17] Early and Lifschutz observed a 10-year-old black Baptist girl during a 3-week period immediately preceding Easter Sunday.^[18] Psychiatric examination did not evidence psychological alterations except an indifference toward bleeding and auditory hallucinations of religious nature.

The issue concerning religious stigmata has always been of particular interest,^[6,15-19] given the complexity of interpretation and discussion. Stigmatized people present bleeding lesions in 5 distinct body sites: hands and feet because of nails (not on wrists, as evidenced by the Shroud of Turin), the ribs in relation to the thrust of the spear, the back (site of flagellation and pressure of the cross), and the forehead because of the crown of thorns. These are all known signs of the passion of Christ. Besides the visible ones, invisible stigmata, such as intense pain affecting the above-mentioned sites, should be considered. The latter could precede or represent the outcome of visible stigmata, but may also be isolated, without any means to ascertain their presence.^[20]

Saint Francis of Assisi (1182–1226) was the first person to receive stigmata, unless Saint Paul implied that he had them when he wrote, "I bear the marks of the Lord Jesus in my body."^[21] Saint Francis developed the stigmata during an ecstatic vision in 1224^[15,19]; these were described 2 years later in his first biography.^[22] The Saint Francis's stigmata were immediately acknowledged by the papacy, despite minor opposition among the clergy, which accused the Saint of heresy.^[15,23] Kluger and Cribier reported 3 other cases of autoinflicted stigmata, which dated back to the XIII century.^[15] During the following centuries, >300 cases of stigmatization were described, most of which were fervent Roman Catholics.^[15] In 1908, Imbert-Gourbeyre found that of the 321 people with stigmata described afterwards Saint Francis, 41 were men and 280 were women during their reproductive years.^[24]

They mainly came from Italy (229 cases), and then France, Spain, Germany, Belgium, Portugal, Switzerland, Holland, Hungary, and Peru.

Among the most famous stigmatized of the XIX century, Louise Lateam (1850–1883) stands out, a Belgian girl who bled every Friday except for 2 from 1868 to 1883; she was visited by various eminent Belgian doctors.^[25] Also Thérèse Neumann (1898–1962), a German peasant girl, developed the stigmata in 1926 during an ecstatic vision; her hands and feet bled regularly on Fridays until her death.^[26] Finally Saint Pio of Petralcina (1887–1968) had his hands bleeding continually from 1918; however, the lesions faded at the time of his death.^[17,19]

As shown by the above-mentioned data, stigmata have been manifesting for centuries in the context of Catholic Europe. In the USA, similar episodes were unheard of until the spreading of Catholicism.^[27] In the XX century, stigmata were reported among other religious contexts (e.g., Muslim ascetics developed lesions in the same sites of injuries endured by the Prophet in his battles).^[27] Cases in America and Australia were also reported.^[15] Kluger and Cribier in 2013 compiled a list of the most praised stigmatized. However, few people with stigmata have been beatified or canonized, and the Catholic Church does not allow for mandatory association between stigmata and sanctity, declining the conventional belief that stigmata represent a supernatural event, and are granted by God as a sign of piety. A proclaim by Pope Urban VIII instructs Catholic writers to attribute a purely human nature to this phenomenon.^[28]

The question of the etiological interpretation of stigmata remains controversial; as a matter of fact, there is no reason to presume that all stigmata should have the same etiology. Excluding ascertained cases of voluntary self-harm for malingering, as our case and few others (Table 2), stigmatization is generally referred to as unconscious self-harm during hysteria, autosuggestion, and hypnotism; these are all characteristic factors that can be associated with "mystical delirium."^[15,17,19,29]

According to various authors, however, cases of "inexplicable" stigmata with spontaneous onset and absence of external triggering factors have been described.^[15,17,19] Discrete exceptional clinical entities, not fully understood in their pathogenesis, could possibly constitute the underlying condition to some of these cases, for example, autoerythrocyte sensitization syndrome (Gardner–Diamond syndrome, or psychogenic purpura), vicarious menstruation (various organs possibly affected), hematidrosis (blood cells in sweat), and hemolacria (secondary to ophthalmic conditions).^[30-38] Such conditions usually affect young women, although often presenting variable features of psychosomatic personality, and are certainly considered "natural."

Table 2**Cases of religious stigmata due to self-harm for malingering purposes.**

Case	Age, y	Sex	Type of lesions and sites	Context	Comment
Ref. ^[16]	42	M	Palmar ulcers	For 3 y consecutively at Easter time	He stated to be blessed by God with Christ's stigmata. His partner revealed that lesions were self-inflicted with a knife
Ref. ^[17]	23	M	Erosion on dorsal surface of hands and right palm. Bullous lesions ("PAX") on left palm	Single episode	Hysterical personality, manipulated by relatives for profit. Intervention of law authority stopped the lucrative affair
Ref. ^[18]	10	F	Religious stigmata	3 wk before Easter	Religious, no psychopathologies, except indifference toward bleeding and auditory hallucinations
Our case	42	F	Erosions on forehead. Ulcers on dorsal surface of hands and feet	For 3 y consecutively at Easter time	Self-induced stigmata for profit. Intervention of law authority stopped lucrative affair

Given all the earlier discussion, religious stigmata are progressively shifting from medieval mysteries to distinct psychiatric and psychosomatic disorders. In studying stigmata, however, we should always consider the possibility that they could be self-inflicted for illegal and/or profit purposes.

Women of catholic faith are mainly affected. In the context of a state of trance, bleeding lesions appear, overlapping those suffered by Jesus Christ, on hands and feet (not on wrists, as evidenced by the Shroud of Turin in relation to the spikes), sometimes on the ribs (in relation to the thrust of the spear), or on the back (site of flagellation), and less often on the forehead (in relation to the crown of thorns) or on the shoulders (in relation to the pressure areas of the cross).

In conclusion, regardless of personal beliefs, we should always consider the possibility that stigmata, whose study "does typify the fascinating relation between physical, psychological and spiritual phenomena," could be self-inflicted for illegal and/or profit purposes.

References

- Gieler U, Consoli SG, Tomas-Aragones L, et al. Self-inflicted lesions in dermatology: terminology and classification. A position paper from the European Society for Dermatology and Psychiatry (ESDaP). *Acta Derm Venereol* 2013;93:4–12.
- Lyell A. Cutaneous artifactual disease. A review, amplified by personal experience. *J Am Acad Dermatol* 1979;1:391–407.
- Fabisch W. What is dermatitis artefacta? *Int J Dermatol* 1981;20:427–8.
- Koblentz CS. Psychosomatic concepts in dermatology. A dermatologist-psychoanalyst's viewpoint. *Arch Dermatol* 1983;119:501–12.
- Consoli SG. Dermatitis artefacta: a general review. *Eur J Dermatol* 1995;5:5–11.
- Angelini G, Bonamonte D. La dermatite artefatta. *G Ital Derm Venereol* 1999;134:99–113.
- Koblentz CS. Dermatitis artefacta. Clinical features and approaches to treatment. *Am J Clin Dermatol* 2000;1:47–55.
- Gieler U, Eckhardt J, Henn A. Factitious disorders. *Dermatol Psychosom* 2004;5:93–8.
- Nielsen K, Jeppesen M, Simmelsgaard L, et al. Self-inflicted skin diseases. A retrospective analysis of 57 patients with dermatitis artefacta seen in a dermatology department. *Acta Derm Venereol* 2005;85:512–5.
- Meneghini CL, Angelini G. Occupational dermatitis artefacta. *Derm Beruf Umwelt* 1979;27:163–5.
- Lachapelle JM. Dermatoses simulées en médecine du travail. Masson, Dermatologie Professionnelle et de l'Environnement. Paris:1992.
- Angelini G, Bonamonte D, Dyal-Smith D, Marks R. Occupational dermatitis artefacta. *Dermatology at the Millennium. The Proceedings of the 19th World Congress of Dermatology, Sydney, June 15–20, 1997.* New York:The Parthenon Publishing Group; 1999. 508–13.
- Angelini G, Kanerva L, Elsner P, Wahlberg JE, et al. Occupational dermatitis artefacta. *Handbook of Occupational Dermatology.* Berlin: Springer Verlag; 2000. 141–7.
- Kluger N. Devotional dermatoses. *Ann Dermatol Venereol* 2012;139:309–20.
- Kluger N, Cribier B. Stigmata: from Saint-Francis of Assisi to idiopathic haematidrosis. *Ann Dermatol Venereol* 2013;140:771–7.
- Hissa-Elián A, Valera CF, Pires GJT, et al. Dermatitis artefacta—stigmata Christ as a self-inflicted disorder. P4937. In: 21st World Congress of Dermatology, Buenos Aires, Argentina, 2007.
- Randazzo SD. Quando la pelle è testimone d'accusa. *Med Psicossom* 1995;4:299–308.
- Early LF, Lifschutz JE. A case of stigmata. *Arch Gen Psychiatry* 1974;30:197–200.
- Simpson CJ. The stigmata: pathology or miracle? *Br Med J* 1984;289:1746–8.
- Mucci C. Il fenomeno delle stimmate. *La Civiltà Cattolica* 1993;144:217–26.
- The Bible. Galatians 6:17.
- Thurston H. *The Physical Phenomena of Mysticism.* London:Burns and Oates; 1952.
- Vanchez A. Les stigmates de Saint François et leur détracteurs dans les derniers siècles du moyen âge. *Melanges Archeol Histoire* 1968;80:595–625.
- Imbert-Gourbeyre A. *La stigmatisation.* Clermont:Bellet; 1894.
- Bion R. *The Riddle of the Stigmata.* London:Burns and Oates; 1962.
- Schimberg AP. *The Story of Therese Neumann.* New York:All Saints Press; 1962.
- Harrison T. *Stigmata: A Medieval Mystery in a Modern Age.* New York: St. Martin's Press; 1994.
- Tocquet R. *L'ombra svelata.* Torino: Dellavalle; 1971.
- Maraldi V. *La scienza moderna e i miracoli.* Torino: SAIE; 1955.
- Boussault P, Doutre MS, Beylot-Barry M, et al. Painful bruising syndrome: a psychogenic disease. *Rev Med Interne* 2005;26:744–7.
- Ivanov OL, Lvov AN, Michenko AV, et al. Autoerythrocyte sensitization syndrome (Gardner–Diamond syndrome): review of the literature. *J Eur Acad Dermatol Venereol* 2009;23:499–504.
- Gauba V, Cooper M, Liu C. Vicarious menstruation in primary localized conjunctival amyloidosis. *Arch Ophthalmol* 2006;124:1361–2.
- Carvalho AC, Machado-Pinto J, Nogueira GC, et al. Hematidrosis: a case report and review of the literature. *Int J Dermatol* 2008;47:1058–9.
- Manonukul J, Wisuthsarewong W, Chantorn R, et al. Hematidrosis: a pathologic process or stigmata. A case report with comprehensive histopathologic and immunoperoxidase studies. *Am J Dermatopathol* 2008;30:135–9.
- Wang Z, Yu Z, Su J, et al. A case of hematidrosis successfully treated with propranolol. *Am J Clin Dermatol* 2010;11:440–3.
- Praveen BK, Vincent J. Hematidrosis and hemolacria: a case report. *Indian J Pediatr* 2012;79:109–11.
- Latorre Martinez N, Betloch Mas I, Monteagudo Paz AF, et al. Recurrent bleeding in an 18-year-old girl. *Arch Dermatol* 2012;148:960–1.
- Ho JZ, de Silva J, Olver J. A very rare case of bloody tears with enigmatic epistaxis and haematuria. *Orbit* 2011;30:116–7.