

World Journal of *Gastroenterology*

World J Gastroenterol 2019 May 28; 25(20): 2402-2538



OPINION REVIEW

- 2402 From 2-dimensional to 3-dimensional: Overcoming dilemmas in intestinal mucosal interpretation
Charlesworth RP, Marsh MN

REVIEW

- 2416 Optimizing radiotherapy with immune checkpoint blockade in hepatocellular carcinoma
Choi C, Yoo GS, Cho WK, Park HC

MINIREVIEWS

- 2430 Contribution of pancreatic enzyme replacement therapy to survival and quality of life in patients with pancreatic exocrine insufficiency
Layer P, Kashirskaya N, Gubergits N
- 2442 Hepatocellular adenoma: An unsolved diagnostic enigma
Renzulli M, Clemente A, Tovoli F, Cappabianca S, Bolondi L, Golfieri R

ORIGINAL ARTICLE**Basic Study**

- 2450 Trimethylamine N-oxide attenuates high-fat high-cholesterol diet-induced steatohepatitis by reducing hepatic cholesterol overload in rats
Zhao ZH, Xin FZ, Zhou D, Xue YQ, Liu XL, Yang RX, Pan Q, Fan JG
- 2463 *Brucea javanica* oil emulsion improves the effect of radiotherapy on esophageal cancer cells by inhibiting cyclin D1-CDK4/6 axis
Qiu ZH, Zhang WW, Zhang HH, Jiao GH

Case Control Study

- 2473 Interleukin-22 receptor 1 is expressed in multinucleated giant cells: A study on intestinal tuberculosis and Crohn's disease
Yu ZQ, Wang WF, Dai YC, Chen XC, Chen JY

Retrospective Study

- 2489 Prognostic significance of lymphovascular invasion in colorectal cancer and its association with genomic alterations
Jiang HH, Zhang ZY, Wang XY, Tang X, Liu HL, Wang AL, Li HG, Tang EJ, Lin MB

Observational Study

- 2503** Rational arrangement of measuring shear wave speed in the liver
Yokoo T, Kanefuji T, Suda T, Nagayama I, Hoshi T, Abe S, Morita S, Kamimura H, Kamimura K, Tsuchiya A, Takamura M, Yagi K, Terai S
- 2514** Effect of Blumgart anastomosis in reducing the incidence rate of pancreatic fistula after pancreatoduodenectomy
Li YT, Zhang HY, Xing C, Ding C, Wu WM, Liao Q, Zhang TP, Zhao YP, Dai MH

SYSTEMATIC REVIEWS

- 2524** Systematic review: Renin-angiotensin system inhibitors in chemoprevention of hepatocellular carcinoma
Barone M, Viggiani MT, Losurdo G, Principi M, Leo AD

ABOUT COVER

Editorial board member of *World Journal of Gastroenterology*, Kentaro Yoshioka, MD, PhD, Professor, Department of Liver, Biliary Tract and Pancreas Diseases, Fujita Health University, Toyoake 470-1192, Japan

AIMS AND SCOPE

World Journal of Gastroenterology (*World J Gastroenterol*, *WJG*, print ISSN 1007-9327, online ISSN 2219-2840, DOI: 10.3748) is a peer-reviewed open access journal. The *WJG* Editorial Board consists of 642 experts in gastroenterology and hepatology from 59 countries.

The primary task of *WJG* is to rapidly publish high-quality original articles, reviews, and commentaries in the fields of gastroenterology, hepatology, gastrointestinal endoscopy, gastrointestinal surgery, hepatobiliary surgery, gastrointestinal oncology, gastrointestinal radiation oncology, etc. The *WJG* is dedicated to become an influential and prestigious journal in gastroenterology and hepatology, to promote the development of above disciplines, and to improve the diagnostic and therapeutic skill and expertise of clinicians.

INDEXING/ABSTRACTING

The *WJG* is now indexed in Current Contents®/Clinical Medicine, Science Citation Index Expanded (also known as SciSearch®), Journal Citation Reports®, Index Medicus, MEDLINE, PubMed, PubMed Central, Scopus and Directory of Open Access Journals. The 2018 edition of Journal Citation Report® cites the 2017 impact factor for *WJG* as 3.300 (5-year impact factor: 3.387), ranking *WJG* as 35th among 80 journals in gastroenterology and hepatology (quartile in category Q2).

RESPONSIBLE EDITORS FOR THIS ISSUE

Responsible Electronic Editor: *Yan-Liang Zhang* Proofing Editorial Office Director: *Ze-Mao Gong*

NAME OF JOURNAL

World Journal of Gastroenterology

ISSN

ISSN 1007-9327 (print) ISSN 2219-2840 (online)

LAUNCH DATE

October 1, 1995

FREQUENCY

Weekly

EDITORS-IN-CHIEF

Subrata Ghosh, Andrzej S Tarnawski

EDITORIAL BOARD MEMBERS

<http://www.wjgnet.com/1007-9327/editorialboard.htm>

EDITORIAL OFFICE

Ze-Mao Gong, Director

PUBLICATION DATE

May 28, 2019

COPYRIGHT

© 2019 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>

Systematic review: Renin-angiotensin system inhibitors in chemoprevention of hepatocellular carcinoma

Michele Barone, Maria Teresa Viggiani, Giuseppe Losurdo, Mariabeatrice Principi, Alfredo Di Leo

ORCID number: Michele Barone (0000-0001-8284-5127); Maria Teresa Viggiani (0000-0002-8689-9103); Giuseppe Losurdo (0000-0001-7038-3287); Mariabeatrice Principi (0000-0003-0545-5656); Alfredo Di Leo (0000-0003-2026-1200).

Author contributions: Barone M and Di Leo A conceived and planned the study. Viggiani MT, Barone M, Losurdo G and Principi M collected the data. Barone M and Losurdo G wrote the manuscript. Viggiani MT and Principi M revised the draft. All authors read and approved the final version of the manuscript.

Conflict-of-interest statement: All authors declare no conflict of interest (e.g., employment, consultancies, honoraria, stock ownership or options, expert testimony, grants/patents received, or royalties).

PRISMA 2009 Checklist statement: The manuscript was prepared according to the PRISMA 2009 Checklist.

Open-Access: This is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Michele Barone, Maria Teresa Viggiani, Giuseppe Losurdo, Mariabeatrice Principi, Alfredo Di Leo, Gastroenterology Section, Department of Emergency and Organ Transplantation, University of Bari Aldo Moro, Bari 70124, Italy

Corresponding author: Michele Barone, MD, PhD, Adjunct Professor, Gastroenterology Section, Department of Emergency and Organ Transplantation, University of Bari Aldo Moro, Piazza Giulio Cesare 11, Bari 70124, Italy. michele.barone@uniba.it

Telephone: +39-80-5593514

Fax: +39-80-5593177

Abstract

BACKGROUND

Neoangiogenesis is one of the key pathogenetic mechanisms in hepatocellular carcinoma (HCC). Modulation of the renin-angiotensin system (RAS) by angiotensin-converting enzyme inhibitors (ACE-Is) and angiotensin receptor blockers (ARBs) seems to be a possible adjuvant therapy for HCC, due to the anti-angiogenic and anti-fibrogenic activity of these drugs.

AIM

To elucidate the role of ARBs and ACE-Is in HCC.

METHODS

We performed an electronic search of the literature using the most accessed online databases (PubMed, Cochrane library, Scopus and Web of Science), entering the query terms "angiotensin-converting enzyme inhibitors" OR "ACE inhibitors" OR "ACE-I" AND "hepatocarcinoma*" OR "hepatocellular carcinoma"; moreover "angiotensin II type 1 receptor blockers" OR "ARBs" AND "hepatocarcinoma*" OR "hepatocellular carcinoma". Eligibility criteria were: (1) prospective or retrospective clinical studies; (2) epidemiological studies; and (3) experimental studies conducted *in vivo* or *in vitro*. Abstracts, conference papers, and reviews were excluded a priori. We limited our literature search to articles published in English, in peer-reviewed journals.

RESULTS

Thirty-one studies were selected. Three interventional studies showed that ACE-Is had a significant protective effect on HCC recurrence only when used in combination with vitamin K or branched chain aminoacids, without a significant increase in overall survival. Of six retrospective observational studies, mainly focused on overall survival, only one demonstrated a prolonged survival in the ACE-Is group, whereas the two that also evaluated tumor recurrence showed

ses/by-nc/4.0/

Manuscript source: Unsolicited manuscript**Received:** January 23, 2019**Peer-review started:** January 23, 2019**First decision:** February 21, 2019**Revised:** March 6, 2019**Accepted:** March 24, 2019**Article in press:** March 25, 2019**Published online:** May 28, 2019**P-Reviewer:** Ikura Y, Ocker M, Streba LAM**S-Editor:** Ma RY**L-Editor:** A**E-Editor:** Zhang YL

conflicting results. All experimental studies displayed beneficial effects of RAS inhibitors on hepatocarcinogenesis. Numerous experimental studies, conducted either on animals and cell cultures, demonstrated the anti-angiogenetic and antifibrotic effect of ACE-Is and ARBs, thanks to the suppression of some cytokines such as vascular endothelial growth factor, hypoxia-inducible factor-1 α , transforming growth factor-beta and tumor necrosis factor alpha. All or parts of these mechanisms were demonstrated in rodents developing fewer HCC and preneoplastic lesions after receiving such drugs.

CONCLUSION

In humans, RAS inhibitors - alone or in combination - significantly suppressed the cumulative HCC recurrence, without prolonging patient survival, but some limitations intrinsic to these studies prompt further investigations.

Key words: Cirrhosis; Renin; Angiotensin; Survival; Cancer prevention; Hepatocarcinoma

©The Author(s) 2019. Published by Baishideng Publishing Group Inc. All rights reserved.

Core tip: All experimental studies strongly support the ability of renin-angiotensin system (RAS) inhibitors to counteract carcinogenesis essentially by inhibiting angiogenesis and possibly fibrogenesis in chemically-induced or xeno-transplanted hepatocarcinomas, whereas additional pathogenetic mechanisms using animal models that mimic the progression of liver disease in humans and have been barely considered. In all randomized clinical trials, RAS inhibitors - in combination with other therapies - significantly reduced cumulative hepatocarcinoma recurrence. However, the data on patient survival obtained by these studies and several observational studies are not conclusive. The beneficial effect of the early use of RAS inhibitors on the progression of cirrhosis toward hepatocarcinoma cannot be excluded.

Citation: Barone M, Viggiani MT, Losurdo G, Principi M, Leo AD. Systematic review: Renin-angiotensin system inhibitors in chemoprevention of hepatocellular carcinoma. *World J Gastroenterol* 2019; 25(20): 2524-2538

URL: <https://www.wjgnet.com/1007-9327/full/v25/i20/2524.htm>

DOI: <https://dx.doi.org/10.3748/wjg.v25.i20.2524>

INTRODUCTION

Liver cancer is the fifth most common cancer and the second most frequent cause of cancer-related death globally^[1]. In particular, it is much more common in men than in women, and if we refer solely to more developed countries, it is the sixth leading cause of cancer death in men^[2].

Hepatocellular carcinoma (HCC) represents about 90% of primary liver cancers and usually develops in patients with underlying liver cirrhosis^[1]. Etiologic factors are most frequently represented by chronic viral hepatitis [hepatitis B virus (HBV) and hepatitis C virus (HCV)], alcohol intake, and non-alcoholic steatohepatitis (NASH), which all together are responsible for about 80% of HCC^[3].

One of the major cancer hallmarks of HCC is the process of neoangiogenesis, which is thought to occur early in the carcinogenic process. Even if the pathogenetic mechanisms responsible for the neoangiogenesis remain to be clarified, it is plausible that the intervention of proangiogenic growth factors occurs as a result of signals mediated by hypoxia-inducible transcription factors^[4].

As a result of the involvement of angiogenesis, clinical studies have been focused on anti-angiogenic biological agents in HCC treatment^[5-7]. However, this therapeutic approach to cirrhotic patients has been frequently associated with severe side effects, and it is burdened by high costs and contraindications to their long-term administration^[8]. Inexpensive and easily manageable drugs could represent an alternative therapy with a proven safety profile on long-term administration. From this point of view, angiotensin-converting enzyme inhibitors (ACE-Is) and angiotensin II type 1 receptor blockers (ARBs) seem to be a possible candidate for adjuvant therapies for HCC, as a consequence of their well-known anti-angiogenic and anti-fibrogenic activity^[9-11]. Theoretically, ACE-I and ARBs could also have a role in HCC primary

prevention since they reduce both neovascularization and the development of liver fibrosis^[10].

We decided to perform this study as the possible role of renin-angiotensin system (RAS) inhibitors in the prevention of HCC is still an attractive field of research, as suggested by the recently published experimental and clinical studies. In order to better clarify this issue, in this systematic review, we performed a search of studies in this field. Using this method, we selected 31 studies, mainly focused on experimental data.

MATERIALS AND METHODS

Data sources and searches

Literature was selected in August 2018, focusing on studies regarding ACE-Is, ARBs and HCCs. For this purpose, an electronic search of the literature was conducted using the online databases PubMed, Cochrane library, Scopus and Web of Science, entering the query terms "angiotensin-converting enzyme inhibitors" OR "ACE inhibitors" OR "ACE-I" AND "hepatocarcinoma*" OR "hepatocellular carcinoma"; moreover "angiotensin II type 1 receptor blockers" OR "ARBs" AND "hepatocarcinoma*" OR "hepatocellular carcinoma" were used. Further relevant articles were hand-searched using the references of the selected studies.

Study selection

Eligibility criteria were: (1) prospective or retrospective clinical studies; (2) epidemiological studies; and (3) experimental studies conducted *in vivo* or *in vitro*. Abstracts, conference papers, and reviews were excluded a priori. We limited our literature search to articles published in English, in peer-reviewed journals.

Data extraction and quality assessment

We conducted the articles selection using a two-step approach, to evaluate whether they matched the eligibility criteria. The first phase was based on titles and abstracts, and the second on full-text examination. This procedure was performed by two of the authors (Viggiani MT and Principi M), and the percentage of agreement demonstrated high inter-rater reliability (97.4%). Lastly, discrepancies in data extraction were resolved by introducing a third, experienced arbitrator (Losurdo G).

In cases of studies analyzing overlapping periods from the same registry/database, we considered only the study that examined the more extended period and the more significant number of patients.

RESULTS

As shown in **Figure 1**, the combined search of electronic databases identified 231 articles. Three additional articles were hand-searched using references of the selected studies. After removing duplicates, 86 articles were screened. On the basis of the abstract evaluation, 33 articles were excluded since they did not match the eligibility criteria. The evaluation of full-text articles led to exclude 15 additional manuscripts. We found a meta-analysis on the beneficial effects of ACE-I/ARBs against several tumors^[12]; this study was excluded since it took in consideration only a trial on HCC that we had already included as a single study^[13]. Finally, ten reviews were excluded since they evaluated the same studies on the relationship between ACE-Is or ARBs already selected. At the end of our selection, only 31 studies were included in the present review.

Clinical studies

Table 1 reports the characteristics of all clinical studies selected. We found only three interventional studies, all performed by the same authors, which examined the influence of a putative antitumoral treatment after randomization of the patients^[13-15]. These studies evaluated the influence of ACE-Is, given alone or in combination with either vitamin K or branched chain aminoacids (BCAA), on HCC recurrence, showing a significant protective effect of ACE-I only when used in combination with vitamin K or BCAA. Patients were enrolled when they had a low number and low size tumors susceptible to radiofrequency ablation (RFA) treatment and were excluded in the event of early recurrence (within 3 mo). No significant increase in overall survival was observed in any of the three studies.

Six more recent retrospective observational studies evaluated the effect of ACE-Is or ARBs on overall survival (OS) and/or time to recurrence in patients who




Figure 1 PRISMA flow diagram describing the process of study selection. HCC: Hepatocellular carcinoma.

underwent different HCC treatments (Table 1). Ho *et al*^[16] evaluated the effect of ACE-I or ARB exposure within the first six months after initiating antiviral agents in a high-risk HBV and HCV large cohorts of patients and did not find any protective effect of these drugs on HCC development adjusted for potential confounders. Surprisingly, in HCV patients without cirrhosis, diabetes, and hyperlipidemia, they found an increased HCC risk associated with ACE-I or ARB use in (HR = 4.53, 95%CI: 1.46-14.1). Hagberg *et al*^[17] restricted the study population to patients with hypertension (both cases and controls). Also, in this case, no protective effect was observed with ACE-I or ARB use. However, patients were already defined exposed (cases) after receiving at least two prescriptions for ACE-I and/or ARBs before the index date. Walker *et al*^[18], included subjects that were exposed to ACE-I treatment 12-24 mo before the first recorded liver cancer diagnosis without finding any protective effect.

Differently from what observed by the previous authors, Pinter *et al*^[19] found a prolonged overall survival in HCC patients treated with sorafenib/supportive therapy and already receiving ACE-I or ARBs for a median duration of treatment of 12.7 mo. Also, Facciorusso *et al*^[20] reported a prolonged overall survival in HCC patients treated with RFA and already receiving ARBs (but not ACE-I) for at least the previous two years. In this study, a significantly longer disease-free survival in patients receiving ARBs (HR = 0.47; 95%CI: 0.27-0.82, $P = 0.009$)^[20] was reported. In the last clinical study, by Karibori *et al*^[21] the treatment of hypertension with ACE-Is or ARBs did not improve overall survival after tumor resection in HCV-HCC patients with or without cirrhosis compared to the related controls. Interestingly, HCC-HCV cirrhotic patients with hypertension treated with other antihypertensive drugs had a significantly reduced overall survival compared to HCC-HCV cirrhotic patients receiving either ACE-Is or ARBs or no antihypertensive treatment ($P = 0.029$). Finally, hypertension had no influence on the prognosis of inpatients with HCC related to other etiologies^[21].

An additional study, not listed in Table 1, analyzing the association between the use of statins in HBV-infected patients and the risk of HCC, reported that the use of ACE-I for more than one year did not significantly influence the protective effect of statins against HCC development^[22]. However, in this study, a large number of patients in the statin group (62%) was diabetic and therefore probably using

Table 1 Clinical studies on the effect of angiotensin-converting enzyme inhibitors or angiotensin II type 1 receptor blockers exposure in patients affected or at high risk of hepatocellular carcinoma

Interventional studies				
Authors	Patients	Treatment	HCC recurrence after 42-54 mo	P value
Yoshiji <i>et al</i> ^[13]	87 with RFA for prior HCC	I. Control	18/25	I vs II; < 0.01
		II. ACE-I + vit. K2	9/25	
		III. ACE-I	19/19	I vs III, n.s.
		IV. vit. K2	18/18	I vs IV; n.s.
Yoshiji <i>et al</i> ^[14]	89 with Insulin resistance and RFA for prior HCC	I. Control	16/26	I vs II; < 0.01
		II. ACE-I + BCAA	9/28	I vs III, n.s.
		III. ACE-I	11/19	
		IV. BCAA	9/16	I vs IV; n.s.
Yoshiji <i>et al</i> ^[15]	54 with HCC randomized in 2 groups, before treatment	I. HCC treatment	77%	I vs II < 0.01
		II. HCC treatment + ACE-I and Vit. K	40%	
Observational studies				
Authors	Patients	Treatment	OS, HR, OR	P value
Ho <i>et al</i> ^[16]	7724 HBV-patients	ACE-I or ARB (46.3% in HBV and 42.5% in HCV) within 6 mo after initiating DAAs	HCC risk after ACE-I and ARB exposure	n.s. ¹
	7873 HCV- patients at high-risk of HCC development		HR = 0.97, 95%CI: 0.81-1.16 HR = 0.96, 95%CI: 0.80-1.16 respectively	
Hagberg <i>et al</i> ^[17]	490 HCC 1909 controls	ACE-I or ARBs	OR = 1.14, 95%CI: 0.85-1.55 users vs non- users	n.s. ²
Walker <i>et al</i> ^[18]	224 HCC	7% ACE-I users in HCC group	OR = 1.29, 95%CI: 0.88-1.88	n.s. ³
	2313 controls	5.9% ACE-I users in control group	unexposed vs exposed	
Pinter <i>et al</i> ^[19]	156, with Sorafenib or supportive therapy	ACE-I or ARBs in 43 pts.	OS = 11.9 mo vs 6.8 mo	P = 0.014
	76, (confirmation cohort) with sorafenib or supportive therapy	ACE-I or ARBs in 38 pts.	HR= 0.6, 95%CI: 0.4-0.9 OS = 19.5 mo vs 10.9 mo HR= 0.5, 95%CI: 0.3-1.0	P = 0.011 P = 0.043 P = 0.038
Facciorusso <i>et al</i> ^[20]	153 with RFA for prior HCC	I: Control, 73 pts	OS = 48 mo	I vs II, n.s.
		II: ACE-I, 49 pts	OS = 72 mo	I < III, P < 0.002
		III. ARBs, 31 pts	OS = 84 mo HR ⁴ = 0.39, 95%CI: 0.22-0.66	
Kabori <i>et al</i> ^[21]	185 HCV-HCC pts. without cirrhosis	I. No hypert.	OS at 5 yr (76/106)	I vs II, n.s.
		II. Hypert. + ACE/ARB	OS at 5 yr (30/37)	I > III, P < 0.001
		III. Hypert. + other anti-hypertensives	OS at 5 yr (11/42)	II > III, P < 0.001
	141 HCV-HCC pts. with cirrhosis	I. No hypertension	OS at 5 yr 51.6%	I and II > III, P = 0.029
		II. Hypert. + ACE/ARB	OS at 5 yr 76.7%	
		III. Hypert. + other	OS at 5 yr 37.3%	
143 pts. with HCC related to other etiologies	I. No hypertension	OS at 5 yr 59%-74%	n.s.	
	II. Hypertension	OS at 5 yr 60%-62%		

¹Adjusted for sex, age, liver cirrhosis, diabetes mellitus, alcohol consumption, hyperlipidemia, malignancies other than hepatocellular carcinoma (HCC), chronic obstructive pulmonary disease, end-stage renal disease, transplantation, aspirin, metformin, and statins.

²Adjusted odds ratio for HCC risk factors (body mass index, smoking status, alcohol abuse, hepatitis B virus and/or hepatitis C virus, rare metabolic disorders, liver disease, diabetes, and use of statins and acetaminophen) and duration of hypertension. The analysis was also repeated to cases and controls without diabetes and to cases and controls without liver disease demonstrating no significant difference.

³Adjusted for alcohol use, smoking, cirrhosis, and hepatitis.

⁴Adjusted for age, gender, Child-Pugh class and α -fetoprotein. Time to recurrence was significantly reduced in patients receiving angiotensin II type 1 receptor blockers ($P = 0.009$). RFA: Radiofrequency ablation; OS: Overall survival; OR: Odds ratio; HR: Hazard ratio; ACE-Is: Angiotensin-converting enzyme inhibitors; ARBs: Angiotensin II type 1 receptor blockers; Vit.: Vitamine; CI: Confidence interval; HCC: Hepatocellular carcinoma; HBV: Hepatitis B virus; HCV: Hepatitis C virus; Pts.: Patients.

metformin or thiazolidinediones, which have been demonstrated to profoundly reduce HCC risk^[23], and therefore could have also overcome the effect of ACE-I/ARBs.

To our knowledge, the only case report that matched our selection criteria describes the protective effect of an ACE-I used in combination with vitamin K2 in a patient with a dysplastic nodule on liver cirrhosis. After one year of treatment, the authors describe a reduction of α -fetoprotein (AFP) and AFP-L3 and the disappearance of the dysplastic nodule^[24].

Experimental data

The most consistent data on the potential role of ACE-Is and ARBs in HCC prevention/cure come from experimental *in vitro* and *in vivo* studies.

Diethylnitrosamine model

Table 2 summarizes all studies performed either in mice or rats using the diethylnitrosamine (DEN) model. Saber *et al*^[25] showed that sorafenib or fosinopril, perindopril and losartan, prolonged overall survival of mice. Both ACE-Is and losartan brought about an improvement in the histological situation, similar to that determined by sorafenib, with regression of malignant histological features. However, only sorafenib and perindopril significantly reduced α -fetoprotein levels as compared to controls. Additionally, sorafenib and anti-hypertensive drugs significantly decreased the transforming growth factor (TGF)- β , tumor necrosis factor (TNF)- α , matrix metalloproteinase (MMP)-2 and vascular endothelial growth factor (VEGF) serum levels compared to controls; however, the administration of perindopril was even more effective than sorafenib in the reduction TNF- α level. The authors conclude their study suggesting that the reduction of inflammatory cytokines, in particular, TNF- α , could have been mediated by the inhibition of phosphorylation-induced NF κ B degradation and phosphorylation of NF κ Bp65. Therefore, the RAS inhibition may have a dual anticancer effect mediated by the suppression of inflammation and angiogenesis. The same group repeated the same experiment^[26] in mice and confirmed the data about the regression of cellular atypia in HCC, but they did not observe a positive effect on animal survival using ACE-Is or ARBs alone, or in combination with sorafenib. In another study^[27], albino DEN-treated mice received perindopril, leflunomide, curcumin alone or in combination. While all DEN-treated mice developed HCC, the prevalence of HCC decreased to 37.5% under curcumin, and 25% under leflunomide and perindopril and no mice developed liver dysplastic lesions in the combination arm. This finding paralleled with the reduction of angiogenesis, immunohistochemically evaluated by CD31, an antigen widely used as a marker of neovascularization^[28]. Finally, the authors also demonstrated a dramatic reduction of hypoxia-inducible factor (HIF)- α . Mansour *et al*^[29] tested captopril, perindopril, and losartan in male albino rats treated with DEN. Some of the rats receiving DEN plus an active drug also received carbon tetrachloride to stimulate cell proliferation. The administration of captopril and losartan induced a marked reduction of α -fetoprotein and almost halved the levels of VEGF, fibroblast growth factor (FGF) and TGF- β 1 only in rats in which hepatocarcinogenesis was promoted by DEN. On the other hand, MMP-2 activity was inhibited by all three drugs in HCC-rats while the levels of tissue inhibitor of metalloproteinase (TIMP)-1, a negative regulator of MMP2, were increased only by the treatment with losartan and perindopril. This study provided exciting results since it demonstrated a prophylactic effect on the mechanisms of hepatocarcinogenesis when ARBs and ACE-Is were given two weeks before DEN injection and not at the same time.

In another model, in addition to DEN treated standard rats, BALB-c mice underwent an injection of HCC cell line. In these mice, the administration of perindopril or 5-fluorouracil alone was not able to reduce the size of HCC^[30], while the combination of the two drugs produced much better results thanks to the increased apoptotic activity and the reduction of both the CD31-mRNA and VEGF expression in HCC tissue. Interestingly, the number of preneoplastic lesions was decreased only by the ACE-I, and the addition of 5-fluorouracil enhanced this effect. Lastly, on the basis of additional *in vitro* studies, they concluded that ACE-I and 5-fluorouracil might reduce the tumor growth by inhibiting neovascularization, which in turn may induce apoptosis of the tumor cells, rather than via direct inhibition of tumor cell proliferation.

Table 2 Studies on the effect of angiotensin-converting enzyme inhibitors or angiotensin II type 1 receptor blockers in diethylnitrosamine-induced hepatocellular carcinoma animal models

	Animal model	Treatment	Results	Comments
Saber <i>et al</i> ^[25]	DEN-induced HCC in mice	I. Sorafenib <i>vs</i> II. ACEIs or ARBs	Both treatments improved liver histology; II. reduced α -feto-protein and VEGF level	Inhibition of proliferation by involvement of NF κ B pathway and cyclin D1
Saber <i>et al</i> ^[26]	DEN-induced HCC in mice	1 Sorafenib 2 ACE-I \pm Sorafenib 3 ARB \pm Sorafenib	ACE-Is and ARBs monotherapy or plus sorafenib improved liver histology with regression to grade 1, almost restoration of lobular architecture	No survival improvement No additional effect when combination therapy was used
Nasr <i>et al</i> ^[27]	DEN-induced HCC in mice	Leflunomide Perindopril Curcumin	All drugs abrogated: hepatic microvessel density, elevated VEGF; only curcumin reduced HIF-1 α . Nodules reduced or absent	Combination of these agents: further inhibited neovascularization
Mansour <i>et al</i> ^[29]	DEN + carbon tetra-chloride in rats	ACE-I ARBs	Significant reduction of tumor markers and hepatic growth factors	Liver histology amelioration correlated with VEGF, CD31 and FGF
Yanase <i>et al</i> ^[30]	DEN-treated rats Male BALB/c mice with injections of BNL-HCC cells.	combined effect of ACE-I and 5-fluorouracil	Inhibition of HCC growth, neovascularization (VEGF and CD31+ vessels suppression), and marked increase of apoptosis	<i>In vitro</i> studies on EC tubular formation confirm the anti-angiogenetic effect
Yoshiji <i>et al</i> ^[31]	DEN-treated mice Male BALB/c mice with injections of BNL-HCC cells	Vit. K and ACE-I used singularly or in combination	Inhibitory effects by each compound on hepato-carcinogenesis, more potent when used in combination	Increased apoptosis in the tumor, w/o any effect on tumor cell proliferation; CD31 mRNA suppression
Yoshiji <i>et al</i> ^[32]	DEN-treated rats and human HCC cell lines	Vit. K or ACE-I alone or in combination	Chemopreventive effect on pre-neoplastic foci formation by single compounds, more potent when combined.	Inhibition of endothelial cell proliferation and tubular formation; reduction of CD31 mRNA expression.
Yoshiji <i>et al</i> ^[33]	DEN-treated rats	Interferon, ACE-I used singularly or in combination	IFN or PE, when used singularly, significantly attenuated, in combination nearly abolished HCC	CD31 and VEGF mRNA expression were reduced; apoptosis was also reduced; no change of cell proliferation
Yoshiji <i>et al</i> ^[34]	DEN-treated rats Endothelial cells <i>in vitro</i>	Perindopril (ACE-I)	Inhibition of neo-vascularization and VEGF expression	Suppression of VEGF-induced tubular formation; no effect on endothelial cell proliferation <i>in vitro</i>

ACE-Is: Angiotensin-converting enzyme inhibitors; ARBs: Angiotensin II type 1 receptor blockers; Vit.: Vitamine; DEN: Diethylnitrosamine; HCC: Hepatocellular carcinoma; VEGF: Vascular endothelial growth factor; HIF: Hypoxia-inducible factor.

Yoshiji *et al*^[31] analyzed the effect of perindopril, vitamin K or their combination, both in DEN model and in mice injected with BNL-HCC cell line. They showed that HCC was significantly inhibited by the treatment with vitamin K and ACE-I, and their combination produced an even more potent inhibitory effect compared to the single agents. These results were observed in both models and were associated with mRNA-CD31 suppression and promotion of apoptosis (the number of TUNEL-positive cells was significantly increased by treatment with vitamin K and ACE-I compared to the control group), while the proliferative activity was not influenced. The same group replicated this protocol^[32] both in rats and HepG2 cells with the aim of studying proliferation and angiogenesis. They found that vitamin K and ACE-I inhibited endothelial cell tubule formation and endothelial cell proliferation in a dose-dependent manner while only vitamin K inhibited tumor cell proliferation. In another similar experience^[33], BALB/c mice treated with DEN and receiving either perindopril or interferon-beta or a combination of them experienced a substantial reduction in the number of nodules/liver that was maximal (less than one nodule/liver) in the group receiving both drugs. These findings paralleled the reduction of CD31-mRNA and the increase of the apoptotic index within nodules. In another study of the same group^[34] perindopril, given at low or high doses, did not influence the production of VEGF in cultured endothelial cells but was able to suppress their migration and formation of tubules. In the same study, perindopril hampered the development of preneoplastic liver lesions in DEN-treated animals, probably induced by the suppression of CD31, which paralleled the reduction of the number of lesions.

Other models

Several studies (Table 3) adopted a xenograft model in which in BALB/c mice received an injection of HCC cell lines. Using this model, Fan *et al.*^[35] demonstrated an enhanced immunohistochemical expression of angiotensin receptor 1 (AT1R) in cancer tissue, which correlated with VEGF receptor A. The administration of candesartan reduced the mean diameter of tumors and suppressed AT1R expression and decreased microvascular density. The authors then examined the effects of angiotensin II and candesartan on the expression of AT1R protein in SMMC-7721 and HepG2 cell lines using Western blotting and found that candesartan down-regulated the expression of AT1R protein induced by angiotensin II with a dose-dependent relationship, as well as angiotensin II-induced production of VEGF-A. On the other hand, when angiotensin receptor 2 was transduced by an adenoviral vector in HCC cell lines and murine orthotopic liver graft, it inhibited cell proliferation with a significant decrease of S-phase cells and an increase of G1-phase cells, probably mediated by the alteration of expression of CDK4 and cyclinD1^[36]. Therefore, it was hypothesized that apoptosis was secondary to activation of p38 MAPK, pJNK, caspase-8 and caspase-3 and inactivation of pp42/44 MAPK (Erk1/2). In another model, using the BNL-HCC cell line transferred into BALB/c mice^[37], both perindopril and interferon-beta reduced the volume of HCC, with an even higher magnitude when administered together. Additionally, the authors showed that this effect was mediated by the promotion of apoptosis, the reduction of cell proliferation and the inhibition of neo-angiogenesis proven by the reduction of CD31 positive cells and the suppression of VEGF protein within the tumor. Yoshiji *et al.*^[38] used a retroviral vector to enhance the expression of VEGF in the BNL-HCC cell line. In this way, they found that perindopril reversed the production of VEGF and that the ability of cells to migrate was directly dependent on VEGF levels. When these manipulated cells were injected in BALB/c mice, they discovered that VEGF over-expression within HCC did not modify the angiotensin-2 expression, thus suggesting that, in this model, VEGF-mediated tumor growth was not influenced by angiotensin 2. Lastly, similarly to previous studies, it was reported that perindopril inhibited CD31 expression. The same group^[39] used the BNL-HCC allograft model in BALB/c mice to show that perindopril, temocapril, and captopril had an inhibitory effect on total HCC volume burden, although the effect of perindopril was much stronger than other ACE-Is. Finally, they showed that perindopril reduced mRNA-VEGF expression and ACE enzyme activity, suppressed microvessel density and inhibited tubule formation in endothelial cells.

The long-term administration of a choline-deficient/low-methionine diet is another way of inducing steatohepatitis, liver cirrhosis, and HCC in rodents. Using this model, Tamaki *et al.*^[40] showed that none of the male Wistar rats receiving telmisartan developed HCC, while 54.6% of the control rats developed the tumor. The latter condition was associated with an increase of HIF-1 α and protein and VEGF mRNA. Furthermore, telmisartan inhibited the up-regulation of TGF- β 1, connective tissue growth factor (CTGF), and TNF- α mRNA levels. The same diet-induced HCC model was adopted by Noguchi^[41]. In this case, male Fisher-344 rats were treated with perindopril and eplerenone, a selective aldosterone blocker. Such drugs reduced both the liver fibrosis index and TGF- β . When administered in combination, they reduced the density of hepatic stellate cells and inhibited their proliferation through the reduction of aldosterone and angiotensin 2. Finally, both eplerenone and aldosterone attenuated the CD31 positive vessels in the liver along with inhibition of pre-neoplastic lesions (less than 4 foci/cm² vs about 8 foci/cm² in the control group) and, even in this report, a suppression of VEGF was recorded. Therefore, authors concluded that the effects of eplerenone and ACE-I were not simply due to the suppression of RAS, but other pathways, notably the hampering of the fibrosis process in this steatohepatitis model, could be involved.

Since insulin resistance status promotes neovascularization and has been associated with an increased risk of HCC, Yoshiji *et al.*^[42] initiated hepatocarcinogenesis by DEN in obese diabetic Otsuka Long-Evans Tokushima Fatty (OLETF) rats undergoing partial hepatectomy and used BCAAs and ACE-I, alone or in combination, at clinically comparable low doses, to assess the antineoplastic activity of these compounds. Both BCAAs and ACE-I significantly attenuated the development of dysplastic lesions, along with the suppression of both angiogenesis and VEGF expression. Used in combination, these two agents exerted an even more potent inhibitory effect. Finally, BCAAs and ACE-I induced an inhibitory effect on tubule formation in cultured endothelial cells showing a similar efficacy^[42].

In a further three studies^[43-45], the effect of ACE-Is or ARBs was tested only on cell lines. In particular, in HLF, HLE, HuH-7 and PLC/PRF/5 lines, it was demonstrated that telmisartan, but not valsartan, losartan and irbesartan, was able to arrest cell proliferation^[43]. Similarly, only telmisartan blocked the cell cycle in phase G1 and

Table 3 Experimental studies on the effect of angiotensin-converting enzyme inhibitors or angiotensin II type 1 receptor blockers in different hepatocellular carcinoma animal models and in hepatocellular carcinoma cell lines

Model	Treatment	Results	Comments
Fan <i>et al</i> ^[35] 1 Tumor cell lines 2 BALB/c nude mice with injections of HCC cell lines 3 Human HCC specimens	Candesartan	Angiotensin II up-regulated AT1R and promoted production of VEGF <i>in vitro</i> . Candesartan reversed this process and downregulated the expression of VEGF-A in tumor xenografts	AT1R expression was associated to angiogenic potential in HCC human tissues
Du <i>et al</i> ^[36] 1 Tumor cell lines 2 BALB/c nude mice with intra-liver injections of human HCC cells	AT2R over- expression by AT2R recombinant adenoviral vector	Overexpression of AT2R in transduced HCC cell lines produced apoptosis and inhibited cell proliferation. Higher AT2R expression could increase the growth of HCC and the proliferation of HCC cells <i>in vivo</i>	A moderate expression of AT2R could increase the growth of HCC and the proliferation of HCC cells <i>in vivo</i> . AT2R mechanisms remain to be elucidated
Noguchi <i>et al</i> ^[37] BALB/c mice injected with HCC cell lines	ACE-I and interferon-beta	Combination therapy was effective even on established tumors. Suppression of VEGF and endothelial cell proliferation and tubular formation, increase of apoptosis;	No effect on HCC cell proliferation.
Yoshiji <i>et al</i> ^[38] BALB/c mice injected with HCC cell lines. Endothelial cell cultures	Retroviral tetracycline up-regulated VEGF gene expression and Perindopril (ACE-I)	Perindopril significantly attenuated VEGF-mediated tumor growth and neovascularization. <i>In vitro</i> , VEGF-induced endothelial cell migration inhibition	Suppression of VEGF
Yoshiji <i>et al</i> ^[39] BALB/c mice injected with HCC cell lines. BNL CL2 cell line	Captopril Perindopril Temocapril Losartan Candesartan	Reduction of tubular formation and microvessel density in the tumor. Higher VEGF mRNA expression reduction and HCC growth inhibition by perindopril compared to temocapril and captopril. Neither candesartan nor losartan significantly inhibited the tumor development	ACE-Is suppressed the tumor development mainly by a mechanism which was independent from AT1-R blockage.
Tamaki <i>et al</i> ^[40] Male Wistar rats receiving modified choline-deficient low-methionine diet	Telmisartan	No HCC in telmisartan group <i>vs</i> 54.6% of HCC in control group. Telmisartan inhibited the dietary-induced up-regulation of VEGF, TGF- β 1, CTGF, HIF1 α and TNF- α mRNA levels	Telmisartan may prevent hepatocarcinogenesis through the inhibition of hepatic angiogenesis
Noguchi <i>et al</i> ^[41] Male Fisher-344 rats receiving modified choline-deficient low-methionine diet	Perindopril Eplerenone (selective aldosterone blocker) Aldosterone	Significant inhibitory effects on the GST-P positive pre-neoplastic lesion development; reduction of VEGF expression tubule formation and neovascularization	Combination of Perindopril + Eplerenone exerted a stronger suppression than single treatment
Yoshiji <i>et al</i> ^[42] Male OLETF rats (spontaneous development of insulin resistance) treated with Diethylnitrosamine	branched-chain amino acids perindopril	Number/size of preneoplastic foci were significantly suppressed by treatment of both drugs suppression of CD31- and VEGF-mRNA	<i>In vitro</i> suppression of tubule formation, regardless of drug concentration
Oura <i>et al</i> ^[43] Five human HCC cell lines (HepG2, HLF, HLE, HuH-7 and PLC/PRF/5)	Telmisartan Valsartan Irbesartan Losartan	Only telmisartan reduced proliferation in most of cell lines. Promotion of apoptosis, Enhancement of bFGF reduction of p-ErbB3	Arrest of cell cycle in G1, and S phase. Effect is dose-dependent and may be mediated by induction of MPK/mTOR signaling.
Santhekadur <i>et al</i> ^[44] Hep3B and QGY-7703 cell lines	Losartan	SND1 increases AT1R level by augmenting AT1R mRNA stability; Losartan inhibited both cell migration and invasion	SND1 induces TGF- β expression through AT1R signaling

Cook <i>et al</i> ^[45]	H4-II-E-C3 rat hepatoma cells transfected with Ang II	Losartan Candesartan	Losartan inhibits Angiotensin II-induced proliferation	Both losartan and candesartan are equally effective in competing with Ang II/AT1 receptor interactions on the cell surface
-----------------------------------	---	----------------------	--	--

Ang II: Angiotensin receptor; AT2R: Angiotensin II type 2 receptor; ACE-Is: Angiotensin-converting enzyme inhibitors; ARBs: Angiotensin II type 1 receptor blockers; HCC: Hepatocellular carcinoma; VEGF: Vascular endothelial growth factor; TNF: Tumor necrosis factor; TGF: Transforming growth factor; OLET: Otsuka Long-Evans Tokushima Fatty.

reduced the population of cells in phase S, possibly by reducing the expression of cyclin D1 and cyclin E. Additionally, it was demonstrated that the block of proliferation could be secondary to the induction of AMPK/mTOR signaling in HCC cells. Finally, telmisartan increased the levels of phospho-p53 and decreased the levels of survivin in HLF cells, thus promoting HCC apoptosis, and it also inhibited angiogenesis by increasing bFGF levels in HLF cells as detected by the protein array. In Hep3B and QGY-7703 cells^[44], it was shown that staphylococcal nuclease domain containing-1 (SND1) could play a role in promoting hepatocarcinogenesis and cell migration by triggering TGF- β signaling, a mechanism that was mediated by increasing activation of angiotensin receptor 1. In this experiment, losartan reduced the migration and invasive potential of HCC cells. Cook *et al*^[45] transfected Ang(-S)Exp/pSVL plasmid, which causes the production of a stable, non-secreted form of angiotensinogen, in H4-II-E-C3 rat hepatoma cells. Thus, they discovered that this novel cell clone exhibited enhanced proliferation, which was blocked by losartan, but not by candesartan, although they were equally effective in competing with Ang II/AT1 receptor interactions on the cell surface. Furthermore, the addition of phenylarsine oxide, a molecule that blocks angiotensin receptor internalization, abolished the effect of losartan. Therefore, the authors concluded that losartan could be effective mainly on intranuclear angiotensin receptor.

DISCUSSION

The interest in the potential beneficial effects of RAS inhibitors against HCC started about two decades ago. Since then, several *in vitro* studies, using either different tumor cell lines or cell lines manipulated through adenoviral vectors, and numerous *in vivo* models have been used to demonstrate the antitumoral effect of RAS inhibitors on hepatocarcinoma development. We had to wait about 10 years for the first clinical study on the possible role of ACE-Is administration in the recurrence of HCC^[13]. This study, designed as a randomized trial, was followed within a couple of years by two similar randomized trials from the same authors, all demonstrating the efficacy of the ACE-I perindopril in reducing tumor recurrence when used in combination with other drugs^[13-15]. Interestingly, this effect was associated with a significant reduction of VEGF^[13-15] and alpha-fetoprotein^[13,14] serum levels. However, a significant effect on survival was not observed in any of these studies. The latter result requires some comments: only the first of these three studies reported a survival analysis (reported as "data not shown" in the other studies), which was interrupted at 42 mo, although the survival rate in the control group and the group receiving a combined treatment was 84% and 96%, respectively^[13]. In consideration of this trend, we do not really know whether a longer period of follow-up would have led to significant survival differences. Moreover, it is plausible that a delayed recurrence in patients prevalently in Child A class (75%-87.5%) could also prolong their survival.

The more recent human studies are all observational retrospective cohort or case-control studies^[7,16-21]. All these studies evaluated the effect of ACEIs and/or ARBs on overall survival. Among them, only one took into consideration tumor recurrence whereas another evaluated disease-free-survival^[20,21]. With regard to tumor recurrence, the study of Facciorusso *et al*^[20] was performed on patients with characteristics very similar to those reported in the randomized trials (patients with a low number and small-size tumors susceptible to RFA treatment). In addition, only patients undergoing antihypertensive therapy for at least two years before RFA were included to allow a reasonable induction period for the antiangiogenic and antifibrotic effects of such drugs. This study demonstrated a significant beneficial effect of ARBs, not only on time to recurrence but also on the overall survival after a period of observation of 96 months, which is twice the one reported in the randomized trials.

The study on disease-free survival was focused on metabolic factors, including

obesity, diabetes, and hypertension, on patients undergoing HCC resection^[21]. In this case, patients had a more advanced tumor disease, as demonstrated by the presence of pulmonary and bone metastasis at the follow-up, and underwent a liver resection including either one segment, more than one segment or even hemihepatectomy, plus removal of additional contiguous segments. In this study, neither overall survival nor disease-survival was improved by the anti-hypertensive treatment. However, the authors underline the fact that patients receiving angiotensin II-blocking agents had a better disease-free and overall survival rate than hypertensive patients not treated with angiotensin II-blocking agents.

The other observational studies conducted only on overall survival rates did not produce conclusive results since the treatment with RAS inhibitors was beneficial in one study^[19], did not influence survival in two studies^[17,18], and resulted either ineffective or even associated with an increased HCC risk in non-cirrhotic HCV patients in another study^[16].

Among the possible explanations for these discrepancies, the different time of exposure to antihypertensive drugs assumes particular importance if we consider that RAS inhibitors would require a reasonable induction period to counteract the carcinogenic process through their antiangiogenic effects. In the various studies, the antihypertensive treatment preceded for a variable period the first recorded liver cancer diagnosis or started at the time of HCC diagnosis, and the comparison of exposure during the follow up was difficult. Other explanations are the characteristics of the tumor (susceptible to surgical resection or locoregional therapy or advanced), the different stages of the underlying liver disease, and the concomitant pharmacological therapies/comorbidities potentially influencing tumor biology.

All experimental studies lead to concordant conclusions on the beneficial effects of RAS inhibitors on hepatocarcinogenesis, with some difference within the same drug class. However, it should be highlighted that this beneficial effect was mainly based on the demonstration that ACE-Is or ARBs inhibit the process of angiogenesis, the most studied and extensively exploited biological mechanism. The main aspects explored included: the involvement of VEGF serum levels or VEGF protein or mRNA expression^[25,28-30,33-35,37-42] in endothelial cell growth and differentiation^[46]; the contribution of HIF-1^[28,40] as a transcription factor for VEGF^[47]; the use of CD31^[29-33,42] [also known as platelet endothelial cell adhesion molecule 1 (PECAM-1)] to recognize endothelial cells^[48]; endothelial cell proliferation and tubular formation *in vitro* and/or *in vivo*^[30,32,34,37,39,41,42], a multi-step process triggered by VEGF^[49].

Inhibition of tumor cells proliferation and induction of apoptosis are two mechanisms that have also been studied to explain how RAS inhibitors could counteract hepatocarcinogenesis. Some authors report an inhibitory effect of RAS inhibitors on tumor cell proliferation^[25] or describe a block of proliferation secondary to the induction of AMPK/mTOR signaling^[43], all studies performed by Yoshiji *et al.*^[30-33] and by others^[35,37] do not confirm these findings, and one study even describes a pro mitogenic effect of angiotensin II^[45]. Moreover, a complete overview of this aspect requires a better definition of the role of VEGF as a potential autocrine growth factor on tumor cells^[46] and the relationship between angiotensin receptor 1 (AT1R) and angiotensin receptor 2 (AT2R)^[36,50,51].

More consistent data seem to demonstrate the proapoptotic effect of RAS inhibitors on hepatic cancer cells^[31,33,36,37,43].

Finally, the last possible antitumor mechanism related to the use of RAS inhibitors could consist of the suppression of inflammation by the reduction of TNF- α ^[25,40], TGF- β ^[25,29,40,41] and MMP-2^[25,29], and an increase of tissue inhibitor of metalloproteinase (TIMP)-1, a negative regulator of MMP2^[29].

All mechanisms mentioned above supporting the anti-tumor activity of RAS inhibitors were studied in HCC models, mostly induced by the carcinogenic agent DEN or in rodents with xenotransplanted tumors, except in two studies^[40,41] that used choline-deficient/low-methionine diets to induce steatohepatitis related HCC. Viral hepatitis still represents the principal cause of HCC in humans. The demonstration of a similar protective effect of ACE-I or ARBs in an animal model that mimics the progression of viral-related liver disease in humans, such as the HBV transgenic mouse model^[52], would strongly support previous experimental data. In addition, since ACE-Is have been able to produce a prophylactic effect on hepatocarcinogenesis^[29,32,33], this model could be used to test the RAS inhibitor influence on the expression of genes that are involved early on in the carcinogenesis^[53]. In fact, the adoption of transgenic mouse models is widely used to test the effect of drugs or nutrients to counteract the process of carcinogenesis^[54]. Finally, to clarify if RAS inhibitors could influence the progression toward overt HCC in humans, it is necessary to design randomized, controlled trials in which the treatment with these anti-hypertensive drugs is initiated in patients at high risk of HCC. Cirrhotics with a low number (≤ 3) of small tumors (≤ 3 cm) or a single larger tumor (≤ 5 cm), with a

complete tumor ablation would be the ideal patients for this study considering that HCC recurrence at 3 years is observed in 48%-53% and 28%-33% in the presence of HCV infection or after HCV eradication, respectively^[55,56].

In conclusion, all experimental data strongly support the ability of RAS inhibitors to counteract HCC development. In humans, this protective effect was confirmed when RAS inhibitors were used, prevalently in combination with other drugs, to prevent HCC recurrence. On the other hand, even if the results of RAS inhibitors on patient survival seem to be inconsistent, these findings are less conclusive. Further studies are necessary to evaluate whether their prolonged use could influence the progression of cirrhosis toward HCC.

ARTICLE HIGHLIGHTS

Research background

Hepatocellular carcinoma (HCC) represents about 90% of primary liver cancers and usually develops in patients with underlying liver cirrhosis. One of the major cancer hallmarks of HCC is the process of neoangiogenesis, which is thought to occur early in the carcinogenetic process. Theoretically, angiotensin-converting enzyme inhibitors (ACE-Is) and angiotensin II type 1 receptor blockers (ARBs) could have a role in HCC primary prevention since they reduce both neovascularization and the development of liver fibrosis.

Research motivation

The possible role of renin-angiotensin system (RAS) inhibitors in the prevention of HCC is still an attractive field of research, as suggested by the recently published experimental and clinical studies.

Research objectives

To elucidate the role of ARBs and ACE-Is in HCC.

Research methods

Literature was selected in August 2018, focusing on studies regarding ACE-Is, ARBs and HCCs. For this purpose, an electronic search of the literature was conducted using the online databases PubMed, Cochrane library, Scopus and Web of Science. Eligibility criteria were: (1) prospective or retrospective clinical studies; (2) epidemiological studies; and (3) experimental studies conducted *in vivo* or *in vitro*. We conducted the articles selection using a two-step approach, to evaluate whether they matched the eligibility criteria.

Research results

Thirty-one studies were selected. Three interventional studies showed that ACE-Is had a significant protective effect on HCC recurrence only when used in combination with vitamin K or branched chain aminoacids, without a significant increase of overall survival. Among six retrospective observational studies, mainly focused on overall survival, only one demonstrated a prolonged survival in ACE-Is group, whereas the two of them that also evaluated tumor recurrence showed conflicting results. All experimental studies displayed beneficial effects of RAS inhibitors on hepatocarcinogenesis. Numerous experimental studies, conducted either in animals and cell cultures, demonstrated the anti-angiogenetic and anti-fibrotic effect of ACE-Is and ARBs, thanks to the suppression of some cytokines such as vascular endothelial growth factor, hypoxia-inducible factor-1 α , transforming growth factor-beta and tumor necrosis factor alpha. All or part of these mechanisms were demonstrated in rodents developing less HCC and preneoplastic lesions after receiving such drugs.

Research conclusions

Our systematic analysis of the literature allowed us to partially reinterpret the data supporting the ability of RAS inhibitors to counteract HCC. While there is reasonable research supporting their ability to reduce the HCC recurrence rate, we think that the conclusions of the randomized clinical trials on overall survival underestimated the potential use of these compounds. These conclusions are further supported by the solid experimental data on the anti-angiogenetic activity of RAS inhibitors, which all suggest their clinical use in the prevention of HCC development.

Research perspectives

We also proposed enlarging the future research on the relationship between RAS inhibitors and HCC to other possible anti carcinogenetic mechanisms.

REFERENCES

- 1 European Association for the Study of the Liver. EASL Clinical Practice Guidelines: Management of hepatocellular carcinoma. *J Hepatol* 2018; **69**: 182-236 [PMID: 29628281 DOI: 10.1016/j.jhep.2018.03.019]
- 2 Torre LA, Bray F, Siegel RL, Ferlay J, Lortet-Tieulent J, Jemal A. Global cancer statistics, 2012. *CA*

- Cancer J Clin* 2015; **65**: 87-108 [PMID: 25651787 DOI: 10.3322/caac.21262]
- 3 **Global Burden of Disease Liver Cancer Collaboration**; Akinyemiju T, Abera S, Ahmed M, Alam N, Alemayohu MA, Allen C, Al-Raddadi R, Alvis-Guzman N, Amoako Y, Artaman A, Ayele TA, Barac A, Bensenor I, Berhane A, Bhutta Z, Castillo-Rivas J, Chitheer A, Choi JY, Cowie B, Dandona L, Dandona R, Dey S, Dicker D, Phuc H, Ekwueme DU, Zaki MS, Fischer F, Fürst T, Hancock J, Hay SI, Hotez P, Jee SH, Kasaeian A, Khader Y, Khang YH, Kumar A, Kutz M, Larson H, Lopez A, Lunevicius R, Malekzadeh R, McAlinden C, Meier T, Mendoza W, Mokdad A, Nagel G, Nguyen Q, Nguyen G, Ogo F, Patton G, Pereira DM, Pourmalek F, Qorbani M, Radfar A, Roshandel G, Salomon JA, Sanabria J, Sartorius B, Satpathy M, Sawhney M, Sepanlou S, Shackelford K, Shore H, Sun J, Mengistu DT, Tran B, Ukwaja KN, Vlassov V, Vollset SE, Vos T, Wakayo T, Weiderpass E, Werdecker A, Yonemoto N, Younis M, Yu C, Zaidi Z, Zhu L, Murray CJL, Naghavi M, Fitzmaurice C, Moradi-Lakeh M, Topór-Mądry R. The Burden of Primary Liver Cancer and Underlying Etiologies From 1990 to 2015 at the Global, Regional, and National Level: Results From the Global Burden of Disease Study 2015. *JAMA Oncol* 2017; **3**: 1683-1691 [PMID: 28983565 DOI: 10.1001/jamaoncol.2017.3055]
 - 4 **Yüksel Ş**, Boylu Akyerli C, Cengiz Yakicier M. Angiogenesis, Invasion, and Metastasis Characteristics of Hepatocellular Carcinoma. *J Gastrointest Cancer* 2017 [PMID: 28785955 DOI: 10.1007/s12029-017-9962-5]
 - 5 **Dhanasekaran R**, Venkatesh SK, Torbenson MS, Roberts LR. Clinical implications of basic research in hepatocellular carcinoma. *J Hepatol* 2016; **64**: 736-745 [PMID: 26450813 DOI: 10.1016/j.jhep.2015.09.008]
 - 6 **Berretta M**, Rinaldi L, Di Benedetto F, Lleshi A, De Re V, Facchini G, De Paoli P, Di Francia R. Angiogenesis Inhibitors for the Treatment of Hepatocellular Carcinoma. *Front Pharmacol* 2016; **7**: 428 [PMID: 27881963 DOI: 10.3389/fphar.2016.00428]
 - 7 **Facciorusso A**, Licinio R, Carr BI, Di Leo A, Barone M. MEK 1/2 inhibitors in the treatment of hepatocellular carcinoma. *Expert Rev Gastroenterol Hepatol* 2015; **9**: 993-1003 [PMID: 25915713 DOI: 10.1586/17474124.2015.1040763]
 - 8 **Raoul JL**, Gilibert M, Adhoute X, Edeline J. An in-depth review of chemical angiogenesis inhibitors for treating hepatocellular carcinoma. *Expert Opin Pharmacother* 2017; **18**: 1467-1476 [PMID: 28893090 DOI: 10.1080/14656566.2017.1378346]
 - 9 **Corey KE**, Shah N, Misdraji J, Abu Dayyeh BK, Zheng H, Bhan AK, Chung RT. The effect of angiotensin-blocking agents on liver fibrosis in patients with hepatitis C. *Liver Int* 2009; **29**: 748-753 [PMID: 19220742 DOI: 10.1111/j.1478-3231.2009.01973.x]
 - 10 **Kaji K**, Yoshiji H, Ikenaka Y, Noguchi R, Aihara Y, Shirai Y, Douhara A, Fukui H. Possible involvement of angiogenesis in chronic liver diseases: interaction among renin-angiotensin-aldosterone system, insulin resistance and oxidative stress. *Curr Med Chem* 2012; **19**: 1889-1898 [PMID: 22376037]
 - 11 **Paizis G**, Gilbert RE, Cooper ME, Murthi P, Schembri JM, Wu LL, Rumble JR, Kelly DJ, Tikellis C, Cox A, Smallwood RA, Angus PW. Effect of angiotensin II type 1 receptor blockade on experimental hepatic fibrogenesis. *J Hepatol* 2001; **35**: 376-385 [PMID: 11592599]
 - 12 **Song T**, Choi CH, Kim MK, Kim ML, Yun BS, Seong SJ. The effect of angiotensin system inhibitors (angiotensin-converting enzyme inhibitors or angiotensin receptor blockers) on cancer recurrence and survival: a meta-analysis. *Eur J Cancer Prev* 2017; **26**: 78-85 [PMID: 27158979 DOI: 10.1097/CEJ.0000000000000269]
 - 13 **Yoshiji H**, Noguchi R, Toyohara M, Ikenaka Y, Kitade M, Kaji K, Yamazaki M, Yamao J, Mito A, Sawai M, Yoshida M, Fujimoto M, Tsujimoto T, Kawaratani H, Uemura M, Fukui H. Combination of vitamin K2 and angiotensin-converting enzyme inhibitor ameliorates cumulative recurrence of hepatocellular carcinoma. *J Hepatol* 2009; **51**: 315-321 [PMID: 19501932 DOI: 10.1016/j.jhep.2009.04.011]
 - 14 **Yoshiji H**, Noguchi R, Ikenaka Y, Kaji K, Aihara Y, Yamazaki M, Yamao J, Toyohara M, Mito A, Sawai M, Yoshida M, Morioka K, Fujimoto M, Uemura M, Fukui H. Combination of branched-chain amino acids and angiotensin-converting enzyme inhibitor suppresses the cumulative recurrence of hepatocellular carcinoma: a randomized control trial. *Oncol Rep* 2011; **26**: 1547-1553 [PMID: 21874260 DOI: 10.3892/or.2011.1433]
 - 15 **Yoshiji H**, Noguchi R, Ikenaka Y, Kaji K, Shirai Y, Aihara Y, Yamao J, Toyohara M, Mito A, Sawai M, Yoshida M, Morioka K, Fujimoto M, Uemura M, Kawaratani H, Tsujimoto T, Fukui H. Soluble VEGF receptor-2 may be a predictive marker of anti-angiogenic therapy with clinically available safe agents. *Oncol Lett* 2011; **2**: 69-73 [PMID: 22870131 DOI: 10.3892/ol.2010.196]
 - 16 **Ho CM**, Lee CH, Lee MC, Zhang JF, Wang JY, Hu RH, Lee PH. Comparative effectiveness of angiotensin-converting enzyme inhibitors and angiotensin II receptor blockers in chemoprevention of hepatocellular carcinoma: a nationwide high-risk cohort study. *BMC Cancer* 2018; **18**: 401 [PMID: 29631561 DOI: 10.1186/s12885-018-4292-y]
 - 17 **Hagberg KW**, Sahasrabudhe VV, McGlynn KA, Jick SS. Does Angiotensin-Converting Enzyme Inhibitor and β -Blocker Use Reduce the Risk of Primary Liver Cancer? A Case-Control Study Using the U.K. Clinical Practice Research Datalink. *Pharmacotherapy* 2016; **36**: 187-195 [PMID: 26846893 DOI: 10.1002/phar.1704]
 - 18 **Walker AJ**, West J, Grainge MJ, Card TR. Angiotensin converting enzyme inhibitors and hepatocellular carcinoma incidence in the General Practice Research Database. *Cancer Causes Control* 2011; **22**: 1743-1747 [PMID: 21909951 DOI: 10.1007/s10552-011-9837-1]
 - 19 **Pinter M**, Weinmann A, Wörns MA, Hucke F, Bota S, Marquardt JU, Duda DG, Jain RK, Galle PR, Trauner M, Peck-Radosavljevic M, Sieghart W. Use of inhibitors of the renin-angiotensin system is associated with longer survival in patients with hepatocellular carcinoma. *United European Gastroenterol J* 2017; **5**: 987-996 [PMID: 29163965 DOI: 10.1177/2050640617695698]
 - 20 **Facciorusso A**, Del Prete V, Crucinio N, Muscatiello N, Carr BI, Di Leo A, Barone M. Angiotensin receptor blockers improve survival outcomes after radiofrequency ablation in hepatocarcinoma patients. *J Gastroenterol Hepatol* 2015; **30**: 1643-1650 [PMID: 25974743 DOI: 10.1111/jgh.12988]
 - 21 **Kaibori M**, Ishizaki M, Matsui K, Kitade H, Matsui Y, Kwon AH. Evaluation of metabolic factors on the prognosis of patients undergoing resection of hepatocellular carcinoma. *J Gastroenterol Hepatol* 2011; **26**: 536-543 [PMID: 21332549 DOI: 10.1111/j.1440-1746.2010.06439.x]
 - 22 **Tsan YT**, Lee CH, Wang JD, Chen PC. Statins and the risk of hepatocellular carcinoma in patients with hepatitis B virus infection. *J Clin Oncol* 2012; **30**: 623-630 [PMID: 22271485 DOI: 10.1200/JCO.2011.36.0917]
 - 23 **Singh PP**, Singh S. Statins and risk reduction in hepatocellular carcinoma: fact or fiction? *J Clin Oncol*

- 2012; **30**: 2569-2570; author reply 2570-2571 [PMID: [22689810](#) DOI: [10.1200/JCO.2012.43.2179](#)]
- 24 **Yoshiji H**, Noguchi R, Yamazaki M, Ikenaka Y, Sawai M, Ishikawa M, Kawaratani H, Mashitani T, Kitade M, Kaji K, Uemura M, Yamao J, Fujimoto M, Mitoro A, Toyohara M, Yoshida M, Fukui H. Combined treatment of vitamin K2 and angiotensin-converting enzyme inhibitor ameliorates hepatic dysplastic nodule in a patient with liver cirrhosis. *World J Gastroenterol* 2007; **13**: 3259-3261 [PMID: [17589909](#)]
- 25 **Saber S**, Mahmoud AAA, Goda R, Helal NS, El-Ahwany E, Abdelghany RH. Perindopril, fosinopril and losartan inhibited the progression of diethylnitrosamine-induced hepatocellular carcinoma in mice via the inactivation of nuclear transcription factor kappa-B. *Toxicol Lett* 2018; **295**: 32-40 [PMID: [29859236](#) DOI: [10.1016/j.toxlet.2018.05.036](#)]
- 26 **Saber S**, Mahmoud A, Helal N, El-Ahwany E, Abdelghany R. Liver Protective Effects of Renin-Angiotensin System Inhibition Have No Survival Benefits in Hepatocellular Carcinoma Induced By Repetitive Administration of Diethylnitrosamine in Mice. *Open Access Maced J Med Sci* 2018; **6**: 955-960 [PMID: [29983784](#) DOI: [10.3889/oamjms.2018.167](#)]
- 27 **Nasr M**, Selima E, Hamed O, Kazem A. Targeting different angiogenic pathways with combination of curcumin, leflunomide and perindopril inhibits diethylnitrosamine-induced hepatocellular carcinoma in mice. *Eur J Pharmacol* 2014; **723**: 267-275 [PMID: [24291100](#) DOI: [10.1016/j.ejphar.2013.11.022](#)]
- 28 **Vermeulen PB**, Gasparini G, Fox SB, Toi M, Martin L, McCulloch P, Pezzella F, Viale G, Weidner N, Harris AL, Dirix LY. Quantification of angiogenesis in solid human tumours: an international consensus on the methodology and criteria of evaluation. *Eur J Cancer* 1996; **32A**: 2474-2484 [PMID: [9059336](#)]
- 29 **Mansour MA**, Al-Ismaeel H, Al-Rikabi AC, Al-Shabanah OA. Comparison of angiotensin converting enzyme inhibitors and angiotensin II type 1 receptor blockade for the prevention of premalignant changes in the liver. *Life Sci* 2011; **89**: 188-194 [PMID: [21699905](#) DOI: [10.1016/j.lfs.2011.06.002](#)]
- 30 **Yanase K**, Yoshiji H, Ikenaka Y, Noguchi R, Kitade M, Kaji K, Yoshii J, Namisaki T, Yamazaki M, Asada K, Tsujimoto T, Akahane T, Uemura M, Fukui H. Synergistic inhibition of hepatocellular carcinoma growth and hepatocarcinogenesis by combination of 5-fluorouracil and angiotensin-converting enzyme inhibitor via anti-angiogenic activities. *Oncol Rep* 2007; **17**: 441-446 [PMID: [17203185](#)]
- 31 **Yoshiji H**, Kuriyama S, Noguchi R, Yoshii J, Ikenaka Y, Yanase K, Namisaki T, Kitade M, Yamazaki M, Akahane T, Asada K, Tsujimoto T, Uemura M, Fukui H. Amelioration of carcinogenesis and tumor growth in the rat liver by combination of vitamin K2 and angiotensin-converting enzyme inhibitor via anti-angiogenic activities. *Oncol Rep* 2006; **15**: 155-159 [PMID: [16328049](#)]
- 32 **Yoshiji H**, Kuriyama S, Noguchi R, Yoshii J, Ikenaka Y, Yanase K, Namisaki T, Kitade M, Yamazaki M, Masaki T, Fukui H. Combination of vitamin K2 and the angiotensin-converting enzyme inhibitor, perindopril, attenuates the liver enzyme-altered preneoplastic lesions in rats via angiogenesis suppression. *J Hepatol* 2005; **42**: 687-693 [PMID: [15826718](#) DOI: [10.1016/j.jhep.2004.12.025](#)]
- 33 **Yoshiji H**, Noguchi R, Kuriyama S, Yoshii J, Ikenaka Y. Combination of interferon and angiotensin-converting enzyme inhibitor, perindopril, suppresses liver carcinogenesis and angiogenesis in mice. *Oncol Rep* 2005; **13**: 491-495 [PMID: [15706423](#)]
- 34 **Yoshiji H**, Noguchi R, Kuriyama S, Yoshii J, Ikenaka Y, Yanase K, Namisaki T, Kitade M, Yamazaki M, Uemura M, Fukui H. Suppression of renin-angiotensin system attenuates hepatocarcinogenesis via angiogenesis inhibition in rats. *Anticancer Res* 2005; **25**: 3335-3340 [PMID: [16101147](#)]
- 35 **Fan F**, Tian C, Tao L, Wu H, Liu Z, Shen C, Jiang G, Lu Y. Candesartan attenuates angiogenesis in hepatocellular carcinoma via downregulating AT1R/VEGF pathway. *Biomed Pharmacother* 2016; **83**: 704-711 [PMID: [27470571](#) DOI: [10.1016/j.biopha.2016.07.039](#)]
- 36 **Du H**, Liang Z, Zhang Y, Jie F, Li J, Fei Y, Huang Z, Pei N, Wang S, Li A, Chen B, Zhang Y, Summers C, Li M, Li H. Effects of angiotensin II type 2 receptor overexpression on the growth of hepatocellular carcinoma cells in vitro and in vivo. *PLoS One* 2013; **8**: e83754 [PMID: [24391821](#) DOI: [10.1371/journal.pone.0083754](#)]
- 37 **Noguchi R**, Yoshiji H, Kuriyama S, Yoshii J, Ikenaka Y, Yanase K, Namisaki T, Kitade M, Yamazaki M, Mitoro A, Tsujinoue H, Imazu H, Masaki T, Fukui H. Combination of interferon-beta and the angiotensin-converting enzyme inhibitor, perindopril, attenuates murine hepatocellular carcinoma development and angiogenesis. *Clin Cancer Res* 2003; **9**: 6038-6045 [PMID: [14676130](#)]
- 38 **Yoshiji H**, Yoshii J, Ikenaka Y, Noguchi R, Yanase K, Tsujinoue H, Imazu H, Fukui H. Suppression of the renin-angiotensin system attenuates vascular endothelial growth factor-mediated tumor development and angiogenesis in murine hepatocellular carcinoma cells. *Int J Oncol* 2002; **20**: 1227-1231 [PMID: [12012003](#)]
- 39 **Yoshiji H**, Kuriyama S, Kawata M, Yoshii J, Ikenaka Y, Noguchi R, Nakatani T, Tsujinoue H, Fukui H. The angiotensin-I-converting enzyme inhibitor perindopril suppresses tumor growth and angiogenesis: possible role of the vascular endothelial growth factor. *Clin Cancer Res* 2001; **7**: 1073-1078 [PMID: [11309359](#)]
- 40 **Tamaki Y**, Nakade Y, Yamauchi T, Makino Y, Yokohama S, Okada M, Aso K, Kanamori H, Ohashi T, Sato K, Nakao H, Haneda M, Yoneda M. Angiotensin II type 1 receptor antagonist prevents hepatic carcinoma in rats with nonalcoholic steatohepatitis. *J Gastroenterol* 2013; **48**: 491-503 [PMID: [22886508](#) DOI: [10.1007/s00535-012-0651-7](#)]
- 41 **Noguchi R**, Yoshiji H, Ikenaka Y, Kaji K, Aihara Y, Shirai Y, Namisaki T, Kitade M, Douhara A, Moriya K, Fukui H. Dual blockade of angiotensin-II and aldosterone suppresses the progression of a non-diabetic rat model of steatohepatitis. *Hepatol Res* 2013; **43**: 765-774 [PMID: [23163573](#) DOI: [10.1111/hepr.12008](#)]
- 42 **Yoshiji H**, Noguchi R, Kaji K, Ikenaka Y, Shirai Y, Namisaki T, Kitade M, Tsujimoto T, Kawaratani H, Fukui H. Attenuation of insulin-resistance-based hepatocarcinogenesis and angiogenesis by combined treatment with branched-chain amino acids and angiotensin-converting enzyme inhibitor in obese diabetic rats. *J Gastroenterol* 2010; **45**: 443-450 [PMID: [19941011](#) DOI: [10.1007/s00535-009-0158-z](#)]
- 43 **Oura K**, Tadokoro T, Fujihara S, Morishita A, Chiyo T, Samukawa E, Yamana Y, Fujita K, Sakamoto T, Nomura T, Yoneyama H, Kobara H, Mori H, Iwama H, Okano K, Suzuki Y, Masaki T. Telmisartan inhibits hepatocellular carcinoma cell proliferation in vitro by inducing cell cycle arrest. *Oncol Rep* 2017; **38**: 2825-2835 [PMID: [29048654](#) DOI: [10.3892/or.2017.5977](#)]
- 44 **Santhekadur PK**, Akiel M, Emdad L, Gredler R, Srivastava J, Rajasekaran D, Robertson CL, Mukhopadhyay ND, Fisher PB, Sarkar D. Staphylococcal nuclease domain containing-1 (SND1) promotes migration and invasion via angiotensin II type 1 receptor (AT1R) and TGFβ signaling. *FEBS Open Bio* 2014; **4**: 353-361 [PMID: [24918049](#) DOI: [10.1016/j.fob.2014.03.012](#)]
- 45 **Cook JL**, Zhang Z, Re RN. In vitro evidence for an intracellular site of angiotensin action. *Circ Res* 2001; **89**: 1138-1146 [PMID: [11739278](#)]

- 46 **Byrne AM**, Bouchier-Hayes DJ, Harmey JH. Angiogenic and cell survival functions of vascular endothelial growth factor (VEGF). *J Cell Mol Med* 2005; **9**: 777-794 [PMID: 16364190]
- 47 **Forsythe JA**, Jiang BH, Iyer NV, Agani F, Leung SW, Koos RD, Semenza GL. Activation of vascular endothelial growth factor gene transcription by hypoxia-inducible factor 1. *Mol Cell Biol* 1996; **16**: 4604-4613 [PMID: 8756616]
- 48 **Pusztaszeri MP**, Seelentag W, Bosman FT. Immunohistochemical expression of endothelial markers CD31, CD34, von Willebrand factor, and Flt-1 in normal human tissues. *J Histochem Cytochem* 2006; **54**: 385-395 [PMID: 16234507 DOI: 10.1369/jhc.4A6514.2005]
- 49 **Lee H**, Kang KT. Advanced tube formation assay using human endothelial colony forming cells for *in vitro* evaluation of angiogenesis. *Korean J Physiol Pharmacol* 2018; **22**: 705-712 [PMID: 30402031 DOI: 10.4196/kjpp.2018.22.6.705]
- 50 **Ozeki K**, Tanida S, Morimoto C, Inoue Y, Mizoshita T, Tsukamoto H, Shimura T, Kataoka H, Kamiya T, Nishiwaki E, Ishiguro H, Higashiyama S, Joh T. Telmisartan inhibits cell proliferation by blocking nuclear translocation of ProHB-EGF C-terminal fragment in colon cancer cells. *PLoS One* 2013; **8**: e56770 [PMID: 23451083 DOI: 10.1371/journal.pone.0056770]
- 51 **Deshayes F**, Nahmias C. Angiotensin receptors: a new role in cancer? *Trends Endocrinol Metab* 2005; **16**: 293-299 [PMID: 16061390 DOI: 10.1016/j.tem.2005.07.009]
- 52 **Barone M**, Scavo MP, Maiorano E, Di Leo A, Francavilla A. Bone marrow-derived stem cells and hepatocarcinogenesis in hepatitis B virus transgenic mice. *Dig Liver Dis* 2014; **46**: 243-250 [PMID: 24286758 DOI: 10.1016/j.dld.2013.10.008]
- 53 **Barone M**, Spano D, D'Apollito M, Centra M, Lasalandra C, Capasso M, Di Leo A, Volinia S, Arcelli D, Rosso N, Francavilla A, Tiribelli C, Iolascon A. Gene expression analysis in HBV transgenic mouse liver: a model to study early events related to hepatocarcinogenesis. *Mol Med* 2006; **12**: 115-123 [PMID: 16953557 DOI: 10.2119/2006-00015.Barone]
- 54 **Barone M**, Notarnicola M, Caruso MG, Scavo MP, Viggiani MT, Tutino V, Polimeno L, Pesetti B, Di Leo A, Francavilla A. Olive oil and omega-3 polyunsaturated fatty acids suppress intestinal polyp growth by modulating the apoptotic process in ApcMin/+ mice. *Carcinogenesis* 2014; **35**: 1613-1619 [PMID: 24632492 DOI: 10.1093/carcin/bgu068]
- 55 **Ng KKC**, Chok KSH, Chan ACY, Cheung TT, Wong TCL, Fung JYY, Yuen J, Poon RTP, Fan ST, Lo CM. Randomized clinical trial of hepatic resection versus radiofrequency ablation for early-stage hepatocellular carcinoma. *Br J Surg* 2017; **104**: 1775-1784 [PMID: 29091283 DOI: 10.1002/bjs.10677]
- 56 **Petta S**, Cabibbo G, Barbara M, Attardo S, Bucci L, Farinati F, Giannini EG, Tovoli F, Ciccarese F, Rapaccini GL, Di Marco M, Caturelli E, Zoli M, Borzio F, Sacco R, Virdone R, Marra F, Felder M, Morisco F, Benvegnù L, Gasbarrini A, Svegliati-Baroni G, Foschi FG, Olivani A, Masotto A, Nardone G, Colecchia A, Persico M, Boccaccio V, Craxi A, Bruno S, Trevisani F, Cammà C; Italian Liver Cancer (ITA. LI.CA) Group. Hepatocellular carcinoma recurrence in patients with curative resection or ablation: impact of HCV eradication does not depend on the use of interferon. *Aliment Pharmacol Ther* 2017; **45**: 160-168 [PMID: 27790734 DOI: 10.1111/apt.13821]



Published By Baishideng Publishing Group Inc
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA
Telephone: +1-925-2238242
Fax: +1-925-2238243
E-mail: bpgoffice@wjgnet.com
Help Desk: <http://www.f6publishing.com/helpdesk>
<http://www.wjgnet.com>

